

## **Assessing sustainable biophysical human-nature connectedness at regional scales**

Dorninger, Christian; Abson, David J.; Fischer, Joern; von Wehrden, Henrik

*Published in:*  
Environmental Research Letters

*DOI:*  
[10.1088/1748-9326/aa68a5](https://doi.org/10.1088/1748-9326/aa68a5)

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication](#)

*Citation for published version (APA):*  
Dorninger, C., Abson, D. J., Fischer, J., & von Wehrden, H. (2017). Assessing sustainable biophysical human-nature connectedness at regional scales. *Environmental Research Letters*, 12(5), 1-11. Article 055001. <https://doi.org/10.1088/1748-9326/aa68a5>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LETTER • OPEN ACCESS

# Assessing sustainable biophysical human–nature connectedness at regional scales

To cite this article: Christian Dörninger *et al* 2017 *Environ. Res. Lett.* **12** 055001

View the [article online](#) for updates and enhancements.

## Related content

- [The surprisingly small but increasing role of international agricultural trade on the European Union's dependence on mineral phosphorus fertiliser](#)  
Thomas Nesme, Solène Roques, Geneviève S Metson *et al*.
- [Mapping and analysing cropland use intensity from a NPP perspective](#)  
Maria Niedertscheider, Thomas Kastner, Tamara Fetzel *et al*.
- [Environmental impacts of food trade via resource use and greenhouse gas emissions](#)  
Carole Dalin and Ignacio Rodríguez-Iturbe

## Recent citations

- [Beyond Biodiversity Conservation: Land Sharing Constitutes Sustainable Agriculture in European Cultural Landscapes](#)  
Jacqueline Loos and Henrik von Wehrden
- [Reconnecting with nature for sustainability](#)  
Christopher D. Ives *et al*
- [Dimensions of Landscape Stewardship across Europe: Landscape Values, Place Attachment, Awareness, and Personal Responsibility](#)  
Maria García-Martín *et al*



Are you our new  
project leader  
marine technology  
development?



MORE INFO? VISIT [WORKINGATNIOZ.NL](#)

# Environmental Research Letters



## LETTER

### OPEN ACCESS

#### RECEIVED

30 November 2016

#### REVISED

20 March 2017

#### ACCEPTED FOR PUBLICATION

23 March 2017

#### PUBLISHED

24 April 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



## Assessing sustainable biophysical human–nature connectedness at regional scales

Christian Dorninger<sup>1</sup>, David J Abson, Joern Fischer and Henrik von Wehrden

Faculty of Sustainability, Leuphana University Lüneburg, D-21335 Lüneburg, Germany

<sup>1</sup> Author to whom any correspondence should be addressed.

E-mail: [christian.dorninger@leuphana.de](mailto:christian.dorninger@leuphana.de)

**Keywords:** biosphere, embodied energy, HANPP, land use, sustainability, teleconnections

### Abstract

Humans are biophysically connected to the biosphere through the flows of materials and energy appropriated from ecosystems. While this connection is fundamental for human well-being, many modern societies have—for better or worse—disconnected themselves from the natural productivity of their immediate regional environment. In this paper, we conceptualize the biophysical human–nature connectedness of land use systems at regional scales. We distinguish two mechanisms by which primordial connectedness of people to regional ecosystems has been circumvented via the use of external inputs. First, ‘biospheric disconnection’ refers to people drawing on non-renewable minerals from outside the biosphere (e.g. fossils, metals and other minerals). Second, ‘spatial disconnection’ arises from the imports and exports of biomass products and imported mineral resources used to extract and process ecological goods. Both mechanisms allow for greater regional resource use than would be possible otherwise, but both pose challenges for sustainability, for example, through waste generation, depletion of non-renewable resources and environmental burden shifting to distant regions. In contrast, biophysically reconnected land use systems may provide renewed opportunities for inhabitants to develop an awareness of their impacts and fundamental reliance on ecosystems. To better understand the causes, consequences, and possible remedies related to biophysical disconnectedness, new quantitative methods to assess the extent of regional biophysical human–nature connectedness are needed. To this end, we propose a new methodological framework that can be applied to assess biophysical human–nature connectedness in any region of the world.

### 1. Introduction

Human societies are inherently connected to and dependent on the biosphere and its functions (Boulding 1966, Daily 1997, Folke *et al* 2011) through the flow of materials and energy (Haberl *et al* 2014, Cooke *et al* 2016). However, modern societies have increasingly disconnected themselves from their immediate regional environment by accessing material and energy flows from distant places (Kastner *et al* 2014, Bergmann and Holmberg 2016) and from outside the biosphere (Wiedmann *et al* 2015). Industry, technology and long-distance trade have enabled a disconnect of human activities from the primary production of their regional environment (Yu *et al* 2013), and from the biosphere by relying on

industrial mineral resources (i.e. fossils, metals, and other minerals extracted from the lithosphere (Cumming *et al* 2014)). Hence, despite growing calls for societal reconnection to the biosphere (Folke *et al* 2011, Andersson *et al* 2014, Folke *et al* 2016), what this means from a biophysical perspective remains poorly understood.

The notion of biophysical human–nature connectedness is in conflict with the notion of decoupling socio-economic activities from natural resource use. In parallel to growing calls to ‘reconnect’ to the biosphere, other scholars have noted a relative decoupling of material throughput and economic growth for some regions (e.g. Fischer-Kowalski and Swilling 2011). Nevertheless, the economy is embedded in the environment (Martínez-Alier and

Muradian 2015, Folke *et al* 2016), all resources used in the economy are drawn from the environment, and all waste must return to the environment. Therefore, complete and sustained decoupling of economic activities from the environment is, by definition, untenable. Moreover, Cumming *et al* (2014) argued that a disconnect weakens direct feedbacks between ecosystems and societies, thereby potentially causing overexploitation and collapse. Others have claimed that disconnections provide opportunities for wild nature to be sustained by decoupling human development from environmental impacts (e.g. Asafu-Adjaye *et al* 2015). The purported benefits of purposeful disconnection are premised on intensified, more efficient land use. However, there is considerable debate regarding the efficacy and sustainability of such moves (Loos *et al* 2014a). For example, human–nature disconnectedness can increase inter- and intragenerational injustice by taxing future generations and distant regions (Haberl *et al* 2002, Martinez-Alier *et al* 2014) via overconsumption, depletion of natural resources, and pollution of the environment (Pearson 2007, Wiedmann 2016). In addition, disconnection from the natural environment may foster a systemic cognitive distancing of land use related activities from their environmental impacts (Cumming *et al* 2014, Seppelt and Cumming 2016).

Both perspectives—increasing or decreasing our distance to nature—share the goal of reducing pressure on ecosystems, but with different underpinning assumptions. By increasing our distance to nature we pin our hopes on the ‘efficiency’ of industrial technology in order to ‘spare’ land (Waggoner 1996)—a core idea of ecomodernism (Asafu-Adjaye *et al* 2015). We argue that this view fails to recognize spillover and distal effects, and is largely blind to issues of justice. For this reason, we instead argue for a reconnection of human activities to the biosphere and its regenerative cycles. This, in turn, implies not only a reduction of industrial material use and a limitation of human domination of ecosystems, but also a strengthened sense of being connected with and knowing the limits of nature (Folke *et al* 2011).

To facilitate constructive debate on whether we should reconnect to or disconnect from the biosphere, in this paper, we propose a conceptual framework to analyze regional-scale biophysical human–nature connectedness. The proposed framework builds on the regional land use system as unit of analysis. Yet it explicitly recognizes not only regional land use, but also global material trade and energy flows. By accounting for both economic and biophysical processes, we integrate concepts such as self-sufficiency, land use intensity, resource use, biophysical and embodied trade flows, waste generation, and environmental feedback loops into the framework. Thus, the framework provides a new lens through which land use sustainability can be investigated, which goes beyond ‘on site’ efficiency thinking (Fischer *et al* 2014, von Wehrden

*et al* 2014). Our focus in this paper is primarily conceptual, but we also provide an outlook for how existing methodological approaches can be used to operationalize the proposed framework.

The paper is structured as follows. First, we outline our conceptual model, distinguishing between different types of biophysical disconnection. Second, we provide concrete examples to illustrate how the proposed framework can help to understand the sustainability challenges facing different regions. Finally, we provide a methodological outlook showing avenues how the proposed framework can be operationalized in order to generate quantitatively robust measures of regional-scale biophysical human–nature connectedness.

## 2. Conceptualizing regional biophysical human–nature connectedness

Regional land use systems are an appropriate unit to analyze biophysical human–nature connectedness because (1) energy and material flows across larger extents are typically too heterogeneous to be usefully aggregated; and (2) humans meaningfully experience life at regional scales (Kissinger and Rees 2010, Wu 2013). The spatial boundary of a ‘region’ will most often be defined by sub-national political-administrative units (e.g. from municipalities to federal states), as this is a vital scale for many political decisions (Dearing *et al* 2014) and usually the finest scale at which relevant material and energy flow data is available.

There are multiple ways in which humans’ connectedness to natural ecological productivity can be conceptualized. For example, Seppelt *et al* (2014) suggest a framework based on distinguishing between renewable and non-renewable resource use. However, for regional assessments clear system boundaries are required, therefore, our framework distinguishes between two realms of land use related disconnectedness from the regional biosphere. The first possible realm is ‘biospheric disconnectedness’, and stems from the use of materials external to the biosphere, such as artificial agrochemicals, fossils or machinery. The second possible realm is ‘spatial disconnectedness’, and relates to the appropriation of distal ecological goods to bolster local production via imports of biomass, including food, timber, or feed for livestock. Moreover, one could consider the import of mineral resources used to extract and process ecological goods in the region as an additional form of spatial disconnect.

Both biospheric and spatial disconnectedness have potentially far reaching consequences for sustainability. Biospheric disconnection is characterized by a strong dependence on industrial inputs which delay or displace ecological constraints (Norgaard 1988, Martinez-Alier *et al* 2014). This raises concerns about intergenerational justice, because it creates societal

structures that cannot be maintained indefinitely, and diminishes the biosphere's life-supporting conditions for future generations (e.g. through causing climate change). Similarly, spatial disconnection can result in the net appropriation of resources which create unsustainable lifestyle patterns (Brand and Wissen 2012, 2013) through teleconnections (Tukker *et al* 2014, Wiedmann *et al* 2015) that potentially disadvantage the 'source' regions. Spatial disconnectedness may thus compromise intragenerational justice, especially if the teleconnections are strong and unbalanced (Dorninger and Eisenmenger 2016, Teixidó-Figueras *et al* 2016).

### 2.1. Intraregional connectedness

Before considering the effects of biospheric and spatial disconnectedness in detail, it is necessary to develop a regional baseline for comparison. To this end, we first define *intraregional connectedness* as comprising the extent to which humans appropriate net primary production (NPP) for their own purposes, in combination with the labor used to appropriate this energy. A balance is required between regionally self-sufficient use of (ecologically derived) material and energy by humans and the availability of such flows to other species. The extent to which humans appropriate the NPP of the terrestrial ecosystems and the amount of trophic energy remaining in the ecosystems for other species indicates the level to which humans directly interact with, and source energy and materials from, ecosystems. In practice, intraregional connectedness may be measured via estimates of human appropriation of net primary production (HANPP) (Imhoff *et al* 2004, Haberl *et al* 2007b) and the labor inputs required to appropriate the NPP.

Direct human and animal labor in land use activities must be considered in the assessment of intraregional connectedness for several reasons. First, labor input is an important factor in the appropriation of net primary production: A system where net primary production is appropriated mainly by human and animal labor is likely to have very different sustainability outcomes than one where the appropriation is largely enabled by fossil fuel usage, even if the two systems have similar levels of HANPP. Second, direct labor is a form of internal input as long as working people and animals are 'fueled' by regional biomass products (Tello *et al* 2016). Third, from a human–nature connectedness perspective direct labor input in land use activities fosters rather than decreases biophysical and cognitive human–nature relationships (Cumming *et al* 2014, Webber *et al* 2015, Soga and Gaston 2016).

### 2.2. Biospheric disconnectedness

The relevant systems boundary for identifying *biospheric disconnectedness* is formed by the biosphere—the sphere of Earth where living organisms are found (Allaby 2008)—excluding, for example, the


lithosphere, where minerals are sourced from. Thus, all mineral and non-renewable material and energy flows, no matter if they were sourced from inside or outside the spatial boundaries of the region, are considered as non-internal flows. However, considering the increase in global trade flows it is still useful to differentiate between regionally sourced and imported minerals that are used for land use related activities, i.e. the production, extraction and processing of ecological goods. In fact, minerals imported for land use related activities create both biospheric and spatial disconnection (see section 2.3 and figure 1).

The degree of biospheric disconnectedness is determined by (1) the direct and embodied flows of mineral inputs (in the form of agrochemicals, fossil fuels, or materials embodied in machinery) that are drawn from outside the biosphere; and (2) waste flows and emissions caused by the use of such inputs (e.g. greenhouse gas emissions). To grasp the full extent of material and energy requirements within the land use system, it is necessary to account not only for direct non-biospheric inflows, for example, the use of fossil fuel based artificial fertilizers, but also for indirect flows, for example, the energy, material, and labor inputs which were necessary to build an agricultural vehicle or the energy required for producing chemical fertilizers (see table 1).

Intensified agricultural practices from the 1950s onwards have led to increased yields (Pimentel *et al* 1973, Pimentel 2009, Martinez-Alier 2011). However, this short-term boost of regional net primary production (NPP) is typically driven by phosphorus, nitrogen and fossil fuels drawn from outside the biosphere (Erb *et al* 2012, Niedertscheider *et al* 2016). The exhaustion of non-renewable materials and the associated production of wastes and pollution during the use of such resources cause serious sustainability problems (Daly 1990). Addressing the 'displaced' impacts of those problems (Haberl *et al* 2002) both temporally (e.g. resource depletion, climate change) and spatially (trade related environmental burden shifting to distant regions), is particularly problematic without a detailed understanding of the non-biospheric energy and material flows that cause them.

### 2.3. Spatial disconnectedness

Regional land use systems are increasingly connected to distal regions via global markets (MacDonald 2013, Henders *et al* 2015, Chaudhary and Kastner 2016). It is, therefore, vital to include and identify interregional exchange relationships in any framework that describes biophysical connectedness. Trade flows of crops and other biomass commodities create biophysical connections to distant places, increasing the disconnect from the regional natural productivity (NPP) (Krausmann *et al* 2008, Mayer *et al* 2015). We define biological resources drawn from within the defined regional boundaries as internal flows, and consequently understand all other biological resources



**Table 1.** External inputs into the land use system (Pimentel *et al* 1973 2008, Tello *et al* 2016). The left column lists material and energy inputs that enter the land use system from outside the biosphere and cause biospheric disconnectedness. These inputs may either come from regional sourcing or imports. The right column includes relevant intraregional trade of biomass based commodities which cause spatial disconnectedness.


External inputs	
Industrial mineral inputs (differentiated by regional sourcing or imports)	Biomass imports
<ul style="list-style-type: none"> <li>- Machinery use</li> <li>- Associated fossil fuels for machinery</li> <li>- Agrochemicals (e.g. artificial fertilizers, pesticides)</li> <li>- Irrigation and water pump facilities</li> <li>- Industrial seed production</li> <li>- Embodied flows of materials, energy, and labor for the production of all that items</li> </ul>	<ul style="list-style-type: none"> <li>- Embodied HANPP flows (section 4) of: <ul style="list-style-type: none"> <li>• Livestock feed</li> <li>• Food</li> <li>• Fiber and timber for textiles and construction</li> <li>• Livestock</li> <li>• Biomass for bioenergy purposes</li> </ul> </li> <li>- Embodied flows of materials, energy, and labor in all that commodities</li> </ul>

flowing into the region as ‘external inputs’ (table 1). *Spatial disconnectedness* can therefore be quantified via the amount of biomass based commodities imported to and exported from a region. Moreover, the import of minerals for land use related activities can be considered as an additional form of spatial disconnection. In order to reveal the full extent of disconnectedness, the embodied flows of material and energy associated with those biomass based imports and exports should also be accounted for.

Here it is important to note that trade-enabled material and energy exchanges between regions do not per se compromise sustainability. Some studies stress

the economic and ecological efficiency gains that arise from free trade and long distance relationships (Bhagwati 2007, Martinez-Melendez and Bennett 2016). Other scholars observe asymmetric power relationships and systematic inequalities in ‘ecologically unequal exchange’ relationships (Hornborg and Jorgenson 2013, Dorninger and Hornborg 2015) which provide only the pretense of efficiency and decoupling gains (Weinzettel *et al* 2013, Wiedmann *et al* 2015, Bergmann and Holmberg 2016). Regardless of the contention regarding the benefits of such land use related trades, we can say that distal trade relations always cause spatial human–nature disconnections.





These distal relations, or ‘teleconnections’ (Adger *et al* 2009, Haberl *et al* 2009, Yu *et al* 2013), not only involve long distance transportation (Cristea *et al* 2013) and environmental load displacement (Peters *et al* 2011, Peng *et al* 2016), but crucially also the substitution of regionally available biospheric resources by distal ones. This increases the complexity of the environmental and societal impacts arising from a given land use and cognitive and psychological disconnectedness from the environment (Kissinger and Rees 2010).

Figure 1 shows the conceptual steps towards assessing the levels of biophysical human–nature connectedness and potential disconnections.

### 3. Archetypical examples

We illustrate our conceptual framework for different regional land use systems with four archetypical systems. For each of the four systems the height of each component indicates the relevant throughput of energy and materials related to intraregional connectedness, spatial and biospheric disconnections (figure 2). A self-sufficient, non-industrialized, subsistence system which does not use any non-renewables in their land use practices, but relies solely on biomass goods and on relatively high labor input, has neither spatial nor biospheric disconnections (figure 2(a)). Such systems represent subsistence farming regions which were

common especially before the 20th century in most parts of the world (Krausmann 2001, Erb *et al* 2008). Moderately industrialized systems exhibit moderate levels of external inputs which allow a comparatively higher NPP appropriation with significant lower labor input (figure 2(b)). Such systems may include regions in transition from an agrarian to a more industrial society, for example, regions of Eastern Europe (Hanspach *et al* 2014, Loos *et al* 2014b). In contrast, a strongly exported oriented, highly industrialized system with high NPP availability is both spatially and biospherically more disconnected (figure 2(c)), for example, export-oriented soybean production regions in Brazil (Wittman *et al* 2016). Finally, an industrialized system with a high HANPP and high external inputs indicates strong regional disconnection and both temporal and spatial displacement of environmental burdens (figure 2(d)). Similar systems are likely to be found in densely populated, largely urbanized and wealthy regions such as Western Europe (Niedertscheider *et al* 2014).

Regions where direct labor input has largely been displaced by external inputs may exhibit a similar HANPP, but differ greatly with regard to the other two dimensions—biospheric and spatial disconnection (figures 2(b) and (d)), this has far reaching sustainability outcomes not only for the focal regions, but also for the distant regions they are connected to. Identifying the nature and extent of such regional disconnections is a crucial first step in addressing the

**Table 2.** Methods register and key references. The left column lists the methods to quantify regional biophysical human–nature connectedness. The relevant key references for each approach are provided in the right column. The ideal units of measurement are given in square brackets (where applicable), where [t] stands for metric tons, [J] for joules, and [h] for hours.

Methods and models of environmental accounting	Key references
Human appropriation of net primary production (HANPP) [t] [J]	• Vitousek <i>et al</i> 1986, Haberl <i>et al</i> 2007a
Material and energy flow analysis (MEFA) [t] [J]	• Haberl <i>et al</i> 2004, Fischer-Kowalski and Haberl 2007
Accounting of embodied flows	
• environmentally extended input output analysis (EEIOA)	• Leontief 1970, Kitzes 2013
• extension factors	• Pimentel <i>et al</i> 2008, Kastner <i>et al</i> 2015
Types of embodied flows	
• embodied HANPP (eHANPP) [t] [J]	• Erb <i>et al</i> 2009, Haberl <i>et al</i> 2009
• raw material equivalents [t] [J]	• Schaffartzik <i>et al</i> 2015b, Eisenmenger <i>et al</i> 2016
• embodied labor [J] [h]	• Alsamawi <i>et al</i> 2014, Simas <i>et al</i> 2015
• embodied energy [J]	• Agostinho and Siche 2014, Aguilera <i>et al</i> 2015

cross-scale sustainability challenges related to such interconnected systems. Without genuine reconnectedness, humans are only at best peripherally aware of the full range of impacts their lifestyle has on other and future generations, and on other species. A more complete understanding of human–nature connectedness and opportunities to reconnect, might increase the leverage potential of actions set in land use systems towards transformational change (Meadows 1999, Abson *et al* 2016). It is to be hoped for that biophysically reconnected regional land use systems ultimately promote a more foresightful, responsible and conscious society, based on a living with rather than dominating nature.

In regionally connected land use systems, largely reliant on (transformed) solar energy and labor as major energy inputs, in- and outputs will then be reconnected to the natural cycles—the regeneration and uptake rate—of the biosphere (Folke *et al* 2016). A reconnected land use system will strengthen self-sufficiency, circularity in production and consumption; involve less teleconnections, less specialization, more diverse land uses, and relations of trust (Tregear 2011, Weatherell *et al* 2003). Here the major balancing challenge is ensuring that sufficient biospheric resources are appropriated for human well-being while retaining resources available for the flourishing of other species.

In all cases the assessment of regional biophysical connectivity, particularly if linked to other regional indicators, can help identify regionally specific challenges in transitioning towards more sustainable land use systems. In addition, multi-scalar assessments may help identify ‘natural’ scales of biophysical connectedness and appropriate scales for managing material and energy flows.

#### 4. From theory to practice: methodological guidelines


In this section we present methodological guidance to operationalize our concept of biophysical human–nature connectedness at regional scales. Building on

well-established methods (table 2) this operationalization will allow assessment of the extent to which systems are built on and driven by intraregional connectedness and biospheric and spatial disconnectness respectively (figure 3).

As discussed above, we consider HANPP as an appropriate starting point to quantify intraregional connectedness. HANPP is based on not only appropriated biomass yields from farming, grazing, and forestry, but also harvest related losses, unused biomass extraction, conversion losses, and land use conversion—changes in the HANPP fraction due to indirect changes to NPP. A land use conversion effect can only be quantified in relation to the potential net primary production ( $NPP_{pot}$ ) that would occur at a certain area without any human interference. A range of different models exist that allow for a computation of site-specific photosynthesis performance (Haberl *et al* 2014). For example, the Miami model (Lieth 1975) calculates  $NPP_{pot}$  from average precipitation and annual mean temperature of an area. Other models additionally include information on soil texture, latitude, and  $CO_2$  availability (Sitch *et al* 2003). By subtracting the HANPP, i.e. all harvest and related flows plus the land use conversion, from the  $NPP_{pot}$ , one arrives at the NPP that remained in the ecosystem after harvest and which is available for other species ( $NPP_{eco}$ ) (Krausmann *et al* 2013, Plutzer *et al* 2015). By going beyond simple harvest or yield assessments HANPP reveals the connectedness to the productivity, and the potentially renewable resources, of ecosystems.

From an ecological perspective, low HANPP may be a desirable goal because it leaves a large amount of energy to other species (Haberl *et al* 2007a). In contrast, if low HANPP values are achieved via the use of non-renewable resources or distant biomass the overall outcomes for sustainability may still be negative (with regards to future generations and distant regions). However, as the conventional HANPP method neither captures external inputs, such as the materials and substances that are used to





produce and harvest goods in the land use system (Haas and Krausmann 2015), nor trade related teleconnections (Haberl *et al* 2009, Kastner *et al* 2015), nor the labor inputs to those systems, further methodological steps are required to evaluate regional scale biophysical human–nature connectedness.

Biospheric disconnections can be assessed via a social–metabolic analysis of the regional land use system. Social metabolism quantifies, similar to the metabolism of organisms, the biophysical inputs and outputs of a social–ecological entity. It is operationalized by material and energy flow accounting analysis (MEFA) (Fischer-Kowalski and Haberl 2007). While analysis of socio-ecological energy and material flows is well established, particularly at the national level (Fischer-Kowalski *et al* 2011, Haberl *et al* 2004), the notion and consequences of changes to ‘regional biophysical human–nature disconnections’ is hardly explored in that literature. However, by conducting such a MEFA analysis one is able to calculate the throughput of materials and energy of the land use system and subsequently relate these flows to regional and cross-scalar sustainability challenges.

The system boundaries of the adopted MEFA analysis are defined by the spatial boundaries of the region (Sastre *et al* 2015) and the boundaries of the biosphere. In order to account for differing levels of teleconnectedness, it is important to differentiate between regionally sourced mineral inputs and imported ones. Doing so can help reveal related additional transport costs, patterns of ecologically

unequal exchange and outsourcing of material and energy intensive processes. Industrial mineral inputs, such as machinery use, fuels, or agrochemicals, enter the system from outside the biosphere (left column of table 1) and potentially from outside the region. Outflows are those materials and substances that are not reused in the land use system but create pollution, wastes, and emissions.

Artificial fertilizer, seeds and machinery production processes are an energy and material intensive endeavor (Pimentel *et al* 2008). In order to reveal the full extent of energy, materials and labor required for the external industrial inputs into the land use system, the embodied flows of those inputs must be accounted for. We suggest using either product and region specific extension factors (Kastner *et al* 2015, Schaffartzik *et al* 2015a), or regionally adjusted environmentally-extended input-output analysis (EEIOA) (Miller and Blair 2009, Kitzes 2013, Schaffartzik *et al* 2014) where inter-sectoral linkages can be retraced, i.e. the flows between the land use sector and other socio-economic sectors of the region.

The third and last methodological measure of the framework is to collect data on the biomass based inter-regionally traded goods for quantifying the spatial disconnect. In short, all directly traded biomass commodities (table 1) such as crops, animal products, textiles, other fibers, bioenergy products, and wood products, and the indirect flows of NPP, materials, energy, and labor embodied in these goods need to be captured. The latter are usually not reported in trade

statistics. Still, the methodological goal is to redistribute the flows embodied in the goods from the place of origin to the place of final consumption (Kastner *et al* 2015, Wiedmann 2016).

To comprehensively reveal the biophysical processes necessary to produce a specific commodity and to disclose how the consumption of traded goods affects connectedness an environmentally-extended input-output table or extension factors, which are adjusted for the specific region and year, would potentially provide the best systematic approach to assess the embodied flows of traded products. The results of embodied HANPP (Erb *et al* 2009), raw material equivalents (Schaffartzik *et al* 2015b, Eisenmenger *et al* 2016), embodied energy (Agostinho and Siche 2014, Perryman and Schramski 2015), or embodied labor (Alsamawi *et al* 2014, Simas *et al* 2015) are established indicators, increasingly used in the scientific literature to reveal international inequalities and related environmental pressures (Teixidó-Figueras *et al* 2016).

Different types of flows have different metrics: NPP and HANPP can be expressed in terms of dry matter [t], carbon [t] or energy units [J]; labor input in time units [h] or energy [J]; materials in units of mass [t] or enthalpy [J]; emissions in GHG potentials [t CO<sub>2</sub> equivalents] and nitrogen leaching [NO<sub>3</sub>]. For achieving comparability between regions we suggest to evaluate connectedness on a per unit area, or per capita basis; comparability within regions may be achieved via an expression of flows in energy units (except emissions), i.e. flows of HANPP, material, and labor. The final result of the framework provides a measure of the degree to which a regional land use system is biophysically connected to the productivity of the regional ecosystems (NPP) and disconnected in terms of external inputs (figure 3).

The empirical application of this framework will likely involve challenges with regards to data availability and the computation of critical embodied flows. In particular, identification and assessment of interregional trade flows from material accounting data will involve region-specific difficulties. For example, physical trade relations between regions might not be reported by authorities. Therefore, we encourage consultation of relevant stakeholders to assure the validity of data where necessary. Likewise, the calculation of embodied flows is a sensitive methodological endeavor (Schaffartzik *et al* 2015b). It will therefore be important to provide detailed information on steps of the decisions that have been made regarding data sources and estimations to ensure transparency and traceability.

## 5. Outlook

We argue that the regional land use system is an appropriate unit of analysis for investigating biophysi-

cal connectedness as it provides a focal unit for understanding cross scale interactions between land use systems, revealing key environmental feedback loops in and between regions. We recognize that we take a relatively pragmatic definition of ‘region’. Yet, in principle it should be possible to use this approach to identify spatial extents within which there are high levels of connectedness or across which significant disconnections occur.

Biophysical human–nature connectedness is increasingly overlain and suppressed by modes of industrial land use, which entails teleconnections and external non-renewable inputs. In this paper we introduced a new approach to conceptualize biophysical human–nature connectedness at regional scales and related it to potential sustainability outcomes. Building on *a priori* state of biophysical connectedness, we identified two major realms of disconnectedness: (1) external non-renewable inputs that enter the land use system and (2) teleconnections with distant systems, both of which decrease regional connectedness.

While the conceptual framework itself represents a novel perspective on land use management, the combined methods for each part are well established. Together these methods allow for comparisons of different ‘types’ and degrees of the connectedness between different regions, which in turn can be related to other regional characteristics or sustainability outcomes (e.g. Wittman *et al* 2016). The framework is designed to be applicable to regions anywhere in the world and to encourage researchers and policymakers to develop a more holistic approach regarding cross-scale, sustainable land management issues not captured by other frameworks (e.g. sustainable intensification (Barnes and Thomson 2014, Loos *et al* 2014a), or land sparing (Fischer *et al* 2014)).

Instead of making human–nature connections evermore complex and opaque by increasing external inputs via industrial technology, a genuinely reconnected system will have a higher internal self-reliance, through a more self-sufficient land use system. Such regionally reconnected systems may facilitate more foresightful, responsible and conscious behaviors. We believe that there are various opportunities to strengthen connectedness of humans to nature. For example, by a re-regionalized economy, a higher degree of self-sufficiency, lower degrees of dependence on external (non-renewable or distant) inputs, by internal biomass reuse (Galán *et al* 2016, Tello *et al* 2016), permaculture, agroforestry, organic farming, small-scale farming, low external input technology farming (Tripp 2005), lower consumption patterns (especially of NPP intensive products, like animal products), less overproduction and consequently less food and biomass ‘wastes’. The operationalization of this model can be applied as a heuristic tool to reveal complex social–ecological interlinkages, raising awareness of the challenge in managing biophysical connections across scales. This in turn might help to shift the

focus from ‘on site’ efficiency thinking in land use management to a more comprehensible and holistic perspective on human–nature connectedness.

## Acknowledgments

This research was supported by the Volkswagen-stiftung and the Niedersächsisches Ministerium für Wissenschaft und Kultur funded project ‘Leverage Points for Sustainable Transformations: Institutions, People and Knowledge’ (Grant Number A112269). We would like to thank Fridolin Krausmann from the Institute of Social Ecology, IFF Vienna, Christopher Ives from the University of Nottingham, and Kathleen Klaniecki from the Leuphana Universität Lüneburg for their helpful comments on earlier drafts of this article. The authors declare no conflict of interest.

## References

- Abson D J *et al* 2016 Leverage points for sustainability transformation *Ambio* **45** 1–10
- Adger W N, Eakin Hallie and Winkels A 2009 Nested and teleconnected vulnerabilities to environmental change *Front. Ecol. Environ.* **7** 150–7
- Agostinho F and Siche R 2014 Hidden costs of a typical embodied energy analysis: Brazilian sugarcane ethanol as a case study *Biomass Bioenergy* **71** 69–83
- Aguilera E, Guzmán G I, Infante-Amate J, García-Ruiz R, Herrera A, Villa I, Torremocha E, Carranza G and González de Molina M 2015 Incorporating a historical perspective *Embodied Energy in Agricultural Inputs Report*
- Allaby M 2008 *Dictionary of Earth Sciences* (Oxford: Oxford Reference)
- Alsamawi A, Murray J and Lenzen M 2014 The employment footprints of nations: Uncovering master-servant relationships *J. Ind. Ecol.* **18** 59–70
- Andersson E, Barthel S, Borgström S, Colding J, Elmqvist T, Folke C and Gren A 2014 Reconnecting cities to the biosphere: stewardship of green infrastructure and urban ecosystem services *Ambio* **43** 445–53
- Asafu-Adjaye J *et al* 2015 *An Ecomodernist Manifesto*
- Barnes A P and Thomson S G 2014 Measuring progress towards sustainable intensification: how far can secondary data go? *Ecol. Indic.* **36** 213–20
- Bergmann L and Holmberg M 2016 Land in Motion *Ann. Am. Assoc. Geogr.* **106** 932–56
- Bhagwati J 2007 *In Defense of Globalization: With a New Afterword* (Oxford: Oxford University Press)
- Boulding K E 1966 The economics of the coming spaceship Earth *Environmental Quality Issues in a Growing Economy* ed H Jarrett (Baltimore: Johns Hopkins University Press) pp 3–14
- Brand U and Wissen M 2013 Crisis and continuity of capitalist society–nature relationships: the imperial mode of living and the limits to environmental governance *Rev. Int. Polit. Econ.* **2290** 1–25
- Brand U and Wissen M 2012 Global environmental politics and the imperial mode of living: articulations of state–capital relations in the multiple crisis *Globalizations* **9** 547–60
- Chaudhary A and Kastner T 2016 Land use biodiversity impacts embodied in international food trade *Glob. Environ. Change* **38** 195–204
- Cooke B, West S and Boonstra W J 2016 Dwelling in the biosphere: exploring an embodied human–environment connection in resilience thinking *Sustain. Sci.* **11** 1–13
- Cristea A, Hummels D, Puzzello L and Avetisyan M 2013 Trade and the greenhouse gas emissions from international freight transport *J. Environ. Econ. Manage.* **65** 153–73
- Cumming G S, Buerkert A, Hoffmann E M, Schlecht E, Von Cramon-Taubadel S and Tschardt T 2014 Implications of agricultural transitions and urbanization for ecosystem services *Nature* **515** 50–7
- Daly G 1997 *Nature's Services: Societal Dependence on Natural Ecosystems* (Washington, DC: Island Press)
- Daly H E 1990 Toward some operational principles of sustainable development *Ecol. Econ.* **2** 1–6
- Dearing J A *et al* 2014 Safe and just operating spaces for regional social–ecological systems *Glob. Environ. Change* **28** 227–38
- Dorninger C and Eisenmenger N 2016 South America's biophysical involvement in international trade: the physical trade balances of Argentina, Bolivia, and Brazil in the light of ecologically unequal exchange *J. Polit. Econ.* **23** 394–409
- Dorninger C and Hornborg A 2015 Can EEMRIO analyses establish the occurrence of ecologically unequal exchange? *Ecol. Econ.* **119** 414–8
- Eisenmenger N, Wiedenhofer D, Schaffartzik A, Giljum S, Bruckner M, Schandl H, Wiedmann T O, Lenzen M, Tukker A and Koning A 2016 Consumption-based material flow indicators—comparing six ways of calculating the Austrian raw material consumption providing six results *Ecol. Econ.* **128** 177–86
- Erb K-H, Gingrich S, Krausmann F and Haberl H 2008 Industrialization, fossil fuels, and the transformation of land use: an integrated analysis of carbon flows in Austria 1830–2000 *J. Ind. Ecol.* **12** 686–703
- Erb K-H, Haberl H, DeFries R, Ellis E C, Krausmann F and Verburg P H 2012 Pushing the planetary boundaries *Science* **338** 1419–20
- Erb K-H, Krausmann F, Lucht W and Haberl H 2009 Embodied HANPP: mapping the spatial disconnect between global biomass production and consumption *Ecol. Econ.* **69** 328–34
- Fischer-Kowalski M and Haberl H 2007 *Socioecological Transitions and Global Change: Trajectories of Social Metabolism and Land Use* (Cheltenham: Edward Elgar Publishing)
- Fischer-Kowalski M, Krausmann F, Giljum S, Lutter S, Mayer A, Bringezu S, Moriguchi Y, Schütz H, Schandl H and Weisz H 2011 Methodology and indicators of economy-wide material flow accounting: State of the art and reliability across sources *J. Ind. Ecol.* **15** 855–76
- Fischer-Kowalski M and Swilling M 2011 *Decoupling natural resources use and environmental impacts from economic growth*
- Fischer J, Abson D J, Butsic V, Chappell M J, Ekroos J, Hanspach J, Kuemmerle T, Smith H G and von Wehrden H 2014 Land sparing versus land sharing: moving forward *Conserv. Lett.* **7** 149–57
- Folke C, Biggs R, Norström A V, Meyers B and Rockström J 2016 Social–ecological resilience and biosphere-based sustainability science *Ecol. Soc.* **21** 41
- Folke C *et al* 2011 Reconnecting to the biosphere *Ambio* **40** 719–38
- Galán E S *et al* 2016 Widening the analysis of Energy Return On Investment (EROI) in agro-ecosystems: socio-ecological transitions to industrialized farm systems (the Vallès County, Catalonia, c.1860 and 1999) *Ecol. Modell.* **336** 13–25
- Haas W and Krausmann F 2015 Rural metabolism: material flows in an Austrian village in 1830 and 2001 *Social Ecology Working Paper* 155 (Vienna: Institute of Social Ecology)
- Haberl H, Erb K-H and Krausmann F 2014 Human appropriation of net primary production: patterns, trends, and planetary boundaries *Annu. Rev. Environ. Resour.* **39** 363–91
- Haberl H, Erb K-H, Krausmann F, Berecz S, Ludwiczek N, Martínez-Alier J, Musel A and Schaffartzik A 2009 Using embodied HANPP to analyze teleconnections in the global land system: conceptual considerations *Geogr. Tidsskr. J. Geogr.* **109** 119–30

- Haberl H, Erb K-H, Plutzar C, Fischer-Kowalski M and Krausmann F 2007a Human appropriation of net primary production (HANPP) as indicator for pressures on biodiversity *Sustainability Indicators. A Scientific Assessment* ed T Hak, B Moldan and A L Dahl (Washington, DC: Island Press) pp 271–88
- Haberl H, Erb K H, Krausmann F, Gaube V, Bondeau A, Plutzar C, Gingrich S, Lucht W and Fischer-Kowalski M 2007b Quantifying and mapping the human appropriation of net primary production in Earth's terrestrial ecosystems *Proc. Natl Acad. Sci. USA* **104** 12942–7
- Haberl H, Fischer-Kowalski M, Krausmann F, Schandl H, Weisz H and Winiwarter V 2002 Theoretical foundations of societal observation of sustainable development *Die Bodenkultur* **53** 55–63
- Haberl H, Fischer-Kowalski M, Krausmann F, Weisz H and Winiwarter V 2004 Progress towards sustainability? What the conceptual framework of material and energy flow accounting (MEFA) can offer *Land Use Policy* **21** 199–213
- Hanspach J, Hartel T, Milcu A I, Mikulcak F, Dorresteijn I, Loos J and Von Wehrden H 2014 A holistic approach to studying social–ecological systems and its application to southern Transylvania *Ecol. Soc.* **19** 32
- Henders S, Persson U M and Kastner T 2015 Trading forests: land-use change and carbon emissions embodied in production and exports of forest-risk commodities *Environ. Res. Lett.* **10** 125012
- Hornborg A and Jorgenson A K 2013 *International Trade and Environmental Justice: Toward a Global Political Ecology* (Lund, Sweden: Nova Science)
- Imhoff M, Bounoua L, Ricketts T, Loucks C, Harriss R and Lawrence W T 2004 Global patterns in human consumption of net primary production *Nature* **429** 870–3
- Kastner T, Erb K-H and Haberl H 2014 Rapid growth in agricultural trade: effects on global area efficiency and the role of management *Environ. Res. Lett.* **9** 034015
- Kastner T, Erb K H and Haberl H 2015 Global human appropriation of net primary production for biomass consumption in the European Union, 1986–2007 *J. Ind. Ecol.* **19** 825–36
- Kissinger M and Rees W E 2010 Importing terrestrial biocapacity: The US case and global implications *Land Use Policy* **27** 589–99
- Kitzes J 2013 An introduction to environmentally-extended input-output analysis *Resources* **2** 489–503
- Krausmann F 2001 Land use and industrial modernization: an empirical analysis of human influence on the functioning of ecosystems in Austria 1830–1995 *Land Use Policy* **18** 17–26
- Krausmann F, Erb K-H, Gingrich S, Haberl H, Bondeau A, Gaube V, Lauk C, Plutzar C and Searchinger T D 2013 Global human appropriation of net primary production doubled in the 20th century *Proc. Natl Acad. Sci.* **110** 10324–9
- Krausmann F, Erb K H, Gingrich S, Lauk C and Haberl H 2008 Global patterns of socioeconomic biomass flows in the year 2000: a comprehensive assessment of supply, consumption and constraints *Ecol. Econ.* **65** 471–87
- Leontief W 1970 Environmental repercussions and the economic structure: an input-output approach *Rev. Econ. Stat.* **52** 262–71
- Lieth H 1975 Modeling the primary productivity of the world *Primary Productivity of the Biosphere* ed H Lieth and R H Whittaker (New York: Springer) pp 237–63
- Loos J, Abson D J, Chappell M J, Hanspach J, Mikulcak F, Tichit M and Fischer J 2014a Putting meaning back into 'sustainable intensification' *Front. Ecol. Environ.* **12** 356–61
- Loos J, Dorresteijn I, Hanspach J, Fust P, Rakosy L and Fischer J 2014b Low-intensity agricultural landscapes in Transylvania support high butterfly diversity: implications for conservation *PLoS One* **9** e103256
- MacDonald G K 2013 Eating on an interconnected planet *Environ. Res. Lett.* **8** 021002
- Martinez-Alier J 2011 The EROI of agriculture and its use by the Via Campesina *J. Peasant Stud.* **38** 145–60
- Martinez-Alier J and Muradian R 2015 Taking stock: the keystones of ecological economics *Handbook of Ecological Economics* (Cheltenham: Edward Elgar Publishing) pp 1–25
- Martinez-Alier J, Temper L and Demaria F 2014 Social metabolism and environmental conflicts in India *Indi@logs* **1** 51–83
- Martinez-Melendez L A and Bennett E M 2016 Trade in the US and Mexico helps reduce environmental costs of agriculture *Environ. Res. Lett.* **11** 055004
- Mayer A, Schaffartzik A, Haas W and Sepulveda A R 2015 Patterns of global biomass trade—Implication for food sovereignty and socio-environmental conflicts *EJOLT Report* 20
- Meadows D 1999 Leverage points. Places to intervene in a system *Sustainability Institute Report*
- Miller R E and Blair P D 2009 *Input–Output Analysis: Foundations and Extensions* (Cambridge: Cambridge University Press)
- Niedertscheider M, Kastner T, Fetzl T, Haberl H, Kroisleitner C, Plutzar C and Erb K-H 2016 Mapping and analysing cropland use intensity from a NPP perspective *Environ. Res. Lett.* **11** 014008
- Niedertscheider M, Kuemmerle T, Müller D and Erb K-H 2014 Exploring the effects of drastic institutional and socio-economic changes on land system dynamics in Germany between 1883 and 2007 *Glob. Environ. Change* **28** 98–108
- Norgaard R B 1988 Sustainable development: a co-evolutionary view *Futures* **20** 606–20
- Pearson C J 2007 Regenerative, semiclosed systems: a priority for twenty-first-century agriculture *Bioscience* **57** 409–18
- Peng S, Zhang W and Sun C 2016 'Environmental load displacement' from the North to the South: a consumption-based perspective with a focus on China *Ecol. Econ.* **128** 147–58
- Perryman M E and Schramski J R 2015 Evaluating the relationship between natural resource management and agriculture using embodied energy and eco-exergy analyses: a comparative study of nine countries *Ecol. Complex.* **22** 152–61
- Peters G P, Minx J C, Weber C L and Edenhofer O 2011 Growth in emission transfers via international trade from 1990 to 2008 *Proc. Natl Acad. Sci. USA* **108** 8903–8
- Pimentel D 2009 Energy inputs in food crop production in developing and developed nations *Energies* **2** 1–24
- Pimentel D, Hurd L E, Bellotti A C, Forster M J, Oka I N, Sholes O D and Whitman R J 1973 Food production and the energy crisis *Science* **182** 443–9
- Pimentel D, Williamson S, Alexander C E, Gonzalez-Pagan O, Kontak C and Mulkey S E 2008 Reducing energy inputs in the US food system *Hum. Ecol.* **36** 459–71
- Plutzar C *et al* 2015 Changes in the spatial patterns of human appropriation of net primary production (HANPP) in Europe 1990–2006 *Reg. Environ. Change* **16** 1225–38
- Sastre S, Carpintero Ó and Lomas P L 2015 Regional material flow accounting and environmental pressures: the spanish case *Environ. Sci. Technol.* **49** 2262–9
- Schaffartzik A, Haberl H, Kastner T, Wiedenhofer D, Eisenmenger N and Erb K H 2015a Trading land: a review of approaches to accounting for upstream land requirements of traded products *J. Ind. Ecol.* **19** 703–14
- Schaffartzik A, Sachs M, Wiedenhofer D and Eisenmenger N 2014 Environmentally extended input-output analysis *Social Ecology Working Paper* 154 (Vienna: Institute of Social Ecology)
- Schaffartzik A, Wiedenhofer D and Eisenmenger N 2015b Raw material equivalents: the challenges of accounting for sustainability in a globalized world *Sustainability* **7** 5345–70
- Seppelt R and Cumming G S 2016 Humanity's distance to nature: time for environmental austerity? *Landsc. Ecol.* **31** 1645–51

- Seppelt R, Manceur A M, Liu J, Fenichel E P and Klotz S 2014 Synchronized peak-rate years of global resources use *Ecol. Soc.* **19** 50
- Simas M, Wood R and Hertwich E 2015 Labor embodied in trade *J. Ind. Ecol.* **19** 343–56
- Sitch S *et al* 2003 Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model *Glob. Change Biol.* **9** 161–85
- Soga M and Gaston K J 2016 Extinction of experience: the loss of human–nature interactions *Front. Ecol. Environ.* **14** 94–101
- Teixidó-Figueras J, Steinberger J K, Krausmann F, Haberl H, Wiedmann T, Peters G P, Duro J A and Kastner T 2016 International inequality of environmental pressures: decomposition and comparative analysis *Ecol. Indic.* **62** 163–73
- Tello E *et al* 2016 Opening the black box of energy throughputs in farm systems: a decomposition analysis between the energy returns to external inputs, internal biomass reuses and total inputs consumed (the Vallès County, Catalonia, c.1860 and 1999) *Ecol. Econ.* **121** 160–74
- Tregear A 2011 Progressing knowledge in alternative and local food networks: critical reflections and a research agenda *J. Rural Stud.* **27** 419–30
- Tripp R 2005 The performance of low external input technology in agricultural development: a summary of three case studies *Int. J. Agric. Sustain.* **3** 143–53
- Tukker A, Bulavskaya T, Giljum S and De Koning A 2014 *The Global Resource Footprint of Nations. Carbon, water, land and materials embodied in trade and final consumption calculated with EXIOBASE*
- Vitousek P M, Ehrlich P R, Ehrlich A H and Matson P A 1986 Human appropriation of the products of photosynthesis *Bioscience* **36** 368–73
- Waggoner P E 1996 How much land can ten billion people spare for nature? *Daedalus* **125** 73–93
- Weatherell C, Tregear A and Allinson J 2003 In search of the concerned consumer: UK public perceptions of food, farming and buying local *J. Rural Stud.* **19** 233–44
- Webber J, Hinds J and Camic P M 2015 The well-being of allotment gardeners: a mixed methodological study *Ecopsychology* **7** 20–8
- von Wehrden H, Abson D J, Beckmann M, Cord A F, Klotz S and Seppelt R 2014 Realigning the land-sharing/land-sparing debate to match conservation needs: considering diversity scales and land-use history *Landsc. Ecol.* **29** 941–8
- Weinzettel J, Hertwich E G, Peters G P, Steen-Olsen K and Galli A 2013 Affluence drives the global displacement of land use *Glob. Environ. Change* **23** 433–8
- Wiedmann T 2016 Impacts embodied in global trade flows *Taking Stock of Industrial Ecology* ed R Clift and A Druckman (Berlin: Springer) pp 159–80
- Wiedmann T O, Schandl H, Lenzen M, Moran D, Suh S, West J and Kanemoto K 2015 The material footprint of nations *Proc. Natl Acad. Sci.* **112** 6271–6
- Wittman H, Chappell M, Abson D, Bezner Kerr R, Blesh J, Hanspach J, Perfecto I and Fischer J 2016 A social–ecological perspective on harmonizing food security and biodiversity conservation *Reg. Environ. Change* (<https://doi.org/10.1007/s10113-016-1045-9>)
- Wu J 2013 Landscape sustainability science: ecosystem services and human well-being in changing landscapes *Landsc. Ecol.* **28** 999–1023
- Yu Y, Feng K and Hubacek K 2013 Tele-connecting local consumption to global land use *Glob. Environ. Change* **23** 1178–86