

Bryophytes and organic layers control uptake of airborne nitrogen in low-N environments

Bähring, Alexandra; Fichtner, Andreas; Friedrich, Uta; Von Oheimb, Goddert; Härdtle, Werner

Published in:
Frontiers in Plant Science

DOI:
[10.3389/fpls.2017.02080](https://doi.org/10.3389/fpls.2017.02080)

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for pulished version (APA):
Bähring, A., Fichtner, A., Friedrich, U., Von Oheimb, G., & Härdtle, W. (2017). Bryophytes and organic layers control uptake of airborne nitrogen in low-N environments. *Frontiers in Plant Science*, 8, Article 2080. <https://doi.org/10.3389/fpls.2017.02080>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Bryophytes and Organic layers Control Uptake of Airborne Nitrogen in Low-N Environments

Alexandra Bähring¹, Andreas Fichtner¹, Uta Friedrich¹, Goddert von Oheimb² and Werner Härdtle^{1*}

¹ Institute of Ecology, Leuphana University of Lüneburg, Lüneburg, Germany, ² Institute of General Ecology and Environmental Protection, Technische Universität Dresden, Tharandt, Germany

OPEN ACCESS

Edited by:

Uwe Rascher,
Forschungszentrum Jülich, Germany

Reviewed by:

Ze-Xin Fan,
Xishuangbanna Tropical Botanical
Garden (CAS), China
Tobias Gebauer,
Albert Ludwigs University of Freiburg,
Germany

*Correspondence:

Werner Härdtle
haerdtle@uni-lueneburg.de

Specialty section:

This article was submitted to
Functional Plant Ecology,
a section of the journal
Frontiers in Plant Science

Received: 22 August 2017

Accepted: 21 November 2017

Published: 04 December 2017

Citation:

Bähring A, Fichtner A, Friedrich U,
von Oheimb G and Härdtle W (2017)
Bryophytes and Organic layers
Control Uptake of Airborne Nitrogen in
Low-N Environments.
Front. Plant Sci. 8:2080.
doi: 10.3389/fpls.2017.02080

The effects of atmospheric nitrogen (N) deposition on ecosystem functioning largely depend on the retention of N in different ecosystem compartments, but accumulation and partitioning processes have rarely been quantified in long-term field experiments. In the present study we analysed for the first time decadal-scale flows and allocation patterns of N in a heathland ecosystem that has been subject to airborne N inputs over decades. Using a long-term ¹⁵N tracer experiment, we quantified N retention and flows to and between ecosystem compartments (above-ground/below-ground vascular biomass, moss layer, soil horizons, leachate). After 9 years, about 60% of the added ¹⁵N-tracer remained in the N cycle of the ecosystem. The moss layer proved to be a crucial link between incoming N and its allocation to different ecosystem compartments (in terms of a short-term capture, but long-term release function). However, about 50% of the ¹⁵N captured and released by the moss layer was not compensated for by a corresponding increase in recovery rates in any other compartment, probably due to denitrification losses from the moss layer in the case of water saturation after rain events. The O-horizon proved to be the most important long-term sink for added ¹⁵N, as reflected by an increase in recovery rates from 18 to 40% within 8 years. Less than 2.1% of ¹⁵N were recovered in the podzol-B-horizon, suggesting that only negligible amounts of N were withdrawn from the N cycle of the ecosystem. Moreover, ¹⁵N recovery was low in the dwarf shrub above-ground biomass (<3.9% after 9 years) and in the leachate (about 0.03% within 1 year), indicating still conservative N cycles of the ecosystem, even after decades of N inputs beyond critical load thresholds. The continuous accumulation of reactive forms of airborne N suggests that critical load-estimates need to account for cumulative effects of N additions into ecosystems.

Keywords: *Calluna vulgaris*, critical load, heathland, nitrogen cycling, nitrogen retention, nitrogen saturation

INTRODUCTION

Atmospheric inputs of reactive nitrogen (comprising all N species with the exception of N₂; Galloway and Cowling, 2002) have tripled since the nineteenth century, resulting in unprecedented impacts on the N status of ecosystems (Galloway and Cowling, 2002; De Schrijver et al., 2011). Since N is the most limiting nutrient in many natural and semi-natural environments, high N loads may affect important ecosystem functions such as biomass production and the complex interplays

within and between plant species, but also their susceptibility to abiotic and biotic stressors such as drought, frost events and herbivory (Bobbink et al., 2010). As a consequence, atmospheric N deposition constitutes one of the most substantial threats to biodiversity today (Sala et al., 2000; Phoenix et al., 2012). This pertains to N-limited ecosystems with conservative N cycles in particular, because many species that are typical of these environments are physiologically adapted to low N availability, for example due to high N use efficiency or mycorrhizal associations (Aerts, 1999; Bobbink et al., 2002; Phoenix et al., 2012).

European heathland ecosystems are typically low-N environments (Härdtle et al., 2006). These heathlands host a huge proportion of the plant and animal diversity typical of open acidic sites, but also provide a wide range of ecosystem services (e.g., groundwater recharge, C-storage, recreation; De Graaf et al., 2009; Fagúndez and Izco, 2016). Agricultural intensification and elevated atmospheric N inputs, however, have contributed to a distinct decline in European heathland area over recent decades (Rose et al., 2000; Fagúndez, 2012; Southon et al., 2012), and nowadays remaining areas are protected by several international protection acts (such as the Natura2000 Habitat Council Directive 92/43/EEC) (Rose et al., 2000; Vandvik et al., 2005; Southon et al., 2012).

Many lowland heaths of NW Europe are characterised by the dominance of the dwarf shrub *Calluna vulgaris* (L.) Hull (henceforth referred to as *Calluna*). This species is competitively superior to grasses (e.g., *Molinia caerulea*, *Deschampsia flexuosa*) under low-N conditions, but becomes less competitive with increasing N availability (Alonso et al., 2001; Friedrich et al., 2011a; Fagúndez, 2012). At N-fertilised sites, a succession towards grass dominance is observable when the dwarf shrub canopy opens as a result of overaging, heather beetle outbreaks or frost and drought damage (Berdowski and Zeilinga, 1987; Bobbink et al., 2010). This process accelerates under increasing N availability, attributable to faster rates of N cycling and plant growth, and finally, shortened life-cycles of *Calluna* (Berendse, 1990; Power et al., 1998; Carroll et al., 1999). Surprisingly, dwarf shrub growth still is N-limited in some heathlands, despite decadal N inputs above critical loads (von Oheimb et al., 2010; Friedrich et al., 2011b). Such findings could be related to both time- and space-dependent allocation and storage processes of airborne N loads, or to underestimated N losses from the N pool of the ecosystem. It is, for example, conceivable that the spodic horizon (B-horizon) of podzols can accumulate considerable amounts of N in organic compounds (fulvic and humic acids; Nielsen et al., 2000; Friedrich et al., 2011b), all of which are to a large extent insoluble and not readily bio-available (Hagedorn et al., 2005). In this way, N inputs might be (at least partly) withdrawn from the N cycle and thus become biologically inactive. However, we still have limited evidence for the long-term efficacy of atmospheric N loads on the functioning of heathlands as well as other ecosystems. In addition, decadal-scale patterns of N flows and sequestration have rarely been quantified, restricting a mechanistic understanding of N-fertilisation effects on ecosystem functions. This in turn emphasises the value of long-term studies that trace the allocation of N to or between

ecosystem compartments on a decadal scale (Silvertown et al., 2010).

Many N-budget calculations from long-term fertilisation experiments have provided valuable knowledge about shifts in N stores of different ecosystem compartments (Power et al., 1998; Pilkington et al., 2005; von Oheimb et al., 2010), but were unable to quantify the origin of N gains and losses over time and, thus, N flows between ecosystem compartments such as plant biomass, organic layers or mineral soil horizons. ^{15}N tracer studies have the potential to overcome this problem and allow for a temporal quantification of N flows (Kleinebecker et al., 2009), as has been demonstrated for grassland and forest ecosystems (Gerzabek et al., 2004; Nadelhoffer et al., 2004). Quantitative analyses of the fate of airborne N loads including different ecosystem compartments are important for an understanding of ecosystem responses to long-term N inputs at the individual plant and community level, but may also provide insights into mechanisms underlying shifts in species composition and losses (Friedrich et al., 2011b). Such information can further help to develop adaptive management strategies for semi-natural N-limited ecosystems and the species they host, particularly in the face of ongoing N inputs and their contribution to biodiversity loss (Maskell et al., 2010).

The objective of the present study was to quantify decadal-scale patterns of N accumulation and partitioning in an N-limited environment that has been subject to high airborne N inputs over decades. We analysed N flows (in terms of ^{15}N tracer enrichment and recovery) to and between different ecosystem compartments, using an experiment that was established in a heathland ecosystem in 2007 (Friedrich et al., 2011b) and continued until 2015. We were particularly interested in the impact of the moss layer on the N cycle of the ecosystem, because this compartment is considered an important sink for incoming N (Proctor, 2008), but also might release N to other compartments, for example via downward transports in the soil profile. In this way, the moss layer could contribute to N enrichment in deeper soil horizons (Aldous, 2002; Proctor, 2008; Startsev et al., 2008). Moreover, we asked whether there are indications of N-saturation, or whether ecosystem-internal sinks for N or unknown pathways of N losses might counteract an accumulation of airborne N inputs. We hypothesised that (i) the moss layer constitutes a crucial component of the N cycle of the ecosystem, based on its capture-release-function for airborne N, (ii) the podzol-B-horizon serves as an important long-term sink for N, and (iii) the heathlands studied still have a high capacity to sequester pollutant N.

MATERIALS AND METHODS

Study Site and Experimental Design

Our study site is located in the Lüneburg Heath nature reserve in NW Germany (Lower Saxony, Schneverdingen, 53°15'N, 9°58'E, 105 m a.s.l.), the site of the largest complex of heathlands (about 5,000 ha) in NW Germany. The study sites is characterised by Pleistocene sandy deposits with a weak relief and nutrient-poor podzol soils. Topsoil-pH_{H2O} values range between 3.3

and 3.5 (Härdtle et al., 2006). The climate is of a humid sub-oceanic type (mean precipitation: 811 mm year⁻¹, annual mean temperature: 8.4°C; Niemeyer et al., 2005). Heathlands of the nature reserve are dominated by the dwarf shrub *Calluna* (cover >80%). The soil surface is covered by mosses (mean cover 93%), with *Hypnum judlandicum* and *Pleurozium schreberi* being the prevailing species. The current background atmospheric N deposition in the study area amounts to 23 kg ha⁻¹ year⁻¹ (von Oheimb et al., 2010 for more detailed information on the quantity of different N-forms deposited in NW Germany and in the study area see Gauger et al. (2000) and Niemeyer et al. (2005), respectively).

The present study made use of the experimental setup of Friedrich et al. (2011b) established in 2007. The experiment comprises a total of seven randomly selected plots, each of which was divided into two subplots (2 × 4 m in size). At the beginning of the experiment (June 2007), in one subplot (per plot) a ¹⁵N tracer was added in liquid form (pulse labeling with 100 mg ¹⁵NH₄¹⁵NO₃ (98 at.%) using a special spray bottle that allowed for a fine nebulisation of the liquid applied), and the corresponding subplot was used to determine the natural ¹⁵N abundance (these subplots are henceforth referred to as labelled and non-labelled subplots, respectively). Labelled and non-labelled subplots were separated by a buffer zone 1 m in width to avoid cross contamination after ¹⁵N tracer addition (for a detailed description of the ¹⁵N application procedure see Friedrich et al., 2011b).

All analyses performed during the present study refer to a time span of 7 years, with all samples collected between 2009 and 2015 (with the exception of one plot, which was sampled from 2008 to 2015, as well as litter and leachate samples; see description below). To present a complete overview of N flows and partitioning since the setup of the experiment in 2007, we included analyses of Friedrich et al. (2011b) in the present study (i.e., analyses from 2007 to 2008). In this way, we were able to document compartment-related shifts in ¹⁵N abundance, enrichment and recovery for a 9-year timespan (2007–2015).

N Allocation Patterns in Biomass and Soil

To quantify N allocation patterns in the biomass and soil we sampled the following ecosystem compartments annually: above-ground biomass of *Calluna* (current year and 1- to 2-year-old shoots), moss layer, and the soil O-, A- and B-horizons (with the latter two representing the albic and spodic horizons, respectively, both of which are typical of podzol soils; nomenclature according to the IUSS Working Group WRB, 2015) in labelled and non-labelled subplots in each August from 2008 to 2015, corresponding to the flowering period of *Calluna*. Twenty randomly chosen current year shoots as well as 1- to 2-year-old shoots of *Calluna* from the top of randomly chosen *Calluna* plants were cut with scissors and were pooled to one sample per subplot. Six randomly chosen squares of the moss layer (2 × 2 cm in size) were sampled and bulked to one sample per subplot. Organic soil samples (O-horizon; mean thickness: 2.7 cm) were cut with knives (six squares of 2 × 2 cm of the entire horizon depth per subplot) and separated from fresh

fallen litter. Mineral soil samples were taken from the A- and B-horizon with a soil auger (Pürckhauer, EcoTech, Bonn, Germany) from six randomly chosen spots per subplot and bulked to one sample each mean thickness of A- and B-horizons: 7.1 cm and 7.5 cm, respectively). The annual litter production of *Calluna* was quantified in labelled and non-labelled subplots by means of fine-mesh nets stretched under one randomly selected plant per subplot and sampled during winter between the end of February and the beginning of April (depending on the duration of snow cover) from 2008 to 2016.

To obtain an overview of the quantities of ¹⁵N accumulated in older *Calluna* twigs and stem wood (age >2 years) and *Calluna* below-ground biomass (i.e., roots), but to avoid an interference of plant growth in plots, these compartments were sampled only twice, in August 2014 and 2015. To this end, six randomly chosen stems of *Calluna* per subplot were cut with clippers at ground level (excluding current year and 1- to 2-year-old shoots). For root analysis, we sampled the O-horizon at six randomly selected sites per subplot (3 × 3 cm samples from across the entire organic layer).

Soil and root samples were stored in a freezer (−18°C) until analysis. Prior to analysis, above-ground biomass and soil samples were air-dried, and the latter one sorted and sieved (2 mm). Root samples were washed using a 0.2 mm sieve to remove soil particles and then dried (48 h at 80°C). All samples were ground with a mixer mill (MM 400, Retsch, Haan, Germany) and re-dried at 105°C for weighing. Total N contents and δ¹⁵N were determined with a continuous flow elemental analyser-isotopic ratio mass spectrometer (vario EL cube, Elementar, Hanau, Germany, coupled to an Isoprime IRMS, Isoprime Ltd., Cheadle Hulme, UK; note that total N analyses of soils hence included microbial biomass N).

N Losses via Leaching

N losses via leaching were quantified by means of four pairs of lysimeters, one of which was labelled with 100 mg ¹⁵NH₄¹⁵NO₃ m⁻² in 2007 (analogous to the labelling of subplots), and one which received a corresponding amount of deionised water (non-labelled lysimeter; for a detailed description of the setup of lysimeters see Friedrich et al., 2011b). Since the collection and analysis of leachate was very laborious and ¹⁵N leaching losses (in terms of inorganic N and DON) proved to be negligible at the beginning of the experiment, leaching losses of ¹⁵N were only quantified on 20 occasions in the period from June 2014 to May 2015 (weekly in June 2014, once every 2 weeks from July to August 2014 and monthly from September 2014 to June 2015). The leachate which accumulated between the sampling occasions was extracted with a temporarily connected vacuum pump. The quantity of leachate per lysimeter was recorded and a subsample of 500 ml was filtered and stored in a freezer (−18°C) for analysis. Samples were carefully defrosted and analysed with a UV/Vis scanning spectrophotometer (UV-3100PC, VWR, Radnor, PA, USA) using the method of Hagedorn and Schleppi (2000). In order to determine δ¹⁵N values, samples were prepared following the diffusion method of Sebilo et al. (2004) modified according to Friedrich et al. (2011b).

Calculation of N Pools and Leaching Losses

To calculate ^{15}N tracer recovery, the mass of N pools in each ecosystem compartment must be known. Therefore, the total N content of a compartment in labelled subplots was multiplied by its mean N pool mass. Pool masses for the moss and soil compartments were taken from Friedrich et al. (2011b). *Calluna* plants were 10–12 years old at the beginning of the experiment (i.e., early mature phase of life-history; Gimingham, 1972). The continuous increment of *Calluna* above-ground biomass (i.e., total above-ground biomass and total biomass of current year and 1- to 2-year-old shoots) was calculated on the basis of age-dependent growth rates of *Calluna* in the nature reserve, derived from our own measurements (based on biomass harvests at six randomly chosen sites adjacent to the experimental plots and comparisons with data from Friedrich et al., 2011b) in the Lüneburg Heath and complemented by literature data (Gimingham, 1972; Milne et al., 2002). *Calluna* below-ground biomass was also quantified in mature stands adjacent to our plots (at six randomly selected sites, 15 soil samples per site taken with a soil auger 8 cm in diameter and to a soil depth of 15 cm; following the procedure for root analyses described above).

N leaching losses were calculated from the total amount of leachate multiplied by the total N contents (N_{inorg} and DON) of the leachate from labelled lysimeters (Friedrich et al., 2011b).

Calculation of ^{15}N Abundance, ^{15}N Enrichment and ^{15}N Tracer Recovery

^{15}N abundance, ^{15}N enrichment and ^{15}N tracer recovery were calculated according to Fry (2006). ^{15}N abundance and the natural ^{15}N abundance in labelled and non-labelled subplots, respectively, are represented in the δ notation:

$$\delta^{15}\text{N} [\text{‰}] = \left(\frac{R_{\text{sample}}}{R_{\text{standard}}} - 1 \right) \cdot 1,000, \quad (1)$$

where R_{sample} and R_{standard} are the $^{15}\text{N}/^{14}\text{N}$ ratio of the sample and the N_2 of the atmosphere, which is used as standard, respectively.

^{15}N enrichment [%] represents the isotopic enrichment of a sample from a labelled subplot ($\delta^{15}\text{N}_{\text{sample}}$) compared to a sample in a non-labelled subplot ($\delta^{15}\text{N}_{\text{ref}}$):

$$^{15}\text{N}_{\text{enrichment}} [\text{‰}] = \frac{\delta^{15}\text{N}_{\text{sample}} - \delta^{15}\text{N}_{\text{ref}}}{\delta^{15}\text{N}_{\text{ref}} + 1,000} \cdot 1,000. \quad (2)$$

^{15}N recovery [g N m^{-2}] is the mass of ^{15}N tracer recovered in the different ecosystem compartments (biomass, soil horizons, and leachate) in labelled subplots and was calculated as follows:

$$^{15}\text{N}_{\text{recovery}} = m_{\text{pool}} \cdot \frac{\text{at.}\%^{15}\text{N}_{\text{pool}} - \text{at.}\%^{15}\text{N}_{\text{ref}}}{\text{at.}\%^{15}\text{N}_{\text{tracer}} - \text{at.}\%^{15}\text{N}_{\text{ref}}}, \quad (3)$$

where m_{pool} is the amount of N in a respective compartment [g N m^{-2}], $\text{at.}\%^{15}\text{N}_{\text{pool}}$ and $\text{at.}\%^{15}\text{N}_{\text{ref}}$ are the $\text{at.}\%^{15}\text{N}$ in the N pool of the samples (or in the leachate) of labelled and non-labelled subplots, respectively, and $\text{at.}\%^{15}\text{N}_{\text{tracer}}$ is the $\text{at.}\%^{15}\text{N}$ of the added ^{15}N tracer (Nadelhoffer et al., 2004).

Statistical Analysis

Differences between ^{15}N abundances of labelled and ^{15}N natural abundances from non-labelled subplots were tested by means of pairwise comparisons of $\delta^{15}\text{N}$ values of each compartment and for each sampling occasion using a Student's t -test ($\alpha = 0.05$). We applied generalised additive mixed models (GAMM) with a log link function and gamma distribution to assess the effects of ecosystem compartments and time (2007–2015) on ^{15}N enrichment. Ecosystem compartments were stratified into five main compartments: above-ground biomass of *Calluna* (including current year shoots, 1- to 2-year-old shoots, and current year litter), moss layer, and O-, A- and B-horizons of the soil. To address the skewed response of the ^{15}N enrichment data, a gamma probability distribution was preferred, because it retains the structure of the data while accounting for a heteroscedastic error structure (Zuur et al., 2014). We added a value of 0.001 to the ^{15}N enrichment data to ensure convergence of the fitting algorithm. The non-linear time effect was modelled using a cubic regression spline with four degrees of freedom to allow some complexity in the ^{15}N enrichment function, while avoiding over-fitting the data (Wood, 2006). We used plot as random factor to account for between-plot variability, and a first-order autocorrelation structure (AR-1) nested within plot to account for temporal correlation (Zuur et al., 2009). The resulting GAMM is:

$$EN_{ij} = \exp(\alpha + f(\text{Time}_{ij}) + \beta \text{Compartment}_{ij} + b_i) \quad (4)$$

where EN_{ij} is the ^{15}N enrichment of sample j in plot i and EN_{ij} is assumed to follow a Gamma distribution with a log link. α is the intercept, f is a smoothing function (cubic regression spline) of the time effect and β is a parametric coefficient of the compartment effect. b_i is a normally distributed random intercept for the plot effect accounting for any dependency between samples from the same plot. The residuals at time s are modelled as a function of the residuals of time $s-1$ along with noise (as $\varepsilon_s = \rho \varepsilon_{s-1} + \eta_s$, in which ρ is the temporal correlation parameter to be estimated; Zuur et al., 2009). The importance of variables for ^{15}N enrichments was assessed by fitting various candidate models accounting for the main effects and their interaction (as $f(\text{Time}_{ij}) \times \text{Compartment}_{ij}$) using maximum likelihood estimation (ML). Model selection was based on the Akaike Information Criterion (AIC) and the model with the smallest AIC and highest Akaike weights (w_i) was identified as the minimum-adequate model (Burnham and Anderson, 2002). Parameter estimates of the minimum-adequate model were based on the restricted maximum likelihood (REML) method (Zuur et al., 2009). Model assumptions were visually assessed following Zuur et al. (2010). We found no trends in the residuals. To avoid biased parameter estimates, we omitted extreme values (outliers, <2% of the data) for the analyses. Outliers were visually identified by Cleveland dotplots (Zuur et al., 2010). All analyses were conducted in R 3.3.1 (<http://www.R-project.org>) with the packages nlme (Pinheiro et al., 2016), mgcv (Wood, 2006), and MuMIn (Bartón, 2016).

TABLE 1 | ^{15}N abundance ($\delta^{15}\text{N}$ in ‰) in non-labelled (Ref.) and labelled subplots (^{15}N) of the ecosystem compartments analysed (means, 1 SE (standard error of the mean) in brackets, $n = 7$ samples) during the study period 2007–2015.

Year	Current year shoots		1- to 2-year-old shoots		Calluna older than 2 years		Calluna roots		Current year litter		Moss layer		O-horizon		A-horizon		B-horizon	
	Ref.	^{15}N	Ref.	^{15}N	Ref.	^{15}N	Ref.	^{15}N	Ref.	^{15}N	Ref.	^{15}N	Ref.	^{15}N	Ref.	^{15}N	Ref.	^{15}N
2007	-5.67 (0.51)	129.30 (25.56)	-5.92 (0.47)	110.42 (23.28)					-6.24 (0.16)	54.66 (25.18)	-6.19 (0.22)	735.74 (106.17)	-4.53 (0.35)	19.74 (4.22)	5.33 (0.40)	8.16 (0.48)	8.71 (0.43)	8.74 (0.69)
2008	-6.54 (0.69)	65.00 (10.43)	-7.67 (0.63)	54.54 (11.40)					-5.40 (0.80)	37.55 (9.70)	-5.93 (0.38)	535.65 (55.71)	-4.79 (0.44)	28.60 (6.03)	4.26 (1.11)	9.43 (0.81)	7.16 (1.32)	8.90 (0.76)
2009	-6.90 (0.52)	47.15 (5.53)	-7.68 (0.57)	41.55 (6.46)					-7.66 (0.61)	40.14 (11.39)	-6.60 (0.31)	432.63 (63.96)	-4.85 (0.48)	30.62 (4.92)	4.49 (1.23)	8.50 (0.95)	6.73 (1.36)	8.30 (1.00)
2010	-6.85 (0.56)	55.23 (7.56)	-7.96 (0.51)	45.77 (6.13)					-8.02 (0.48)	38.11 (8.71)	-6.11 (0.34)	338.70 (32.94)	-3.99 (0.59)	18.57 (2.91)	4.76 (1.02)	9.26 (1.39)	7.68 (1.11)	9.82 (1.03)
2011	-6.02 (0.46)	51.47 (5.23)	-7.50 (0.42)	47.30 (5.14)					-7.60 (0.47)	36.68 (7.32)	-6.32 (0.49)	177.92 (18.88)	-4.50 (0.42)	25.12 (2.17)	4.69 (1.12)	9.26 (0.64)	7.21 (1.25)	8.78 (1.01)
2012	-6.30 (0.51)	66.69 (8.60)	-7.07 (0.46)	59.44 (7.55)					-6.64 (0.64)	31.11 (6.29)	-6.79 (0.18)	141.69 (13.06)	-3.52 (0.53)	38.77 (7.28)	4.14 (1.18)	10.55 (0.25)	6.62 (1.28)	10.24 (0.46)
2013	-6.55 (0.50)	49.12 (6.05)	-7.58 (0.44)	40.61 (5.05)					-6.62 (0.52)	29.57 (5.95)	-6.75 (0.35)	109.67 (9.92)	-3.58 (0.63)	29.41 (2.84)	3.91 (1.14)	13.97 (2.08)	7.14 (1.10)	10.22 (0.30)
2014	-6.06 (0.56)	36.13 (5.51)	-6.72 (0.50)	31.92 (4.72)	-7.00 (0.46)	38.44 (4.85)	-5.34 (0.61)	60.77 (10.48)	-6.78 (0.43)	24.09 (5.36)	-6.74 (0.21)	50.42 (7.22)	-3.82 (0.53)	46.60 (7.72)	3.53 (1.01)	15.63 (3.97)	6.52 (1.26)	10.35 (0.90)
2015	-6.30 (0.53)	33.70 (6.35)	-7.28 (0.41)	31.90 (5.41)	-7.40 (0.47)	34.10 (5.90)	-5.39 (0.52)	61.74 (5.87)	-8.60 (0.39)	18.33 (4.69)	-6.63 (0.24)	40.02 (6.87)	-3.78 (0.55)	59.32 (6.17)	5.17 (1.43)	10.38 (0.54)	6.13 (1.08)	9.48 (0.47)

Significantly ($P < 0.05$) enhanced $\delta^{15}\text{N}$ values in labelled subplots are indicated in bold.

RESULTS

^{15}N Abundances and ^{15}N Enrichment in Biomass and Soil

$\delta^{15}\text{N}$ values (natural abundance) varied between -8.0 and $+8.7\text{‰}$ across compartments and were lowest in *Calluna* biomass (range between -8.0 and -5.3‰). The moss layer showed $\delta^{15}\text{N}$ values between -6.8 and -5.9‰ . In the soils, $\delta^{15}\text{N}$ values increased continuously with soil depth (organic layer: between -4.9 and -3.5‰ , A-horizon: between 3.5 and 5.3‰ and B-horizon: between 6.1 and 8.7‰ ; **Table 1**).

As was to be expected, most compartments showed a significant ($P < 0.05$) increase in $\delta^{15}\text{N}$ values (enrichment) after tracer addition (exceptions were the B-horizon, some measurements of the A-horizon and the litter samples in the first year of the experiment). The highest enrichment values were found for the moss layer and *Calluna* shoots (**Table 1**).

The direction and magnitude of temporal changes in ^{15}N enrichment distinctly varied among compartments (**Figure 1**), as indicated by the strong support for interacting effects of time and compartment (Akaike model weight of 1.0; **Table 2**). 86 % of the variation in ^{15}N enrichment was explained by the interaction between time and compartment, whereas compartment or time effects explained 45 and 7% of the variance, respectively (**Table 2**). The partial effects of the above-ground biomass of *Calluna* and the moss layer on ^{15}N enrichment showed a significant decrease over time ($P < 0.001$ for both compartments;

Table 3; **Figure 1**), while a significant increase of ^{15}N enrichment was observed in the soil (O- and B-horizon: $P < 0.001$; A-horizon: $P < 0.01$; **Table 3**; **Figure 1**). ^{15}N enrichment in the A-horizon also increased within the course of the experiment, but decreased in 2015 (**Table 1**).

^{15}N Tracer Recovery

Total ^{15}N recovery was highest in the first year (about 93%); it then (almost linearly) decreased until 2010 to a value of about 54% (**Figure 2**; **Table 4**). Interestingly, recovery rates remained largely constant in subsequent years and achieved values between 49 and 54% until 2015. The recovery rates calculated increase by

TABLE 2 | Model selection statistics for four candidate models describing ^{15}N enrichment as a function of ecosystem compartment (above-ground biomass, moss layer, O-, A-, and B-horizons) and time (2007–2015).

Model	Fixed effects	ΔAIC	w_i	$R^2(\text{adj.})$
1	Time	476.80	0	0.07
2	Ecosystem compartment	227.95	0	0.45
3	Time + Ecosystem compartment	147.60	0	0.60
4	Time \times Ecosystem compartment	0.00	1	0.86

The minimum-adequate model is highlighted in bold. ΔAIC is the difference in AIC (Akaike Information Criterion) with respect to the best-fitting model (lowest value of AIC). The Akaike weight (w_i) is the relative likelihood of each model being the best-fitting model, given the complete set of candidate models. R^2 is the variance explained by the model.





FIGURE 1 | Estimated smoother for time-related shifts in ^{15}N enrichment in the ecosystem compartments analysed. The solid line corresponds to fitted relationships of generalised additive mixed models, with shaded areas showing the 95% confidence interval range. The y-axis indicates the contribution of the smoother (effect, ‰) to the fitted values of ^{15}N enrichment (see Equation 4 and **Table 3**; $n = 7$ per compartment and year).

TABLE 3 | Minimum-adequate generalised mixed-effects model (GAMM) for the effects of ecosystem compartment and time on ^{15}N enrichment.

	Estimate	SE	t/F-value	edf	P-value
Parametric coefficients:					
Intercept	3.941	0.09	41.84		<0.001
Compartment (moss layer)	1.358	0.08	17.80		<0.001
Compartment (O-horizon)	-0.349	0.08	-4.47		<0.001
Compartment (A-horizon)	-2.321	0.08	-29.76		<0.001
Compartment (B-horizon)	-3.364	0.08	-43.41		<0.001
Smooth terms:					
$f(\text{Time}) * \text{Compartment (above-ground biomass)}$			20.61	3.64	<0.001
$f(\text{Time}) * \text{Compartment (moss layer)}$			214.69	1.00	<0.001
$f(\text{Time}) * \text{Compartment (O-horizon)}$			12.41	1.90	<0.001
$f(\text{Time}) * \text{Compartment (A-horizon)}$			4.03	3.02	0.006
$f(\text{Time}) * \text{Compartment (B-horizon)}$			16.36	2.97	<0.001
Variance components:					
SD (plot)	0.226				
SD (residuals)	0.439				
ϕ (residuals)	0.197				

The intercept corresponds to the above-ground biomass compartment. The estimated degrees of freedom (edf) indicate the amount of smoothing. SE, Standard error (of the mean); SD, Standard deviation; ϕ , Estimated temporal correlation of the residuals (AR-1).



about 7% if the compartments which were not sampled annually are taken into account: “*Calluna* biomass older than 2 years” (about 2.5%), and “*Calluna* roots” (about 4.2%; means across sampling years; Table 4).

The moss layer proved to be the most important short-term sink and longer-term source for applied ^{15}N , and showed a distinct decrease in recovery rates within the first 5 years (from 66% in 2007 to 14% in 2011; Figures 2, 3). This loss of ^{15}N from the moss layer was not compensated for by a corresponding increase in recovery rates in any other compartment (Figure 3). We did find increasing recovery rates in the O- and A-horizons in the course of the experiment (e.g., from 18 to 40% for the O-horizon). This increase corresponded to about 50% of the total ^{15}N losses from the moss layer between 2007 and 2015 (Figure 2). For the B-horizon, we found a slight but continuous increase in ^{15}N recovery, with a maximum of 2.1% in the last year of the experiment.

Recovery rates in both the current year and 1- to 2-year-old shoots of *Calluna* were negligibly low in all study years, with values below 2%, and in most years even below 1% (Figures 2, 3, Table 4). Accordingly, we also found low recovery rates for the annual litter of *Calluna*, which amounted to $\leq 1\%$. In addition, leaching losses of ^{15}N proved to be very low, as indicated by an annual recovery of 0.03% of the total ^{15}N applied. Thus, leaching losses were even lower than at the beginning of the experiment (0.05% in 2007).

DISCUSSION

Decadal-Scale Allocation Patterns of N

Our findings support our first hypothesis that the moss layer acts as both an important short-term sink and a longer-term source of captured ^{15}N tracer. The moss layer thus represents a crucial link between airborne N inputs and the allocation of N to the heathland's N cycle (in term of its capture-release function). The moss layer's ability to store incoming N in the short term may be related to the fact that mosses represent ectohydric plants that

TABLE 4 | ^{15}N enrichment [%] and ^{15}N recovery [% of the total ^{15}N applied] of the sampled ecosystem compartments (means, 1 SE in brackets) during the study period 2007–2015.

Year	Current year shoots	1- to 2-year-old shoots	<i>Calluna</i> shoots/woody parts older than 2 years	<i>Calluna</i> roots	Current year litter	Moss layer	O-horizon	A-horizon	B-horizon	total
^{15}N ENRICHMENT [%]										
2007	135.72 (25.42)	117.00 (23.10)			61.27 (25.18)	746.60 (106.95)	24.38 (4.03)	2.81 (0.34)	0.46 (0.22)	
2008	72.03 (10.72)	62.69 (11.54)			43.21 (10.10)	544.87 (56.34)	33.55 (5.98)	5.15 (1.00)	1.74 (0.64)	
2009	54.44 (5.91)	49.63 (6.92)			48.18 (11.67)	442.26 (64.83)	35.61 (5.78)	3.99 (0.50)	1.66 (0.62)	
2010	62.54 (8.17)	54.18 (6.62)			46.51 (9.01)	346.97 (33.39)	22.59 (3.58)	4.72 (1.11)	2.19 (0.63)	
2011	57.85 (5.64)	55.23 (5.48)			44.63 (7.62)	185.41 (19.09)	29.86 (2.47)	4.55 (1.01)	1.57 (0.43)	
2012	73.48 (9.11)	67.00 (7.97)			38.03 (6.85)	149.50 (13.27)	42.43 (7.18)	6.39 (1.05)	2.64 (0.68)	
2013	56.05 (6.37)	48.56 (5.36)			36.45 (6.44)	117.21 (10.07)	33.12 (3.44)	6.98 (0.44)	2.02 (0.58)	
2014	42.47 (5.88)	38.91 (4.97)	45.77 (5.16)	66.47 (10.69)	31.09 (5.76)	57.55 (7.35)	50.61 (7.69)	7.24 (1.10)	1.89 (0.43)	
2015	40.26 (6.61)	39.47 (5.64)	41.81 (6.21)	67.49 (5.73)	27.18	46.97 (7.11)	63.35 (6.40)	5.19 (1.07)	3.34 (0.78)	
^{15}N RECOVERY [%]										
2007	1.69 (0.41)	0.89 (0.21)			0.56 (0.21)	66.38 (10.30)	17.80 (2.98)	5.01 (1.36)	0.43 (0.24)	92.75
2008	0.96 (0.14)	0.44 (0.08)			0.62 (0.14)	42.08 (5.92)	22.08 (3.41)	7.05 (1.46)	0.89 (0.18)	74.13
2009	0.74 (0.08)	0.40 (0.05)			0.74 (0.16)	32.48 (5.42)	22.55 (3.10)	6.28 (1.66)	0.96 (0.29)	64.15
2010	1.04 (0.14)	0.48 (0.07)			0.88 (0.17)	24.44 (2.73)	17.11 (2.74)	8.08 (2.68)	1.62 (0.59)	53.65
2011	0.94 (0.10)	0.54 (0.06)			1.25 (0.22)	14.42 (1.37)	24.27 (2.59)	6.91 (1.93)	0.94 (0.19)	49.26
2012	1.24 (0.14)	0.62 (0.09)			0.94 (0.16)	10.76 (1.35)	30.05 (3.63)	8.43 (1.94)	1.95 (0.48)	53.99
2013	0.91 (0.10)	0.42 (0.06)			1.04 (0.20)	8.85 (0.81)	24.52 (3.14)	12.73 (2.55)	1.63 (0.37)	50.11
2014	0.74 (0.10)	0.33 (0.05)	2.49 (0.25)	3.60 (0.57)	0.80 (0.14)	3.97 (0.52)	36.11 (6.02)	11.02 (2.99)	1.39 (0.58)	54.37 60.47*
2015	0.83 (0.13)	0.41 (0.07)	2.59 (0.54)	4.86 (0.77)	0.93 (0.15)	2.91 (0.46)	39.89 (4.89)	6.61 (1.46)	2.08 (0.55)	53.65 61.10*

*Total recovery of all compartments, including compartments not sampled annually (*Calluna* older than 2 years and roots).

absorb dissolved nutrients at the cell surface (Proctor, 2008). The decadal-scale availability and release of N could be attributable to a longer-term sequestration of N due to N assimilation by mosses and the microbial community associated with the moss phytomass (Gordon et al., 2001; Tye et al., 2005).

An important finding in terms of the N balance of the ecosystem was that, during the first 4 years of the experiment, ^{15}N losses from the moss layer were not compensated for

by a corresponding increase in recovery rates in any other compartment. These losses accounted for more than 30% of the total ^{15}N applied and suggest that N losses from the moss layer in gaseous forms are very likely (Startsev and Lieffers, 2007; for a further discussion of this aspect see paragraph ‘Leaching losses and putative flows’ below). In contrast, the largely constant recovery rates found from 2010 onwards point to a conservative N cycling with marginal ^{15}N losses from



Larsen et al. (2012) in a subarctic heathland, where 65% of the applied ^{15}N was sequestered by soil microbes. In this context, it is important to note that high immobilisation rates by soil microbes are also indicative of a low N saturation level (Curtis et al., 2005).

Leaching Losses and Putative Flows

^{15}N losses via leaching proved to be negligibly low. This is an important indication of conservative N cycles (Tye et al., 2005) and contradicts the notion that the heathlands of the study area have achieved an advanced stage of N saturation (Aber et al., 1998), despite decades of N inputs beyond critical load thresholds. Obviously, the sites analysed still have a high capacity to sequester pollutant N (Choudhary et al., 2016), suggesting that a constant proportion of incoming N enters the heathland's N cycle and contributes to a slow but steady increase in the N level (De Schrijver et al., 2011). This confirms our third hypothesis.

In contrast to the observed accumulation of N, the first 4 years of the experiment (2007–2010) also exhibited a considerable loss of ^{15}N . Tracer mass balances suggest that this loss was attributable to a leak of N from the moss layer in gaseous forms (Startsev and Lieffers, 2007). High N losses from bryophyte layers due to denitrification have been found for different species and ecosystems, and increase with increasing humidity (e.g., with water saturation of the moss layer after precipitation events), temperature, and N deposition (Laverman et al., 2000; Opelt and Berg, 2004; Lenhart et al., 2015). As a consequence, gaseous N losses from moss carpets are considered to have significant impacts on ecosystem N budgets (Fang et al., 2015), and also contribute to global nitrous oxide emissions from terrestrial ecosystems (Lenhart et al., 2015). For example, Calvo-Fernández et al. (2015) found high N losses as a result of denitrification in Cantabrian Mountain heathlands, when soils were water-saturated during winter. Fang et al. (2015) concluded that the effects of denitrification losses on ecosystem N balances are often underestimated. NH_3 volatilisation might also have contributed to ^{15}N losses from the moss layer (Mahendrappa and Ogden, 1973; Startsev et al., 2008). Further mechanisms of N losses could be lateral N transport by mycorrhiza or soil fauna. However, since the natural ^{15}N abundances in non-labelled subplots remained unchanged in the course of the experiment, we consider these losses (including conceivable lateral transports due to surface run-off) to be of minor importance. Low lateral transports also might be related to high absorption rates of ammonium, which is the prevailing N_{inorg} form at podzol sites with strongly acidic soil condition (cf. study site description). Due to the avoidance of destructive sampling, N contents of roots below a soil depth of 15 cm were not quantified. Since *Calluna* is considered a shallow-rooted plant (with more than 90% of root biomass being located in the humus horizons; Gimingham, 1972), and CN

ratios of deep rooting woody roots are exceptionally high (>120 ; own unpublished data), we consider the N pool related to this compartment to be very low. We also rule out a significant effect of N deposition on ^{15}N recovery patterns, since N deposition is evenly distributed across the study area, and thus between-plot differences in N deposition are low (including differences in N forms and isotope signatures; cf. deposition measurements of Niemeyer et al., 2005).

In conclusion, we found evidence that about 60% of the added ^{15}N remained in the N cycle of the ecosystem at least for about one decade. In this context moss layers may play a crucial functional role for the N cycle of ecosystems: On the one hand, moss layers may contribute to high N losses by fostering denitrification processes (particularly under oceanic to sub-oceanic climates), and thus may decelerate N eutrophication processes in ecosystems. On the other hand, mosses represent a considerable short-term sink for incoming N which is released to the N cycle of an ecosystem in the long term. As a result, a considerable proportion of airborne N inputs can accumulate in different ecosystem compartments. Since the O-horizon proved to be an important long-term sink, it is likely that N captured by this horizon remains bio-available in the long term.

This has important implications for both critical load estimates and the development of management strategies to counteract airborne N inputs. A continuous accumulation of bio-available N (even under low N inputs) would implicate that critical load revisions need to account for cumulative effects of (low) N additions into ecosystems (Payne et al., 2011; Phoenix et al., 2012; Bähring et al., 2017). Mitigation strategies (in terms of management) should comprise high-intensity measures (e.g., choppers and sod-cutting; Härdtle et al., 2006; Frouz et al., 2009) as an important means to achieve balanced N budgets in the long term, because only these measures affect upper soil horizons with high contents of accumulated N.

AUTHOR CONTRIBUTIONS

UF, WH, and GvO conceived of the study. All authors contributed to the field and lab work, the analysis and interpretation of data, and the writing of the manuscript. All authors approved the final version of the manuscript.

ACKNOWLEDGMENTS

We would like to thank the Verein Naturschutzpark e.V. for permission to conduct this study in the Lüneburg Heath nature reserve.

REFERENCES

- Aber, J., McDowell, W., Nadelhoffer, K., Magill, A., Berntson, G., Kamakea, M., et al. (1998). Nitrogen saturation in temperate forest ecosystems-Hypotheses revisited. *Bioscience* 48, 921–934. doi: 10.2307/13296
- Aerts, R. (1999). Interspecific competition in natural plant communities: mechanisms, trade-offs and plant-soil feedbacks. *J. Exp. Bot.* 50, 29–37. doi: 10.1093/jxb/50.330.29
- Aldous, A. R. (2002). Nitrogen translocation in *Sphagnum* mosses: effects of atmospheric nitrogen deposition. *New Phytol.* 156, 241–253. doi: 10.1046/j.1469-8137.2002.00518.x

- Alonso, I., Hartley, S. E., and Thurlow, M. (2001). Competition between heather and grasses on Scottish moorlands: interacting effects of nutrient enrichment and grazing regime. *J. Vegetation Sci.* 12, 249–260. doi: 10.2307/3236609
- Andresen, L. C., Jonasson, S., Ström, L., and Michelsen, A. (2008). Uptake of pulse injected nitrogen by soil microbes and mycorrhizal and non-mycorrhizal plants in a species-diverse subarctic heath ecosystem. *Plant Soil* 313, 283–295. doi: 10.1007/s11104-008-9700-7
- Bähring, A., Fichtner, A., Ibe, K., Schütze, G., Temperton, V. M., von Oheimb, G., et al. (2017). Ecosystem functions as indicators for heathland responses to nitrogen fertilization. *Ecol. Indic.* 72, 183–195. doi: 10.1016/j.ecolind.2016.08.013
- Bartón, K. (2016). *MuMIn: Multi-Model Inference*. R package version 1.15.6. Available online at: <https://CRAN.R-project.org/package=MumIn>
- Bates, J. W. (2008). “Mineral nutrition and substratum ecology,” in *Bryophyte Ecology*, 2nd Edn., eds B. Goffinet and A. J. Shaw (Cambridge: Cambridge University Press), 299–356.
- Berdowski, J. J. M., and Zeilinga, R. (1987). Transition from heathland to grassland: damaging effects of the heather beetle. *J. Ecol.* 75, 159–175. doi: 10.2307/2260542
- Berendse, F. (1990). Organic matter accumulation and nitrogen mineralization during secondary succession in heathland ecosystems. *J. Ecol.* 78, 413–427. doi: 10.2307/2261121
- Bobbink, R., Ashmore, M., Braun, S., Flückiger, W., and van den Wyngaert, I. J. (2002). *Empirical Nitrogen Critical Loads for Natural and Semi-natural Ecosystems: 2002 Update* (Berne: Swiss Agency for Environment, Forest and Landscape SAEFL), 1–128.
- Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., et al. (2010). “Review and revision of empirical critical loads and dose-response relationships,” *Proceedings of an Expert Workshop. Noordwijkerhout*, 23–25 June 2010, eds R. Bobbink and J.-P. Hettelingh (Noordwijkerhout), 246.
- Brady, N. C., and Weil, R. R. (2008). *The Nature and Properties of Soils*, 14th Edn. Upper Saddle River, NJ: Pearson/Prentice Hall.
- Burnham, K. P., and Anderson, D. R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. New York, NY: Springer Science+Business Media.
- Calvo-Fernández, J., Marcos, E., Calvo, L., and Härdtle, W. (2015). Allocation patterns of airborne nitrogen in mountainous heathlands – A ^{15}N tracer study in the Cantabrian Mountains (NW Spain). *Ecol. Eng.* 84, 128–135. doi: 10.1016/j.ecoleng.2015.07.027
- Carroll, J. A., Caporn, S. J. M., Cawley, L., Read, D. J., and Lee, J. A. (1999). The effect of increased deposition of atmospheric nitrogen on *Calluna vulgaris* in upland Britain. *New Phytol.* 141, 423–431. doi: 10.1046/j.1469-8137.1999.00358.x
- Choudhary, S., Blaud, A., Osborn, A. M., Press, M. C., and Phoenix, G. K. (2016). Nitrogen accumulation and partitioning in a High Arctic tundra ecosystem from extreme atmospheric N deposition events. *Sci. Total Environ.* 554, 303–310. doi: 10.1016/j.scitotenv.2016.02.155
- Curtis, C. J., Emmett, B. A., Grant, H., Kernan, M., Reynolds, B., and Shilland, E. (2005). Nitrogen saturation in UK moorlands: the critical role of bryophytes and lichens in determining retention of atmospheric N deposition. *J. Appl. Ecol.* 42, 507–517. doi: 10.1111/j.1365-2664.2005.01029.x
- De Graaf, M. C. C., Bobbink, R., Smits, N. A. C., van Diggelen, R., and Roelofs, J. G. M. (2009). Biodiversity, vegetation gradients and key biogeochemical processes in the heathland landscape. *Biol. Conserv.* 142, 2191–2201. doi: 10.1016/j.biocon.2009.04.020
- De Schrijver, A., de Frenne, P., Ampoorter, E., van Nevel, L., Demey, A., Wuyts, K., et al. (2011). Cumulative nitrogen input drives species loss in terrestrial ecosystems. *Glob. Ecol. Biogeogr.* 20, 803–816. doi: 10.1111/j.1466-8238.2011.00652.x
- Fagúndez, J. (2012). Heathlands confronting global change: drivers of biodiversity loss from past to future scenarios. *Ann. Bot.* 111, 151–172. doi: 10.1093/aob/mcs257
- Fagúndez, J., and Izco, J. (2016). Spatial analysis of heath toponymy in relation to present-day heathland distribution. *Int. J. Geogr. Inform. Sci.* 30, 51–60. doi: 10.1080/13658816.2015.1017729
- Fang, Y. T., Koba, K., Makabe, A., Takahashi, C., Zhu, W. X., Hayashi, T., et al. (2015). Microbial denitrification dominates nitrate losses from forest ecosystems. *Proc. Natl. Acad. Sci. U.S.A.* 112, 1470–1474. doi: 10.1073/pnas.1416776112
- Friedrich, U., von Oheimb, G., Dziedek, C., Kriebitzsch, W. U., Selbmann, K., and Härdtle, W. (2011a). Mechanisms of purple moor-grass (*Molinia caerulea*) encroachment in dry heathland ecosystems with chronic nitrogen inputs. *Environ. Pollut.* 159, 3553–3559. doi: 10.1016/j.envpol.2011.08.010
- Friedrich, U., Falk, K., Bahlmann, E., Marquardt, T., Meyer, H., Niemeyer, T., et al. (2011b). Fate of airborne nitrogen in heathland ecosystems: a ^{15}N tracer study. *Glob. Chang. Biol.* 17, 1549–1559. doi: 10.1111/j.1365-2486.2010.02322.x
- Frouz, J., van Diggelen, R., Pizl, V., Stary, J., Hanel, L., Tajovsky, K., et al. (2009). The effect of topsoil removal in restored heathland on soil fauna, topsoil microstructure, and cellulose decomposition: implications for ecosystem restoration. *Biodivers. Conserv.* 18, 3963–3978. doi: 10.1007/s10531-009-9692-5
- Fry, B. (2006). *Stable Isotope Ecology*. New York, NY: Springer Science+Business Media.
- Gabet, E. J., Reichmann, O. J., and Seabloom, E. W. (2003). The effects of bioturbation on soil processes and sediment transport. *Ann. Rev. Earth Planet. Sci.* 31, 249–273. doi: 10.1146/annurev.earth.31.100901.141314
- Galloway, J. N., and Cowling, E. B. (2002). Reactive nitrogen and the world: 200 years of change. *Ambio* 31, 64–71. doi: 10.1579/0044-7447-31.2.64
- Gauger, T., Köble, R., and Anshelm, F. (2000). *Kritische Luftschadstoffkonzentrationen und Eintragsraten sowie ihre Überschreitung für Wald und Agrarökosysteme sowie naturnahe waldfreie Ökosysteme*. Federal Environmental Agency Report No. 29785079, (Berlin).
- Gerzabek, M. H., Haberhauer, G., Stemmer, M., Klepsch, S., and Haunold, E. (2004). Long-term behaviour of ^{15}N in an alpine grassland ecosystem. *Biogeochemistry* 70, 59–69. doi: 10.1023/B:BIOG.0000049336.84556.62
- Gimingham, C. H. (1972). *Ecology of Heathlands*. London: Chapman and Hall Ltd.
- Gordon, C., Wynn, J. M., and Woodin, S. J. (2001). Impacts of increased nitrogen supply on high Arctic heath: the importance of bryophytes and phosphorus availability. *New Phytol.* 149, 461–471. doi: 10.1046/j.1469-8137.2001.00053.x
- Gurmesa, G. A., Lu, X., Gundersen, P., Mao, Q., Zhou, K., Fang, Y., et al. (2016). High retention of ^{15}N -labeled nitrogen deposition in a nitrogen saturated old-growth tropical forest. *Glob. Chang. Biol.* 22, 3608–3620. doi: 10.1111/gcb.13327
- Hagedorn, F., Maurer, S., Bucher, J. B., and Siegwolf, R. T. W. (2005). Immobilization, stabilization and remobilization of nitrogen in forest soils at elevated CO_2 : A ^{15}N and ^{13}C tracer study. *Glob. Chang. Biol.* 11, 1816–1827. doi: 10.1111/j.1365-2486.2005.01041.x
- Hagedorn, F., and Schleppe, P. (2000). Determination of total dissolved nitrogen by persulfate oxidation. *J. Plant Nutr. Soil Sci.* 163, 81–82. doi: 10.1002/(SICI)1522-2624(200002)163:1<81::AID-JPLN81>3.0.CO;2-1
- Härdtle, W., Niemeyer, M., Niemeyer, T., Assmann, T., and Fottner, S. (2006). Can management compensate for atmospheric nutrient deposition in heathland ecosystems? *J. Appl. Ecol.* 43, 759–769. doi: 10.1111/j.1365-2664.2006.01195.x
- IUSS Working Group WRB (2015). *World Reference Base for Soil Resources 2014, Updated 2015. International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*, Vol. 106. (Rome: FAO).
- Jonasson, S., Michelsen, A., Schmidt, I. K., Nielsen, E. V., and Callaghan, T. V. (1996). Microbial biomass C, N and P in two arctic soils and responses to addition of NPK fertilizer and sugar: implications for plant nutrient uptake. *Oecologia* 106, 507–515. doi: 10.1007/BF00329709
- Kleinebecker, T., Schmidt, S. R., Fritz, C., Smolders, A. J. P., and Hölzel, N. (2009). Prediction of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in plant tissues with near-infrared reflectance spectroscopy. *New Phytol.* 184, 732–739. doi: 10.1111/j.1469-8137.2009.02995.x
- Kristensen, H. L. (2001). High immobilization of NH_4^+ in Danish heath soil related to succession, soil and nutrients: implications for critical loads of N. *Water Air Soil Pollut.* 1, 211–230. doi: 10.1023/A:1011588007257
- Kristensen, H. L., and McCarty, G. W. (1999). Mineralization and immobilization of nitrogen in heath soil under intact *Calluna*, after heather beetle infestation and nitrogen fertilization. *Appl. Soil Ecol.* 13, 187–198. doi: 10.1016/S0929-1393(99)00036-0
- Larsen, K. S., Michelsen, A., Jonasson, S., Beier, C., and Grogan, P. (2012). Nitrogen uptake during fall, winter and spring differs among plant functional groups in a subarctic heath ecosystem. *Ecosystems* 15, 927–939. doi: 10.1007/s10021-012-9555-x
- Laverman, A. M., Zoomer, H. R., Engelbrecht, D., Berg, M. P., van Straalen, N. M., van Verseveld, H. W., et al. (2000). Soil layer-specific variability in net

- nitrification and denitrification in an acidic coniferous forest. *Biol. Fertil. Soils* 32, 427–434. doi: 10.1007/s003740000274
- Lenhart, K., Werber, B., Elbert, W., Steinkamp, J., Clough, T., Crutzen, P., et al. (2015). Nitrous oxide and methane emissions from cryptogamic covers. *Glob. Chang. Biol.* 21, 3889–3900. doi: 10.1111/gcb.12995
- Mahendrapa, M. K., and Ogden, E. D. (1973). Patterns of ammonia volatilization from forest soil. *Plant Soil* 38, 257–265. doi: 10.1007/BF00779010
- Maskell, L. C., Smart, S. M., Bullock, J. M., Thompson, K., and Stevens, C. J. (2010). Nitrogen deposition causes widespread loss of species richness in British habitats. *Glob. Chang. Biol.* 16, 671–679. doi: 10.1111/j.1365-2486.2009.02022.x
- Milne, J. A., Pakeman, R. J., Kirkham, F. W., Jones, I. P., and Hossell, J. E. (2002). Biomass production of upland vegetation types in England and Wales. *Grass Forage Sci.* 57, 373–388. doi: 10.1046/j.1365-2494.2002.00339.x
- Nadelhoffer, K. J., Colman, B. P., Currie, W. S., Magill, A., and Aber, J. D. (2004). Decadal-scale fates of ^{15}N tracers added to oak and pine stands under ambient and elevated N inputs at the Harvard Forest (USA). *For. Ecol. Manage.* 196, 89–107. doi: 10.1016/j.foreco.2004.03.014
- Nielsen, K. E., Hansen, B., Ladekarl, U. L., and Nørnberg, P. (2000). Effects of N-deposition on ion trapping by B-horizons of Danish heathlands. *Plant Soil* 223, 265–276. doi: 10.1023/A:1004853802637
- Niemeyer, T., Niemeyer, M., Mohamed, A., Fottner, S., and Hårdtle, W. (2005). Impact of prescribed burning on the nutrient balance of heathlands with particular reference to nitrogen and phosphorus. *Appl. Vegetat. Sci.* 8, 183–192. doi: 10.1111/j.1654-109X.2005.tb00644.x
- Nordin, A., Schmidt, I. K., and Shaver, G. R. (2004). Nitrogen uptake by arctic soil microbes and plants in relation to soil nitrogen supply. *Ecology* 85, 955–962. doi: 10.1890/03-0084
- Opelt, K., and Berg, G. (2004). Diversity and antagonistic potential of bacteria associated with bryophytes from nutrient-poor habitats of the Baltic Sea coast. *Appl. Environ. Microbiol.* 70, 6569–6579. doi: 10.1128/AEM.70.11.6569-6579.2004
- Payne, R. J., Stevens, C. J., Dise, N. B., Gowing, D. J., Pilkington, M. G., Phoenix, G. K., et al. (2011). Impacts of atmospheric pollution on the plant communities of British acid grasslands. *Environ. Pollut.* 159, 2602–2608. doi: 10.1016/j.envpol.2011.06.009
- Phoenix, G. K., Emmett, B. A., Britton, A. J., Caporn, S. J. M., Dise, N. B., Helliwell, R., et al. (2012). Impacts of atmospheric nitrogen deposition: responses of multiple plant and soil parameters across contrasting ecosystems in long-term field experiments. *Glob. Chang. Biol.* 18, 1197–1215. doi: 10.1111/j.1365-2486.2011.02590.x
- Pilkington, M. G., Caporn, S. J. M., Carroll, J. A., Cresswell, N., Lee, J. A., Reynolds, B., et al. (2005). Effects of increased deposition of atmospheric nitrogen on an upland moor: nitrogen budgets and nutrient accumulation. *Environ. Pollut.* 138, 474–485. doi: 10.1016/j.envpol.2005.04.011
- Pinheiro, J., Bates, D., DebRoy, S., and Sarkar, D. (2016). *nlme: Linear and Nonlinear Mixed Effects Models*. R package version 3.1–121.
- Power, S. A., Ashmore, M. R., and Cousins, D. A. (1998). Impacts and fate of experimentally enhanced nitrogen deposition on a British lowland heath. *Environ. Pollut.* 102, 27–34. doi: 10.1016/S0269-7491(98)80011-0
- Proctor, M. C. F. (2008). “Physiological ecology,” in *Bryophyte Ecology*, 2nd Edn., eds B. Goffinet and A. J. Shaw (Cambridge: Cambridge University Press), 237–268.
- Rose, R. J., Webb, N. R., Clarke, R. T., and Traynor, C. H. (2000). Changes on the heathlands in Dorset, England, between 1987 and 1996. *Biol. Conserv.* 93, 117–125. doi: 10.1016/S0006-3207(99)00047-6
- Sala, O. E., Chapin, F. S., Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., et al. (2000). Global biodiversity scenarios for the year 2100. *Science* 287, 1770–1774. doi: 10.1126/science.287.5459.1770
- Schimel, J. P., and Bennett, J. (2004). Nitrogen mineralization: challenges of a changing paradigm. *Ecology* 85, 591–602. doi: 10.1890/03-8002
- Schmidt, I. K., Tietema, A., Williams, D., Gundersen, P., Beier, C., Emmett, B. A., et al. (2004). Soil solution chemistry and element fluxes in three European heathlands and their responses to warming and drought. *Ecosystems* 7, 638–649. doi: 10.1007/s10021-004-0217-5
- Sebilio, M., Mayer, B., Grably, M., Billion, D., and Mariotti, A. (2004). The use of the “ammonium diffusion” method for $\delta^{15}\text{N-NH}_4^+$ and $\delta^{15}\text{N-NO}_3^-$ measurements: comparison with other techniques. *Environ. Chem.* 1, 99–103. doi: 10.1071/EN04037
- Silvertown, J., Tallwin, J., Stevens, C., Power, S. A., Morgan, V., Emmett, B., et al. (2010). Environmental myopia: a diagnosis and a remedy. *Trends Ecol. Evol.* 25, 556–561. doi: 10.1016/j.tree.2010.06.015
- Southon, G. E., Green, E. R., Jones, A. G., Barker, C. G., and Power, S. A. (2012). Long-term nitrogen additions increase likelihood of climate stress and affect recovery from wildfire in a lowland heath. *Glob. Chang. Biol.* 18, 2824–2837. doi: 10.1111/j.1365-2486.2012.02732.x
- Startsev, N. A., and Lieffers, V. J. (2007). Emission of nitrogen gas, nitrous oxide, and carbon dioxide on rehydration of dry feathermosses. *Soil Sci. Soc. Am. J.* 71, 214–218. doi: 10.2136/sssaj2004.0376
- Startsev, N. A., Lieffers, V. J., Landhäusser, S. M., and Velazquez-Martinez, A. (2008). N-transfer through aspen litter and feather moss layers after fertilization with ammonium nitrate and urea. *Plant Soil* 311, 51–59. doi: 10.1007/s11104-008-9657-6
- Templer, P. H., Mack, M. C., Chapin, F. S., Christenson, L. M., Compton, J. E., Crook, H. D., et al. (2012). Sinks for nitrogen inputs in terrestrial ecosystems: a meta-analysis of ^{15}N tracer field studies. *Ecology* 93, 1816–1829. doi: 10.1890/11-1146.1
- Tye, A. M., Young, S. D., Crout, N. M. J., West, H. M., Stapleton, L. M., Poulton, P. R., et al. (2005). The fate of ^{15}N added to high Arctic tundra to mimic increased inputs of atmospheric nitrogen released from a melting snowpack. *Glob. Chang. Biol.* 11, 1640–1654. doi: 10.1111/j.1365-2486.2005.01044.x
- Vandvik, V., Heegaard, E., Maren, I. E., and Aarrestad, P. A. (2005). Managing heterogeneity: the importance of grazing and environmental variation on post-fire succession in heathlands. *J. Appl. Ecol.* 42, 139–149. doi: 10.1111/j.1365-2664.2005.00982.x
- von Oheimb, G., Power, S. A., Falk, K., Friedrich, U., Mohamed, A., Krug, A., et al. (2010). N:P ratio and the nature of nutrient limitation in *Calluna*-dominated heathlands. *Ecosystems* 13, 317–327. doi: 10.1007/s10021-010-9320-y
- Wood, S. N. (2006). *Generalized Additive Models: An Introduction with R*. London; New York, NY: Chapman & Hall/CRC; Boca Raton.
- Zuur, A. F., Fichtner, A., Ieno, E. N., and Saveliev, A. A. (2014). “Gamma GAMM applied on tree growth data,” in *A beginner's Guide to Generalized Additive Mixed Models with R*, eds A. F. Zuur, A. A. Saveliev, and E. N. Ieno (Newburgh, NY: Highland Statistics Ltd.), 217–239.
- Zuur, A. F., Ieno, E. N., and Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems. *Methods Ecol. Evol.* 1, 3–14. doi: 10.1111/j.2041-210X.2009.00001.x
- Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., and Smith, G. M. (2009). *Mixed Effects Models and Extensions in Ecology with R*. New York, NY: Springer Science+Business Media.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Bähring, Fichtner, Friedrich, von Oheimb and Hårdtle. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.