

COVID-19 stressors and health behaviors

Keng, Shian Ling; Stanton, Michael V.; Haskins, Lee Ann B.; Almenara, Carlos A.; Ickovics, Jeannette; Jones, Antwan; Grigsby-Toussaint, Diana; Agostini, Maximilian; Bélanger, Jocelyn J.; Gützkow, Ben; Kreienkamp, Jannis; Lemay, Edward P.; vanDellen, Michelle R.; Abakoumkin, Georgios; Abdul Khaiyom, Jamilah Hanum; Ahmedi, Vjollca; Akkas, Handan; Atta, Mohsin; Bagci, Sabahat Cigdem; Basel, Sima; Berisha Kida, Edona; Bernardo, Allan B.I.; Buttrick, Nicholas R.; Chobthamkit, Phatthanakit; Choi, Hoon Seok; Cristea, Mioara; Csaba, Sára; Damnjanovic, Kaja; Danyliuk, Ivan; Dash, Arobindu; Di Santo, Daniela; Douglas, Karen M.; Enea, Violeta; Faller, Daiane G.; Fitzsimons, Gavan; Gheorghiu, Alexandra; Gómez, Ángel; Hamaidia, Ali; Han, Qing; Helmy, Mai; Hudiyana, Joevarian; Jeronimus, Bertus F.; Jiang, Ding Yu; Jovanović, Veljko; Kamenov, Željka; Kende, Anna; Kieu, Tra Thi Thanh; Koc, Yasin; Kovyazina, Kamila; Kozytska, Inna; Krause, Joshua; Kruglanski, Arie W.; Kurapov, Anton; Kutlaca, Maja; Lantos, Nóra Anna; Lesmana, Cokorda Bagus Jaya; Louis, Winnifred R.; Lueders, Adrian; Maj, Marta; Malik, Najma Iqbal; Martinez, Anton; McCabe, Kira O.; Mehulić, Jasmina; Milla, Mirra Noor; Mohammed, Idris; Molinario, Erica; Moyano, Manuel; Muhammad, Hayat; Mula, Silvana; Muluk, Hamdi; Myroniuk, Solomiia; Najafi, Reza; Nisa, Claudia F.; Nyúl, Boglárka; O'Keefe, Paul A.; Osuna, Jose Javier Olivas; Osin, Evgeny N.; Park, Joonha; Pica, Gennaro; Pierro, Antonio; Rees, Jonas; Reitsema, Anne Margit; Resta, Elena; Rullo, Marika; Ryan, Michelle K.; Samekin, Adil; Santtila, Pekka; Sasin, Edyta M.; Schumpe, Birga M.; Selim, Heyla A.; Stroebe, Wolfgang; Sultana, Samiah; Sutton, Robbie M.; Tseliou, Eleftheria; Utsugi, Akira; van Breen, Jolien Anne; Van Lissa, Caspar J.; Van Veen, Kees; Vázquez, Alexandra; Wollast, Robin; Yeung, Victoria Wai Ian; Zand, Somayeh; Žeželj, Iris Lav; Zheng, Bang; Zick, Andreas; Zúñiga, Claudia; Leander, N. Pontus

Published in:
Preventive Medicine Reports

DOI:
[10.1016/j.pmedr.2022.101764](https://doi.org/10.1016/j.pmedr.2022.101764)

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for pulished version (APA):

Keng, S. L., Stanton, M. V., Haskins, L. A. B., Almenara, C. A., Ickovics, J., Jones, A., Grigsby-Toussaint, D., Agostini, M., Bélanger, J. J., Gützkow, B., Kreienkamp, J., Lemay, E. P., vanDellen, M. R., Abakoumkin, G., Abdul Khaiyom, J. H., Ahmed, V., Akkas, H., Atta, M., Bagci, S. C., ... Leander, N. P. (2022). COVID-19 stressors and health behaviors: A multilevel longitudinal study across 86 countries. *Preventive Medicine Reports*, 27, Article 101764. <https://doi.org/10.1016/j.pmedr.2022.101764>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 04. Dez.. 2025



COVID-19 stressors and health behaviors: A multilevel longitudinal study across 86 countries

Shian-Ling Keng^{a,b,q,*}, Michael V. Stanton^{b,1}, LeeAnn B. Haskins^c, Carlos A. Almenara^d, Jeannette Ickovics^{e,bq}, Antwan Jones^f, Diana Grigsby-Toussaint^g, Maximilian Agostini^h, Jocelyn J. Bélangerⁱ, Ben Gützkow^h, Jannis Kreienkamp^h, Edward P. Lemay Jr.^{am}, Michelle R. vanDellen^c, Georgios Abakoumkin^j, Jamilah Hanum Abdul Khaiyom^k, Vjollca Ahmedi^l, Handan Akkas^m, Mohsin Attaⁿ, Sabahat Cigdem Bagci^o, Sima Baselⁱ, Edona Berisha Kida^l, Allan B.I. Bernardo^p, Nicholas R. Buttrick^q, Phatthanakit Chobthamkit^r, Hoon-Seok Choi^s, Mioara Cristea^t, Sára Csaba^u, Kaja Damjanovic^v, Ivan Danyliuk^w, Arobindu Dash^x, Daniela Di Santo^y, Karen M. Douglas^z, Violeta Enea^{aa}, Daiane G. Faller^{bn}, Gavan Fitzsimons^{ab}, Alexandra Gheorghiu^{aa}, Ángel Gómez^{ac}, Ali Hamaidia^{ad}, Qing Han^{ae}, Mai Helmy^{af,bo}, Joevarian Hudiyana^{ag}, Bertus F. Jeronimus^h, Ding-Yu Jiang^{ah}, Veljko Jovanović^{ai}, Željka Kamenov^{aj}, Anna Kende^u, Tra Thi Thanh Kieu^{ak}, Yasin Koc^h, Kamila Kovyazina^{al}, Inna Kozytska^w, Joshua Krause^h, Arie W. Kruglanski^{am}, Anton Kurapov^w, Maja Kutlaca^{an}, Nóra Anna Lantos^u, Cokorda Bagus Jaya Lesmana^{ao}, Winnifred R. Louis^{ap}, Adrian Lueders^{aq}, Marta Maj^{ar}, Najma Iqbal Malikⁿ, Anton Martinez^{as}, Kira O. McCabe^{at}, Jasmina Mehulić^{aj}, Mirra Noor Milla^{ag}, Idris Mohammed^{au}, Erica Molinario^{bp}, Manuel Moyano^{av}, Hayat Muhammad^{aw}, Silvana Mula^y, Hamdi Muluk^{ag}, Solomiia Myroniuk^h, Reza Najafi^{ax}, Claudia F. Nisaⁱ, Boglárka Nyúl^u, Paul A. O'Keefe^{bq}, Jose Javier Olivas Osuna^{br}, Evgeny N. Osin^{ay}, Joonha Park^{az}, Gennaro Pica^{ba}, Antonio Pierro^y, Jonas Rees^{bb}, Anne Margit Reitsema^h, Elena Resta^y, Marika Rullo^{bc}, Michelle K. Ryan^{h,bd}, Adil Samekin^{be}, Pekka Santtila^{bf}, Edyta M. Sasinⁱ, Birga M. Schumpe^{bu}, Heyla A. Selim^{bg}, Wolfgang Stroebe^h, Samiah Sultana^h, Robbie M. Sutton^z, Eleftheria Tseliou^j, Akira Utsugi^{bh}, Jolien Anne van Breen^{bi}, Caspar J. Van Lissa^{bj}, Kees Van Veen^h, Alexandra Vázquez^{ac}, Robin Wollast^{bs}, Victoria Wai-lan Yeung^{bk}, Somayah Zand^{bt}, Iris Lav Žeželj^v, Bang Zheng^{bl}, Andreas Zick^{bb}, Claudia Zúñiga^{bm}, N. Pontus Leander^{h,bv}

^a Monash University Malaysia, Malaysia

^b California State University, East Bay, USA

^c University of Georgia, USA

^d Universidad Peruana de Ciencias Aplicadas, Peru

^e Yale University, USA

^f The George Washington University, USA

^g Brown University, USA

^h University of Groningen, The Netherlands

ⁱ New York University Abu Dhabi, United Arab Emirates

^j University of Thessaly, Volos, Greece

^k International Islamic University Malaysia, Gombak, Malaysia

^l University of Pristina, Pristina, Kosovo

^m Ankara Science University, Ankara, Turkey

ⁿ University of Sargodha, Sargodha, Pakistan

* Corresponding author at: Monash University Malaysia, Jalan Lagoon Selatan, Bandar Sunway, 47500 Subang Jaya, Selangor, Malaysia.

E-mail address: keng.sl@monash.edu (S.-L. Keng).

<https://doi.org/10.1016/j.pmedr.2022.101764>

Received 7 July 2021; Received in revised form 6 March 2022; Accepted 13 March 2022

Available online 17 March 2022

2211-3355/© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

- ^o Sabanci University, Istanbul, Turkey
^p De La Salle University, Manila, Philippines
^q University of Virginia, Charlottesville, USA
^r Thammasat University, Bangkok, Thailand
^s Sungkyunkwan University, Seoul, South Korea
^t Heriot Watt University, United Kingdom
^u Eötvös Loránd University (ELTE), Budapest, Hungary
^v University of Belgrade, Belgrade, Serbia
^w Taras Shevchenko National University of Kyiv, Kiev, Ukraine
^x Leuphana University Lüneburg, Lüneburg, Germany
^y Sapienza University of Rome, Rome, Italy
^z University of Kent, Canterbury, UK
^{aa} Alexandru Ioan Cuza University of Iasi, Iasi, Romania
^{ab} Duke University, Durham, USA
^{ac} Universidad Nacional de Educación a Distancia (UNED), Madrid, Spain
^{ad} University Setif 2, Sétif, Algeria
^{ae} University of Bristol, Bristol, UK
^{af} Menoufia University, Al Minufiyah, Egypt
^{ag} Universitas Indonesia, Depok, Indonesia
^{ah} National Chung-Cheng University, Chiayi, Taiwan
^{ai} University of Novi Sad, Novi Sad, Serbia
^{aj} University of Zagreb, Zagreb, Croatia
^{ak} HCMC University of Education, Ho Chi Minh City, Viet Nam
^{al} Independent Researcher, Nur-Sultan, Kazakhstan
^{am} University of Maryland, College Park, USA
^{an} Durham University, Durham, UK
^{ao} Udayana University, Denpasar, Indonesia
^{ap} University of Queensland, Brisbane, Australia
^{aq} University of Limerick, Ireland
^{ar} Jagiellonian University, Kraków, Poland
^{as} University of Sheffield, Sheffield, UK
^{at} Carleton University, Canada
^{au} Usmanu Danfodiyo University Sokoto, Sokoto, Nigeria
^{av} University of Cordoba, Córdoba, Spain
^{aw} University of Peshawar, Peshawar, Pakistan
^{ax} University of Padova, Italy
^{ay} National Research University Higher School of Economics, Moscow, Russia
^{az} NUCB Business School, Nagoya, Japan
^{ba} University of Camerino, Camerino, Italy
^{bb} University of Bielefeld, Bielefeld, Germany
^{bc} University of Siena, Siena, Italy
^{bd} University of Exeter, Exeter, UK
^{be} School of Liberal Arts, M. Narikbayev KAZGUU University, Nur-Sultan, Kazakhstan
^{bf} New York University Shanghai, Shanghai, China
^{bg} King Saud University, Riyadh, Saudi Arabia
^{bh} Nagoya University, Nagoya, Japan
^{bi} Leiden University, Leiden, The Netherlands
^{bj} Utrecht University, Utrecht, The Netherlands
^{bk} Lingnan University, Tuen Mun, Hong Kong
^{bl} Imperial College London, London, UK
^{bm} Universidad de Chile, Santiago, Chile
^{bn} National University of Singapore, Singapore, Singapore
^{bo} Sultan Qaboos University, Egypt
^{bp} Florida Gulf Coast University, USA
^{bq} Yale-NUS College, Singapore, Singapore
^{br} National Distance Education University, Spain
^{bs} Université Clermont-Auvergne, France
^{bt} University of Milano-Bicocca, Italy
^{bu} University of Amsterdam, The Netherlands
^{bv} Wayne State University, USA

ARTICLE INFO

Keywords:
 COVID-19
 Health behaviors
 Infection risk
 Economic burden

ABSTRACT

Anxiety associated with the COVID-19 pandemic and home confinement has been associated with adverse health behaviors, such as unhealthy eating, smoking, and drinking. However, most studies have been limited by regional sampling, which precludes the examination of behavioral consequences associated with the pandemic at a global level. Further, few studies operationalized pandemic-related stressors to enable the investigation of the impact of different types of stressors on health outcomes. This study examined the association between perceived risk of COVID-19 infection and economic burden of COVID-19 with health-promoting and health-damaging behaviors using data from the PsyCorona Study: an international, longitudinal online study of psychological and behavioral correlates of COVID-19. Analyses utilized data from 7,402 participants from 86 countries across three waves of assessment between May 16 and June 13, 2020. Participants completed self-report measures of COVID-19 infection risk, COVID-19-related economic burden, physical exercise, diet quality, cigarette smoking,

¹ Shian-Ling Keng and Michael Stanton are co-first authors on this paper.

sleep quality, and binge drinking. Multilevel structural equation modeling analyses showed that across three time points, perceived economic burden was associated with reduced diet quality and sleep quality, as well as increased smoking. Diet quality and sleep quality were lowest among respondents who perceived high COVID-19 infection risk combined with high economic burden. Neither binge drinking nor exercise were associated with perceived COVID-19 infection risk, economic burden, or their interaction. Findings point to the value of developing interventions to address COVID-related stressors, which have an impact on health behaviors that, in turn, may influence vulnerability to COVID-19 and other health outcomes.

1. Introduction

The COVID-19 pandemic has caused profound adverse health, economic, and psychological consequences. To contain the spread of the pandemic, many countries have imposed lockdowns, limiting citizens' participation in regular social and physical activities. Though essential to slow the rate of infection, lockdowns have been found to be positively associated with negative mental health consequences, such as depression and anxiety (Huang and Zhao, 2020; Nguyen et al., 2020). Furthermore, such measures are likely to impact health-related behaviors: restricted mobility decreases physical activity, and heightened psychological distress increases the propensity to engage in unhealthy eating, smoking, and binge drinking (Grzywacz and Almeida, 2008; Kassel et al., 2003). These unhealthy behaviors are risk factors for non-communicable diseases, including obesity, diabetes, and cardiovascular diseases (Thornton et al., 2016; Stang et al., 2000; Hu et al., 2000), which in turn increase the risk of contracting COVID-19 and greater disease severity and may eventually lead to increased mortality (Esai, 2020; Zheng et al., 2020).

To date, results are mixed across extant cross-sectional studies looking at the relationship between stress related to COVID-19 and unhealthy behaviors. In the United States, pandemic-related psychological distress was positively associated with alcohol use, with women being significantly more likely to consume greater amounts of alcohol on a typical evening and during their recent heaviest drinking occasion (Rodríguez et al., 2020). In Vietnam, fear of COVID-19 was associated with greater alcohol consumption and smoking among college students (Nguyen et al., 2020). In contrast, a study based in Spain reported less alcohol consumption and better dietary behaviors during the COVID-19 lockdown (Rodríguez-Pérez et al., 2020). In China, pandemic-related home isolation was associated with improvements in dietary behaviors and sleep quality, even though time spent being sedentary increased during lockdown compared to pre-lockdown (Wang et al., 2020). These varying associations could in part be attributed to regional variations in lockdown policies, which affect ease of access to health-relevant resources such as exercise facilities, and outdoor dining options.

Even though these studies provide some insight into the potential impact of the pandemic on health behaviors, several caveats can be identified. First, the majority of the studies are regionally focused and do not explore global trends. One exception is a study involving over 1000 adults in Asia, Europe, and Africa, which documented a decrease in physical activity and binge drinking and an increase in unhealthy food consumption during COVID-19 home confinement (Ammar et al., 2020). The analyses however did not control for potential confounding variables, such as gender, age, and education that may have explained the changes in these health behaviors. Though most individuals likely experienced heightened anxiety about contracting COVID-19, the degree of anxiety and perceived risk may also vary globally depending on access to protective measures, as well as perceived effectiveness of the government and/or the community in curbing the pandemic.

Further, few studies have operationalized stressors related to the pandemic. Two critical stressors faced by many individuals during the pandemic include infection risk and economic burden. During the ongoing pandemic, many individuals experience varying degrees of financial impact, with millions facing unemployment and loss of income and housing, which may adversely impact health-related behaviors and

outcomes. It remains to be examined whether perceived risk of infection and economic burden may differentially impact health behaviors and whether these stressors may interact to predict engagement in specific health behaviors. Importantly, these effects should be assessed while controlling for sociodemographic characteristics, which are known to impact health behaviors, such as binge drinking, smoking, and healthy eating (Wilsnack et al., 2018; Wardle et al., 2004; Bauer et al., 2007; Cavelaars et al., 2000).

In this study, we utilized data from a multinational, longitudinal online study on psychological and behavioral correlates of COVID-19 to examine the association between perceived risk of infection and economic burden with several health-promoting (exercise, diet quality, sleep quality) and health-damaging (binge drinking, smoking) behaviors. We hypothesized that perceived risk of infection and economic burden would be associated with reduced engagement in healthier behaviors. Specifically, we predicted that higher levels of perceived infection risk and economic burden would each independently be associated with less exercise, poorer diet, and worse sleep quality, as well as more binge drinking and smoking, independent of the effects of demographic factors. Additionally, we expected the interaction between perceived infection risk and economic burden would be a particularly strong predictor of health-damaging behaviors. Recruitment of a large international sample enabled us to observe the association between pandemic-related stressors and health behaviors on a global scale.

2. Method

2.1. Participants and procedure

The sample consisted of adult participants (aged 18 and above) of an online, longitudinal study as part of the PsyCorona project (<https://psycorona.org/>), a multinational research project examining behavioral and psychological responses to the COVID-19 pandemic. Research participants initially completed a baseline cross-sectional survey, and a subset of participants signed up for a longitudinal study involving follow-up surveys over the course of the pandemic (Jin et al., 2021; Han et al., 2021; Romano et al., 2020). Our analysis focused on a self-selected cohort of participants ($N = 7,402$) who completed Wave 7, 9, and 11 of assessments (administered in two-week intervals) between May 16 and June 13 of 2020. Each assessment lasted approximately 10 min. The surveys were translated into 30 languages and distributed by members of the research team (consisting of over 100 behavioral scientists) in their respective countries using social media campaigns, press releases, and social and academic networks.

This study complies with ethical regulations for research on human subjects. All participants gave informed consent, as approved by the Institutional Review Board at New York University Abu Dhabi (HRPP-2020-42) and the Ethics Committee of Psychology at Groningen University (PSY-1920-S-0390).

2.2. Measures

2.2.1. Perceived Stressors: COVID-19 infection risk and economic burden

Perceived stress was measured by the item: "How likely is it that the following will happen to you in the next few months?" (1) COVID-19 infection risk – "you will get infected with coronavirus", and (2)

Economic burden – “your personal situation will get worse due to economic consequences of coronavirus.” Responses were based on a Likert-type scale of 1 (*very unlikely*) to 8 (*already happened*).

2.2.2. Health behaviors

Five health-related behaviors were measured with single-item questions:

- (1) *Physical Exercise* was measured with the question: “During the past week, how many days did you do 20 min of vigorous (sweating and puffing) or 30 min of moderate (increasing your heart rate but not vigorous) physical activity?” (adapted from the Brief Physical Activity Assessment Tool) (Marshall et al., 2005). Participants responded using a range of 0 to 7 days.
- (2) *Diet quality* was assessed with the question: “During the past week, how healthy was your overall diet? Consider how many sweets you have been eating as well as how many portions of fruit and/or vegetables you ate each day” (adapted from National Health and Nutrition Examination Survey Questionnaire) (National Health and Nutrition Examination Survey Questionnaire, 2018). Participants were asked to provide a rating on a 1 (*poor*) to 5 (*excellent*) scale.
- (3) *Sleep quality* was measured with the question: “During the past week, how would you rate your sleep quality overall?” (adapted from Pittsburgh Sleep Quality Index) (Buysse et al., 1989). Participants were asked to provide a rating on a 1 (*poor*) to 5 (*excellent*) scale.
- (4) *Binge drinking* was measured with the item: “During the past week, how many days did you have >4 drinks in a day?” (adapted from a screening test for unhealthy alcohol use recommended by the National Institute on Alcohol Abuse and Alcoholism) (Smith et al., 2009). Participants responded using a range of 0 to 7 days.
- (5) *Smoking* was assessed with the item: “During the past week, how many cigarettes did you smoke each day?”, with an open response option (adapted from National Health and Nutrition Examination Survey Questionnaire) (National Health and Nutrition Examination Survey Questionnaire, 2018). This variable was transformed into four categories: 0 cigarettes per day coded as non-smoker, 1–10 cigarettes per day coded as light smoker, 11–19 cigarettes per day coded as moderate smoker, >=20 cigarettes per day coded as heavy smoker, following the criteria of the Government of Canada (Government of Canada, 2008). After a visual inspection of the dataset, plots, and measures of dispersion, we excluded outliers, particularly those who reported smoking >75 cigarettes per day ($n = 37$, $n = 24$, and $n = 28$, in waves 7, 9, and 11, respectively).

2.2.3. Sociodemographic characteristics

Participants provided information about age, categorized on a scale from 1 (18–24 years old) to 7 (75 + years old); education, categorized on a scale from 1 (*elementary*) to 6 (*doctorate*); and gender, categorized as 1 (*female*), 2 (*male*), and 3 (*other*). For the purpose of our analyses, gender was re-coded into a binary variable (0 = female, 1 = male, whereas “other” was excluded from analyses).

2.3. Statistical analyses

Demographic information was assessed using SAS. Mplus 8.4 was used to conduct multilevel structural equation modeling (MSEM) bivariate correlations and regression. Data from Waves 7, 9, and 11 (time points; level 1) were nested within the participants (level 2). All health behavior outcomes had sufficient variance across the two levels ($ICCs > 0.68$), so MSEM was employed to estimate the structural relationships at both levels (i.e., within and between persons). Acknowledging that participants were nested within geographical region (i.e., North America, Europe, Asia, Africa, Oceania, Caribbean, Central, and

Table 1

Sample Characteristics (N = 7402).

Variable	n (percentage)
Gender	
Female	4959 (67%)
Male	2443 (33%)
Age	
18 to 24 years old	794 (10.73%)
25 to 34 years old	1235 (16.68%)
35 to 44 years old	1260 (17.02%)
45 to 54 years old	1386 (18.72%)
55 to 64 years old	1400 (18.91%)
65 to 74 years old	1143 (15.44%)
75 and older	184 (2.48%)
Region	
Europe	4510 (61.01%)
North America	1387 (18.74%)
Asia	633 (8.56%)
Caribbean, Central and South America	486 (6.57%)
Oceania	197 (2.67%)
Africa	179 (2.42%)
Country Not Indicated	10 (0.14%)
Education	
Elementary and Secondary Education	907 (12.25%)
Vocational Education	831 (11.23%)
Higher Education (Without a Bachelor's Degree)	1504 (20.32%)
Bachelor's Degree	2018 (27.26%)
Master's degree	1590 (21.48%)
Doctorate Degree	552 (7.46%)

South America) (United Nations. World Population Prospectus, 2019), we evaluated the intraclass correlations (ICCs) of each of the health behaviors by adding region as a level 3 variable (time points within participants within region). We evaluated region as opposed to country as a level 3 variable because of limited samples from some countries (e.g., $n < 10$), which precluded sufficient data for analyses of country as a higher order variable. However, because all ICCs were at or below 0.05, we did not include region as a level 3 variable in the final MSEM analyses (LeBreton and Senter, 2008).

Because the current research interest was to evaluate the effects of COVID-19 stressors on health behaviors across individuals, all results reported are at the between-person level and over three time periods. As part of preliminary analyses, we conducted MSEM bivariate correlational analyses to examine the association between demographic factors and COVID-19 related stressors, as well as each of the health behaviors. Next, we conducted MSEM regression with random intercepts and fixed slopes to examine the role of perceived infection risk, economic burden, and their interaction as predictors of each of the health behaviors. All MSEM regression analyses included age, gender, and education as between-person covariates. Analyses were conducted using full-information maximum likelihood estimation, which provides standard errors that are robust to data non-normality and non-independence (Heck and Thomas, 2015).

3. Results

3.1. Sample characteristics and preliminary analyses

The sample consisted of 7,402 participants from 86 countries. Table 1 provides a detailed breakdown of demographic information in this sample. Sixty-seven percent ($n = 4959$) of the participants were female. Regionally, more than one-half of the sample was based in Europe (60.9%), followed by North America (14.8%) and Asia (6.7%). There was a relatively even distribution of individuals across age groups: 63.1% were between 18 and 54 years. More than half (56.2%) had at

Table 2
Descriptive Statistics for COVID-19 Stressors and Health Behaviors.

Variable	N	Scale	Mean	SD
Perceived Infection Risk	7402	1 (<i>very unlikely</i>) – 8 (<i>already happened</i>)	3.56	1.33
Perceived Economic Burden	7402	1 (<i>very unlikely</i>) – 8 (<i>already happened</i>)	3.93	1.76
Exercise	7401	Days in the past week	2.54	2.19
Diet Quality	7401	1 (<i>poor</i>) – 5 (<i>excellent</i>)	3.00	0.96
Sleep Quality	7400	1 (<i>poor</i>) – 5 (<i>excellent</i>)	2.73	1.04
Binge Drinking Variable	7401	Days in the past week (0–7)	0.65	1.49
Smoking	4664	0 = Non-smoker 1 = Light Smoker 2 = Moderate Smoker 3 = Heavy Smoker	Frequency (Percentage) 3654 (78.34%) 495 (10.61%) 213 (4.57%) 282 (6.05%)	

least a college degree. A list of all countries included in this study is provided in S1, under [Supplementary Materials](#). Table 2 presents the descriptive statistics of COVID-19 stressors and health behavior outcomes across the whole sample.

We next examined demographic factors (age, gender, and education) as potential correlates of the two COVID-19 stressors and each of the health behaviors (see [Table 3](#)). Older age predicted significantly lower perceived COVID infection risk and economic burden, better diet and sleep quality, and more cigarettes smoked in the past week, all $ps < 0.001$. Being male was associated with lower perceived infection risk, better perceived diet and sleep quality, and more smoking and binge drinking, all $ps < 0.01$. Higher education levels were associated with significantly greater perceived COVID infection risk, better diet quality, more days spent engaging in moderate to vigorous exercise, and fewer cigarettes smoked, all $ps < 0.01$.

3.2. Perceived infection Risk, economic Burden, and their interaction as predictors of each health behavior and outcome

Between-person results of the multilevel structural equation modeling analyses are presented in [Table 4](#). Post hoc power analyses were conducted to determine achieved power for each parameter coefficient in the five models. Power analysis was conducted using Monte Carlo simulation with 500 replications using Robust Maximum Likelihood (MLR) estimation in *Mplus*. The analyses indicated adequate power (>80%) to detect the majority of effects, with the exception of physical

exercise, binge drinking, and select parameter estimates for smoking. Within-person results are reported in S2, under [Supplementary Materials](#).

COVID-related infection risk and economic burden were both negatively associated with perceived diet quality during the previous week. These main effects were qualified by a significant interaction between perceived infection risk and perceived economic burden, $b = 0.01$, $SE = 0.01$, $p < .05$. As shown in [Fig. 1](#), those who reported high economic burden (top 10%) reported lower diet quality regardless of levels of perceived infection risk, $b = 0.008$, $SE = 0.02$, $p = .693$, whereas those perceiving low economic burden (bottom 10%) reported better diet quality if their perceived infection risk was also low, $b = -0.057$, $SE = 0.02$, $p = .002$.

COVID-related infection risk and perceived economic burden were both negatively associated with sleep quality during the previous week. These main effects were qualified by a significant interaction, $b = 0.67$, $SE = 0.01$, $p < .001$. As shown in [Fig. 2](#), those who reported high economic burden (top 10%) reported decreased sleep quality regardless of levels of perceived infection risk, $b = -0.02$, $SE = 0.02$, $p = .325$, whereas people perceiving low economic burden (bottom 10%) reported better sleep quality if their perceived infection risk was also low, $b = -0.111$, $SE = 0.02$, $p < .001$.

Perceived economic burden was positively associated with the number of cigarettes smoked. COVID-related infection risk was not associated with the number of cigarettes smoked in the previous week. There was no significant interaction between infection risk and economic burden in predicting the number of cigarettes smoked.

No relationship was observed between perceived COVID-related infection risk, economic burden or their interaction and the number of days spent binge drinking or the number of days spent exercising moderately or vigorously. Across these analyses, none of the associations at the within-person level were significant, indicating stability in participants' responses over time.

4. Discussion

This longitudinal study of health behaviors during the COVID-19 pandemic found that two pandemic-related stressors – perceived infection risk and perceived economic burden – were associated with a range of health-related behaviors and outcomes. In particular, perceived economic burden related to the pandemic was found to have the most consistent negative impact across several health behavior outcomes, including diet quality, sleep quality, and cigarette smoking. Economic burden may lead to individuals engaging in unhealthy behaviors as a

Table 3
Bivariate Relationships among Demographic Variables, COVID-19 Stressors, and Health Behaviors.

	Age	Gender	Education	Perceived Infection Risk	Perceived Economic Burden	Physical Exercise	Diet Quality	Sleep Quality	Binge Drinking	Smoking
Age	–									
Gender	0.18***	–								
Education	-0.28***	-0.04***	–							
Perceived Infection Risk	-0.27***	-0.04***	0.18***	–						
Perceived Economic Burden	-0.31***	-0.02	-0.04	0.67***	–					
Exercise	0.06	0.03	0.41***	-0.02	-0.25***	–				
Diet Quality	0.20***	0.02**	0.11***	-0.10***	-0.27***	0.65***	–			
Sleep Quality	0.15***	0.04***	0.04	-0.19***	-0.39***	0.35***	0.39***	–		
Binge Drinking	0.05	0.09***	-0.03	-0.02	0.07	0.05	-0.02	0.00	–	
Smoking	0.09***	0.02***	-0.15***	-0.04**	0.13***	-0.15***	-0.04***	-0.01	0.14***	–

Notes. Gender is coded as 0 (female) and 1 (male); Education is coded on a scale from 1 (elementary) to 6 (doctorate); ** $p < .01$; *** $p < .001$.

Table 4

Test Statistics for Multilevel Regression with Each Health Behavior Predicted by Infection Risk, Economic Burden, and Their Interaction.

Physical Exercise						
	<i>b</i>	SE	<i>p</i>	95% CI (Lower)	95% CI (Upper)	Achieved Power to Detect Parameter Estimate
Infection Risk	0.06	0.06	0.34	−0.06	0.18	0.25
Economic Burden	−0.06	0.05	0.25	−0.16	0.04	0.35
Infection Risk* Economic Burden	−0.01	0.01	0.43	−0.04	0.02	0.20
Diet Quality						
	<i>b</i>	SE	<i>p</i>	95% CI (Lower)	95% CI (Upper)	Achieved Power to Detect Parameter Estimate
Infection Risk	−0.08	0.03	0.004	−0.13	−0.03	0.98
Economic Burden	−0.14	0.02	< 0.001	−0.19	−0.09	>0.99
Infection Risk* Economic Burden	0.01	0.01	0.028	0.00	0.03	0.90
Sleep Quality						
	<i>b</i>	SE	<i>p</i>	95% CI (Lower)	95% CI (Upper)	Achieved Power to Detect Parameter Estimate
Infection Risk	−0.15	0.03	< 0.001	−0.20	−0.09	>0.99
Economic Burden	−0.20	0.03	< 0.001	−0.25	−0.15	>0.99
Infection Risk* Economic Burden	0.02	0.01	0.002	0.01	0.03	0.99
Binge Drinking						
	<i>b</i>	SE	<i>p</i>	95% CI (Lower)	95% CI (Upper)	Achieved Power to Detect Parameter Estimate
Infection Risk	−0.06	0.04	0.14	−0.14	0.02	0.50
Economic Burden	−0.00	0.04	0.93	−0.07	0.07	0.06
Infection Risk* Economic Burden	0.01	0.01	0.27	−0.01	0.03	0.33
Smoking						
	<i>b</i>	SE	<i>p</i>	95% CI (Lower)	95% CI (Upper)	Achieved Power to Detect Parameter Estimate
Infection Risk	−0.04	0.02	0.075	−0.08	0.00	0.55
Economic Burden	0.06	0.02	0.002	0.02	0.10	0.97
	0.00	0.01	0.96	−0.01	0.01	0.05

Table 4 (continued)

Physical Exercise						
	<i>b</i>	SE	<i>p</i>	95% CI (Lower)	95% CI (Upper)	Achieved Power to Detect Parameter Estimate
Infection Risk*						
Economic Burden						

Note. The above analyses included age, gender, and education as covariates.




Fig. 1. Interaction between Infection Risk and Economic Burden in Predicting Diet Quality. Note: Low economic burden is represented as the 10th percentile, equal to 1.67 on the economic burden scale of 1 to 8; High economic burden is represented as the 90th percentile, equal to 6.33 on the economic burden scale of 1 to 8. Thin dotted lines represent 95% confidence intervals.




Fig. 2. Interaction between Infection Risk and Economic Burden in Predicting Sleep Quality. Note: Low economic burden is represented as the 10th percentile, equal to 1.67 on the economic burden scale of 1 to 8; High economic burden is represented as the 90th percentile, equal to 6.33 on the economic burden scale of 1 to 8. Thin dotted lines represent 95% confidence intervals.

coping mechanism, consistent with theoretical and empirical work demonstrating an association between stress and health-damaging behaviors (Park and Iacocca, 2014). A recent report suggests that cash-based assistance in the form of stimulus check in the United States was linked to a robust 20% reduction in symptoms of depression and

anxiety during the pandemic (Fottrell, 0000). Therefore, economic burden might be related to unhealthy behaviors through symptoms of depression or anxiety, and when economic burden is alleviated, this may reduce unhealthy behaviors as well.

The finding that economic burden was associated with greater cigarette use is in line with previous research demonstrating a positive association between financial stress and tobacco use across households of varying incomes (Siahpush et al., 2003). Notably, the association between perceived economic burden and negative health outcomes may be bi-directional: heightened economic stress may increase smoking behaviors, and greater expenditure on acquiring tobacco products may pose further economic strain.

Consistent with past research, the present study documented a negative association between COVID-19 economic burden and sleep quality (Hall et al., 2009; Onder et al., 2020). This association may be accounted for by an increased tendency to engage in financial rumination and worry (de Bruijn and Antonides, 2020) which have been found to predict worsened sleep quality and mental health outcomes (Thorsteinsson et al., 2019). Financial stress may also be linked to unemployment, which affords greater unstructured time and likely more time for smoking and drinking, and fewer resources available for healthy food consumption (French and McKillop, 2017). In the context of the COVID-19 pandemic, stress and isolation resulting from government-imposed lockdowns and home quarantine may leave individuals more prone to engaging in unhealthy coping behaviors.

Importantly, the study found that perceived economic burden interacted with COVID-19 infection risk to predict worsened diet and sleep quality. This suggests that the main effects of perceived COVID-19-related stressors can only be meaningfully examined in the context of an interaction between the stressors. This finding highlights the need to develop interventions that address these stressors simultaneously to mitigate the negative impact of the COVID-19 pandemic on health outcomes. Specifically, economically disadvantaged populations are likely to be disproportionately impacted by the pandemic. There is therefore an urgent need to develop measures to lower their infection risk and economic burden, in order to mitigate the pandemic's long-term negative health consequences.

Contrary to our hypotheses, the study found no significant association between perceived infection risk and binge drinking, and only a trending, positive association between infection risk and smoking. It is plausible that attempts to drink or smoke may be driven more by general distress associated with the pandemic, as suggested by a study by Rodriguez and colleagues (Rodriguez et al., 2020), as opposed to the perception of infection risk, per se. The finding does not rule out the possibility that perceived infection risk is linked with more drinking that does not reach the threshold of a binge. The absence of a significant association between perceived infection risk and these behaviors may also reflect individual variations in response to infection risk: while some may be motivated to reduce engagement in health-damaging behaviors following awareness of high infection risk, others may engage in more of such behaviors as a coping mechanism (Park and Iacocca, 2014). Likewise, the lack of an association between the stressors and physical exercise may be attributable to significant individual variations in exercise habits during the pandemic, along with varying access to exercise facilities due to lockdowns.

The present study also identified a few demographic correlates of COVID-19 stressors and associated health behaviors. In particular, older individuals reported lower levels of perceived infection risk and economic burden, as well as better sleep and diet quality. The perception of lower infection risk could be due to several factors, such as the fact that older adults are less socially mobile. Compared to younger adults, they are also more likely to engage in prosocial COVID-19 protective behaviors like social distancing and mask-wearing (Jin et al., 2021). The finding that older individuals have better sleep quality suggests they may be less psychologically impacted by the pandemic, consistent with other studies' findings that older adults experience lower levels of

psychological symptoms and stress reactivity compared to younger adults, likely due to a higher degree of resilience (Nwachukwu et al., 2020; Nelson et al., 2021, 2021). Relative to females, males tend to perceive lower infection risk, in line with other research finding similar gender differences in the perception of seriousness of the COVID-19 pandemic (Galasso et al., 2020). Compared to females, males also smoke a greater number of cigarettes and spend more days binge drinking. Lastly, higher levels of education are identified consistently as a correlate of greater engagement in health-promoting behaviors and lower engagement in health-damaging behaviors. These findings point to the value of tailoring public campaigns to certain demographics such as young males, in order to reduce infection risk and likelihood of engaging in health-damaging behaviors.

This study is characterized by several strengths, such as recruitment of a large, multinational sample, a longitudinal design, and use of a multilevel analytical approach that takes into consideration potential variances accounted for by region and within-person variances across time. Limitations of the study include lack of representativeness and use of self-report measures, subject to recall and social desirability biases. Although several of the outcome measures were single-item, several of them were derived from established and validated scales. Due to limitations in survey length, some measures such as income and general mental health were not available. We did not examine patterns of behavior change over time because each of the 86 participating countries were in a different stage of dealing with the pandemic at the time of the surveys.

Future research could examine health behaviors using multimodal and/or objective measures (e.g., food diaries to assess diet, polysomnography to assess sleep quality). Future work should control for the effects of generalized anxiety or mental health symptoms to examine the unique effects of perceived infection risk and economic burden on health behaviors. Beyond infection risk and economic burden, social isolation is an additional stressor that should be examined as a potential contributor to health outcomes. Future research could also examine coping styles that may moderate the effects of pandemic-related stressors on health behaviors. Efforts should be made to examine specific communities (e.g., lower income groups) who may be at higher risk for contracting COVID-19 due to jobs that may not support social distancing. It would be of value to examine mechanisms underlying the associations between COVID-19 related stressors and health behaviors, including decisions about vaccinations, which were not yet available at the time of the surveys.

The COVID-19 pandemic persists, with >410 million confirmed cases and 5.8 million deaths globally as of February 14, 2022 (World Health Organization, 2021). Vaccination roll-out is moving quickly in a few countries, with marked delays in many more. Moreover, coronavirus variants are of grave concern. As such, it is critical that each country develops effective interventions tailored to the context of the local community, particularly to those who are economically disadvantaged and/or at higher infection risk, to mitigate the negative impact of the pandemic on health behaviors (Han et al., 2021; Nisa et al., 2021).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors would like to acknowledge Maleyka Mammadova for her assistance with literature review and data coding. This research received support from the New York University Abu Dhabi (VCDSF/75-71015), the University of Groningen (Sustainable Society & Ubbo Emmius Fund), and the Instituto de Salud Carlos III (COV20/00086). The COVID-19 risk perception item measured at baseline was previously

reported in unrelated test of effects on subjective well-being and mental health.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.pmedr.2022.101764>.

References

- Ammar, A., Brach, M., Trabelsi, K., et al., 2020. Effects of COVID-19 home confinement on physical activity and eating behaviour Preliminary results of the ECLB-COVID19 international online-survey. *Nutrients*.
- Bauer, T., Göhlmann, S., Sinning, M., 2007. Gender differences in smoking behavior. *Health Econ.* 16 (9), 895–909.
- Buyse, D.J., Reynolds, C.F., Charles, F., Monk, T.H., Berman, S.R., Kupfer, D.J., 1989. The Pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 28 (2), 193–213.
- Cavelaars, A.E.J.M., Kunst, A.E., Geurts, J.J.M., et al., 2000. Educational differences in smoking: international comparison. *BMJ* 320 (7242), 1102–1107.
- de Bruijn, E.-J., Antonides, G., 2020. Determinants of financial worry and rumination. *J. Econ. Psychol.* 76.
- Esai, Selvan M., 2020. Risk factors for death from COVID-19. *Nat. Rev. Immunol.* 20 (7), 407.
- Fottrell, Q. Stimulus checks played a major role in nationwide decline in anxiety and depression, new analysis says. *MSN2021*.
- French, D., McKillop, D., 2017. The impact of debt and financial stress on health in Northern Irish households. *Journal of European Social Policy.* 27 (5), 458–473.
- Galasso, V., Pons, V., Profeta, P., Becher, M., Brouard, S., Foucault, M., 2020. Gender differences in COVID-19 attitudes and behavior: Panel evidence from eight countries. *PNAS* 117 (44), 27285–27291.
- Government of Canada. Tobacco Use Statistics: Terminology. <https://www.canada.ca/en/health-canada/services/health-concerns/tobacco/research/tobacco-use-statistics/terminology.html>. Published 2008. Updated 2008-11-28. Accessed.
- Grzywacz, J.G., Almeida, D.M., 2008. Stress and binge drinking: A daily process examination of stressor pile-up and socioeconomic status in affect regulation. *Int. J. Stress Manage.* 15 (4), 364–380.
- Hall, M.H., Matthews, K.A., Kravitz, H.M., et al., 2009. Race and financial strain are independent correlates of sleep in midlife women: The SWAN sleep study. *Sleep* 32 (1), 73–82.
- Han, Q., Zheng, B., Agostini, M., Bélanger, J.J., Gützkow, B., Kreienkamp, J., Reitsema, A.M., van Breen, J.A., Collaboration, PsyCorona, Leander, N.P., 2021. Associations of risk perception of COVID-19 with emotion and mental health during the pandemic. *J. Affect. Disord.* 284, 247–255.
- Heck, R.H., Thomas, S.L., 2015. An introduction to multilevel modeling techniques: MLM and SEM approaches using Mplus, 3rd ed. Routledge.
- Hu, F.B., Stamper, M.J., Manson, J.E., Grodstein, F., Colditz, G.A., Speizer, F.E., Willett, W.C., 2000. Trends in the incidence of coronary heart disease and changes in diet and lifestyle in women. *N. Engl. J. Med.* 343 (8), 530–537.
- Huang, Y., Zhao, N., 2020. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res.* 288.
- Jin, S., Balliet, D., Romano, A., Spadaro, G., van Lissa, C.J., Agostini, M., Bélanger, J.J., Gützkow, B., Kreienkamp, J., Leander, N.P., Abakoumkin, G., Khaiyom, J.H.A., Ahmedi, V., Akkas, H., Almenara, C.A., Kurapov, A., Atta, M., Bagci, S.C., Basel, S., Kida, E.B., Buttrick, N.R., Chobthamkit, P., Choi, H.-S., Cristea, M., Csaba, S., Damjanovic, K., Danyliuk, I., Dash, A., Di Santo, D., Douglas, K.M., Enea, V., Faller, D.G., Fitzsimons, G., Gheorghiu, A., Gómez, A., Han, Q., Helmy, M., Hudiyana, J., Jeronimus, B.F., Jiang, D.-Y., Jovanović, V., Kamenov, Ž., Kende, A., Keng, S.-L., Kieu, T.T.T., Koc, Y., Kovyazina, K., Kozytska, I., Krause, J., Kruglanski, A.W., Kutlaca, M., Lantos, N.A., Lemay, E.P., Lesmana, C.B.J., Louis, W.R., Lueders, A., Malik, N., Martinez, A., McCabe, K.O., Mehulić, J., Milla, M.N., Mohammed, I., Molinaro, E., Moyano, M., Muhammad, H., Mula, S., Muluk, H., Myroniuk, S., Najafi, R., Nisa, C.F., Nyúl, B., O'Keefe, P.A., Osuna, J.J.O., Osin, E.N., Park, J., Pica, G., Pierro, A., Rees, J., Reitsema, A.M., Resta, E., Rullo, M., Ryan, M.K., Samekin, A., Santtila, P., Sasin, E., Schumpe, B.M., Selim, H.A., Stanton, M.V., Stroebe, W., Sultana, S., Sutton, R.M., Tseliou, E., Utsugi, A., van Breen, J.A., Van Veen, K., van Dellen, M.R., Vázquez, A., Wollast, R., Yeung, V.-L., Zand, S., Žeželj, I. L., Zheng, B., Zick, A., Zúñiga, C., 2021. Intergenerational conflicts of interest and prosocial behavior during the COVID-19 pandemic. *Personality Individ. Differ.* 171.
- Kassel, J.D., Stroud, L.R., Paronis, C.A., 2003. Smoking, stress, and negative affect: Correlation, causation, and context across stages of smoking. *Psychol. Bull.* 129 (2), 270–304.
- LeBreton, J.M., Senter, J.L., 2008. Answers to 20 questions about interrater reliability and interrater agreement. *Org. Res. Meth.* 11 (4), 815–852.
- Marshall, A.L., Smith, B.J., Bauman, A.E., Kaur, S., 2005. Reliability and validity of a brief physical activity assessment for use by family doctors. *Br. J. Sports Med.* 39 (5), 294.
- National Health and Nutrition Examination Survey Questionnaire. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2017–2018. https://wwwnchs.cdc.gov/Nchs/Nhanes/2017-2018/SMQRTU_J.htm.
- Nelson, N.A., Bergeman, C.S., 2020. Daily stress processes in a pandemic: The effects of worry, age, and affect. *The Gerontologist*.
- Nguyen, H.T., Do, B.N., Pham, K.M., Kim, G.B., Dam, H.T.B., Nguyen, T.T., Nguyen, T.T.P., Nguyen, Y.H., Sørensen, K., Pleasant, A., Duong, T.V., 2020. Fear of COVID-19 Scale: Associations of its scores with health literacy and health-related behaviors among medical students. *Int. J. Environ. Res. Public Health* 17 (11), 4164.
- Nguyen, H.C., Nguyen, M.H., Do, B.N., Tran, C.Q., Nguyen, T.T.P., Pham, K.M., Pham, L.V., Tran, K.V., Duong, T.T., Tran, T.V., Duong, T.H., Nguyen, T.T., Nguyen, Q.H., Hoang, T.M., Nguyen, K.T., Pham, T.T.M., Yang, S.-H., Chao, J.-J., Duong, T.V., 2020. People with suspected COVID-19 symptoms were more likely depressed and had lower health-related quality of life: The potential benefit of health literacy. *J. Clin. Med.* 9 (4), 965.
- Nisa, C.F., Bélanger, J.J., Faller, D.G., Buttrick, N.R., Mierau, J.O., Austin, M.M.K., Schumpe, B.M., Sasin, E.M., Agostini, M., Gützkow, B., Kreienkamp, J., Abakoumkin, G., Abdol Khaiyom, J.H., Ahmedi, V., Akkas, H., Almenara, C.A., Atta, M., Bagci, S.C., Basel, S., Kida, E.B., Bernardo, A.B.I., Chobthamkit, P., Choi, H.-S., Cristea, M., Csaba, S., Damjanovic, K., Danyliuk, I., Dash, A., Di Santo, D., Douglas, K.M., Enea, V., Fitzsimons, G., Gheorghiu, A., Gómez, A., Grzymala-Moszczynska, J., Hamaidia, A., Han, Q., Helmy, M., Hudiyana, J., Jeronimus, B.F., Jiang, D.-Y., Jovanović, V., Kamenov, Ž., Kende, A., Keng, S.-L., Kieu, T.T.T., Koc, Y., Kovyazina, K., Kozytska, I., Krause, J., Kurapov, A.W., Kurapov, A., Kutlaca, M., Lantos, N.A., Lemay, E.P., Lesmana, C.B.J., Louis, W.R., Lueders, A., Malik, N.I., Martinez, A., McCabe, K.O., Mehulić, J., Milla, M.N., Mohammed, I., Molinaro, E., Moyano, M., Muhammad, H., Mula, S., Muluk, H., Myroniuk, S., Najafi, R., Nyúl, B., O'Keefe, P.A., Osuna, J.J.O., Osin, E.N., Park, J., Pica, G., Pierro, A., Rees, J., Reitsema, A.M., Resta, E., Rullo, M., Ryan, M.K., Samekin, A., Santtila, P., Selim, H. A., Stanton, M.V., Sultana, S., Sutton, R.M., Tseliou, E., Utsugi, A., van Breen, J.A., Van Lissa, C.J., Van Veen, K., van Dellen, M.R., Vázquez, A., Wollast, R., Yeung, V.-L., Zand, S., Žeželj, I.L., Zheng, B., Zick, A., Zúñiga, C., Leander, N.P., 2021. Lives versus Livelihoods? Perceived economic risk has a stronger association with support for COVID-19 preventive measures than perceived health risk. *Sci. Rep.* 11 (1).
- Nwachukwu, I., Nkire, N., Shalaby, R., Hrabok, M., Vuong, W., Gusnowski, A., Surood, S., Urichuk, L., Greenshaw, A.J., Agyapong, V.I.O., 2020. COVID-19 pandemic: Age-related differences in measures of stress, anxiety and depression in Canada. *Int. J. Environ. Res. Public Health* 17 (17), 6366.
- Onder, G., Rezza, G., Brusaferro, S., 2020. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA* 323 (18), 1775–1776.
- Park, C.L., Iacocca, M.O., 2014. A stress and coping perspective on health behaviors: theoretical and methodological considerations. *Anxiety, Stress, & Coping.* 27 (2), 123–137.
- Rodriguez, L.M., Litt, D.M., Stewart, S.H., 2020. Drinking to cope with the pandemic: The unique associations of COVID-19-related perceived threat and psychological distress to drinking behaviors in American men and women. *Addict. Behav.* 110.
- Rodríguez-Pérez, C., Molina-Montes, E., Verardo, V., Artacho, R., García-Villanova, B., Guerra-Hernández, E.J., Ruíz-López, M.D., 2020. Changes in dietary behaviours during the COVID-19 outbreak confinement in the Spanish COVIDiet study. *Nutrients*. 12 (6), 1730.
- Romano, A., Spadaro, G., Balliet, D., Joireman, J., Van Lissa, C.J., Jin, S., Leander, N.P., 2020. Cooperation and trust across societies during the COVID-19 pandemic. *J. Cross-Cult. Psychol.* in press.
- Siahpush, M., Borland, R., Scollo, M., 2003. Smoking and financial stress. *Tobacco Control*. 12 (1), 60.
- Smith, P.C., Schmidt, S.M., Allensworth-Davies, D., Saitz, R., 2009. Primary Care Validation of a Single-Question Alcohol Screening Test. *J. Gen. Intern. Med.* 24 (7), 783–788.
- Stang, P., Lydick, E., Silberman, C., Kempel, A., Keating, E.T., 2000. The Prevalence of COPD. *Chest* 117 (5), 354S–359S.
- Thornton, J.S., Frémont, P., Khan, K., et al., 2016. Physical activity prescription: a critical opportunity to address a modifiable risk factor for the prevention and management of chronic disease: a position statement by the Canadian Academy of Sport and Exercise Medicine. *Br. J. Sports Med.* 50 (18), 1109.
- Thorsteinsson, E.B., Brown, R.F., Owens, M.T., 2019. Modeling the Effects of Stress, Anxiety, and Depression on Rumination, Sleep, and Fatigue in a Nonclinical Sample. *J. Nerv. Ment. Dis.* 207 (5), 355–359.
- United Nations. World Population Prospects. <https://population.un.org/wpp/DefinitionOfRegions/> Published 2019. Accessed May 2, 2021, 2021.
- Wang, X., Lei, S.M., Le, S., Yang, Y., Zhang, B., Yao, W.u., Gao, Z., Cheng, S., 2020. Bidirectional influence of the COVID-19 pandemic lockdowns on health behaviors and quality of life among Chinese adults. *Int. J. Environ. Res. Public Health* 17 (15), 5575.
- Wardle, J., Haase, A.M., Steptoe, A., Nillapun, M., Jonwutiwes, K., Bellis, F., 2004. Gender differences in food choice: The contribution of health beliefs and dieting. *Ann. Behav. Med.* 27 (2), 107–116.
- Wilsnack, R.W., Wilsnack, S.C., Gmel, G., Kantor, L.W., 2018. Gender differences in binge drinking: Prevalence, predictors, and consequences. *Alcohol Research: Current Reviews.* 39 (1), 57–76.
- World Health Organization. Coronavirus disease (COVID-19) pandemic. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>. Published 2021. Accessed March 3, 2021, 2021.
- Zheng, Z., Peng, F., Xu, B., et al., 2020. Risk factors of critical & mortal COVID-19 cases: A systematic literature review and meta-analysis. *J. Infect.* 81 (2), e16–e25.