

A health economic outcome evaluation of an internet-based mobile-supported stress management intervention for employees

Ebert, David Daniel; Kählke, Fanny; Buntrock, Claudia; Berking, Matthias; Smit, Filip; Heber, Elena; Baumeister, Harald; Funk, Burkhardt; Riper, Heleen; Lehr, Dirk

Published in:
Scandinavian Journal of Work, Environment and Health

DOI:
[10.5271/sjweh.3691](https://doi.org/10.5271/sjweh.3691)

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for pulished version (APA):
Ebert, D. D., Kählke, F., Buntrock, C., Berking, M., Smit, F., Heber, E., Baumeister, H., Funk, B., Riper, H., & Lehr, D. (2018). A health economic outcome evaluation of an internet-based mobile-supported stress management intervention for employees. *Scandinavian Journal of Work, Environment and Health*, 44(2), 171-182. <https://doi.org/10.5271/sjweh.3691>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Scand J Work Environ Health 2018;44(2):171-182
Published online: 16 Nov 2017, Issue date: 01 Mar 2018

doi:10.5271/sjweh.3691

A health economic outcome evaluation of an internet-based mobile-supported stress management intervention for employees

by Ebert DD, Kählke F, Buntrock C, Berking M, Smit F, Heber E, Baumeister H, Funk B, Riper H, Lehr D

Occupational internet-based stress management interventions (iSMI) have shown to be effective, but evidence for cost-effectiveness is scarce. We provide evidence for the cost-effectiveness of a guided iSMI from the employer's perspective and show a high probability that this intervention is associated with cost savings due to less absenteeism and greater productivity at work. Offering iSMI represents good value for money in occupational healthcare.

Affiliation: Friedrich-Alexander University Erlangen-Nuremberg, Department of Clinical Psychology and Psychotherapy, Nögelsbachstrasse 25a, Erlangen, 91052, Germany. david.ebert@fau.de

Refers to the following texts of the Journal: 2016;42(5):382-394
2015;41(2):164-174 2014;40(6):582-596 2006;32(1):1-4

Key terms: CBT; cost-benefit analysis; cost-effectiveness analysis; e-health; economic evaluation; evaluation; internet-based intervention; m-health; randomized controlled trial; RCT; stress; stress management; stress management intervention

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/29144535

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

A health economic outcome evaluation of an internet-based mobile-supported stress management intervention for employees

by David Daniel Ebert, PhD,^{1,2} Fanny Kählke, MA Ed MPH,¹ Claudia Buntrock, MSC,^{1,2} Matthias Berking, PhD,^{1,2} Filip Smit, PhD,^{3,4,5} Elena Heber, PhD,^{2,6} Harald Baumeister, PhD,⁷ Burkhardt Funk, PhD,² Heleen Riper, PhD,^{2,5} Dirk Lehr, PhD²

Ebert DD, Kählke F, Buntrock C, Berking M, Smit F, Heber E, Baumeister H, Funk B, Riper H, Lehr D. A health economic outcome evaluation of an internet-based mobile-supported stress management intervention for employees. *Scand J Work Environ Health*. 2018;44(2):171–182. doi:10.5271/sjweh.3691

Objective This study aimed to estimate and evaluate the cost-effectiveness and cost-benefit of a guided internet- and mobile-supported occupational stress-management intervention (iSMI) for employees from the employer's perspective alongside a randomized controlled trial.

Methods A sample of 264 employees with elevated symptoms of perceived stress (Perceived Stress Scale, PSS-10 ≥ 22) was randomly assigned either to the iSMI or a waitlist control (WLC) group with unrestricted access to treatment as usual. The iSMI consisted of seven sessions of problem-solving and emotion-regulation techniques and one booster session. Self-report data on symptoms of perceived stress and economic data were assessed at baseline, and at six months following randomization. A cost-benefit analysis (CBA) and a cost-effectiveness analysis (CEA) with symptom-free status as the main outcome from the employer's perspective was carried out. Statistical uncertainty was estimated using bootstrapping (N=5000).

Results The CBA yielded a net-benefit of €181 [95% confidence interval (CI) -6043–1042] per participant within the first six months following randomization. CEA showed that at a willingness-to-pay ceiling of €0, €1000, €2000 for one additional symptom free employee yielded a 67%, 90%, and 98% probability, respectively, of the intervention being cost-effective compared to the WLC.

Conclusion The iSMI was cost-effective when compared to WLC and even lead to cost savings within the first six months after randomization. Offering stress-management interventions can present good value for money in occupational healthcare.

Key terms CBT; cost-benefit analysis; cost-effectiveness analysis; economic evaluation; e-health; internet-based intervention; m-health; randomized controlled trial; RCT.

Work-related stress has shown to be a major risk factor for a range of adverse health outcomes, such as depression (1), coronary disease (2), and related mortality (3). It is also associated with reduced workplace safety (4) and substantial economic costs due to higher sick leave, lower work performance and staff turnover (5, 6). Absenteeism and presenteeism have found to be the major cost-drivers, often exceeding the costs of treatment (7, 8).

The efficacy of interventions for occupational stress has been demonstrated in a large number of randomized

controlled trials (RCT) (9, 10). Despite the availability of evidence-based interventions, however, the majority of stressed employees do not use stress interventions to improve their distress.

Using the internet to provide self-help interventions may help to overcome some of the limitations of traditional stress-management interventions (SMI), such as limited availability, high threshold for participation and delivery costs (11, 12). A large amount of studies has shown that internet-based interventions can be effective

¹ Department of Clinical Psychology and Psychotherapy, University of Erlangen-Nuremberg, Germany.

² Division of Online Health Training, Innovation Incubator, Leuphana University of Lüneburg, Lüneburg, Germany.

³ Department of Public Mental Health, Trimbos-Institute (Netherlands Institute of Mental Health and Addiction), Utrecht, The Netherlands.

⁴ Department of Epidemiology and Biostatistics, Amsterdam Public Health Research Institute, VU Medical Centre, Amsterdam, The Netherlands.

⁵ Department of Clinical, Neuro- and Developmental Psychology, VU University, Amsterdam, The Netherlands.

⁶ GET.ON Institut, Hamburg, Germany.

⁷ Department of Clinical Psychology and Psychotherapy, Institute of Psychology and Education, University of Ulm, Ulm, Germany.

(13) in clinical populations, such as depression (14–17), anxiety (18), and risky alcohol use (19) as well as in the field of prevention of mental disorders (20–23). Recently a number of trials showed that internet- and mobile-based interventions can be a valuable addition in occupational health including the treatment of insomnia among employees with work-related rumination (12, 24, 25), improving work-related stress (26–33) and the prevention and treatment of depression in workers (11, 34, 35). However, one trial revealed no significant results of internet-based interventions on depression in employees (36–38).

Whereas the efficacy of both face-to-face as well as internet-based approaches for managing work-related stress is well-documented, high quality data on whether such approaches provide good value for money is scarce. The most recent systematic review on economic analyses of occupational health programs was not able to identify any economic evaluation based on a randomized controlled design that focused on stress-management among workers (39).

As resources are restricted, however, employers are not just interested in the effectiveness of such an intervention, but also in the return-on-investment (ROI) (40). However, various intervention outcomes (ie, treatment response) cannot be easily monetarized and cannot thus be included in ROI analyses. Therefore, cost-effectiveness analyses (CEA) that compare the incremental costs and effects of the intervention and the comparator are also important (41).

Hence, the aim of the present study is to evaluate the cost-effectiveness and net-benefits of an internet-based SMI (iSMI) compared to a waitlist control (WLC) condition with unrestricted access to treatment as usual among employees with heightened levels of perceived stress. Analyses in the present study are conducted from the employer's perspective focusing on costs and savings that are relevant to the employer (ie, productivity losses).

Method

Design

A two-armed RCT was conducted to compare a guided iSMI (GET.ON Stress) with a WLC. Both conditions had full access to treatment as usual. Assessments took place at baseline (T1), post-treatment (7 weeks, T2), and 6-month follow-up. Participants were primarily recruited via the occupational health program of a large health insurance company in Germany. Recruitment was directed at the general working population and not restricted to members of the healthcare insurance company. It occurred through announcements on the health

care insurance company's website, newspaper articles and advertisements in the membership magazine of the insurance company. Moreover, the insurance company's occupational health management workers informed human resource departments of collaborating small- and medium-sized companies about the possibility for their employees to participate in the trial.

The present health economic evaluation followed guidelines from the ISPOR RCT-CEA Task Force Report and the recommendations of the Consolidated Health Economic Evaluation Reporting Standard (CHEERS) (42, 43). The ISPOR RCT-CEA Task Force report addresses issues related to trial design, selecting data elements, database design and management, analysis, and reporting of results. The aim of the CHEERS statement is to provide recommendations, in the form of a checklist, to optimize reporting of health-economic evaluations. A more detailed description of the study's design can be found in the study protocol (29). In brief, this study was designed as a health-economic evaluation of an occupational stress-management intervention in employees alongside a randomized controlled trial from the employer's perspective.

The sample consisted of 264 employees in Germany, who were randomly allocated with a ratio of 1:1 to the intervention (iSMI) or WLC group. An independent statistician performed the randomization with a web-based program (RandList). Both groups had full access to treatment as usual. Online-assessments took place at baseline (T1), post-treatment (seven weeks, T2), and 6-month follow up (T3).

Participants were primarily recruited via the occupational health program of a large health insurance company in Germany. Recruitment was directed at the general working population and not restricted to those who were insured by the healthcare insurance company. Recruitment was facilitated by announcements on the healthcare insurance company's website, newspaper articles and advertisements in the membership magazine of the insurance company. Moreover, the insurance company's occupational health management workers informed human resource departments of collaborating small- and medium-sized companies about the possibility for their employees to participate in the trial.

Participants were included in the study if they were 18 years or older, currently employed and scored ≥ 22 on the perceived stress scale (PSS-10). A cut-off value of one standard deviation above the mean PSS-10 of 15.3 [standard deviation (SD) 6.2] in a large working population (44) was chosen. The exclusion criteria were suicide risk, and presence of dissociative and / or psychotic symptoms. The Ethics Committee of the Philipps-University of Marburg, Germany, approved the study. The trial was registered (DRKS00004749) in the German clinical trial registry.

Intervention

The iSMI (called “GET.ON Stress”) consisted of seven sessions and one booster session four weeks after training completion. The primary aim of this intervention was to reduce stress levels in employees. GET.ON Stress is based on Lazarus’ transactional model of stress (45) (for a detailed description of the intervention see (29, 46). It consists of two main components: problem solving and emotion regulation. Problem solving is an evidence-based method that has been proven to be successful in improving mental health (47). On the other hand, employees are also frequently faced with situations that require dealing with unsolvable problems; such situations are often associated with strong negative affective reactions and require effective regulation strategies. Numerous studies indicate that deficits in emotion regulation may be a relevant factor for the development and persistence of mental health symptoms (31, 32). Targeting emotion regulation skills has shown to be promising for reducing a broad range of psychopathological symptoms (48) and a mechanism of change in previous studies on GET.ON Stress (32). While problem-focused coping by means of problem-solving techniques is a well-established component of most CBT stress management trainings, the emotion-focused ways of coping could be regarded as the forgotten component. The intervention consists of eight sessions composed of modules for psycho-education (session 1), problem solving (sessions 2–3), emotion regulation (sessions 4–6), planning for the future (session 7) and a booster session (session 8). Additionally, participants are offered 11 modules that are integrated in sessions 2–6 and that can be chosen based on individual need and/or preference. Additional modules are directed at time management, rumination and worrying, psychological detachment from work, sleep hygiene, rhythm and regularity of sleeping habits, nutrition and exercise, organization of breaks during work, and social support (see table 1 for a session overview). Each session can be completed in approximately 45–60 minutes. We advise participants to do at least one and maximum two sessions a week. Consequently, the training lasts about 4–7 weeks (plus a booster session after 4 weeks). Once participants finish a lesson, they need to wait until they receive feedback from the accompanying e-coach to be able to start with the next lesson. Lessons consist of texts, exercises, and testimonials and also include interactive elements such as audio and video clips. Participants are encouraged to keep a daily online stress diary. A strong focus of the intervention lies on transfer tasks (homework assignments) to integrate newly acquired strategies and techniques into daily life. The training is adaptive as the content is tailored to the specific needs of the individual participant by continuously asking participants to choose among various response options. Subsequent content is then tailored to each participant’s

response. Using responsive web-design, participants can follow the program on the internet, a tablet or mobile phone. An integrated read-aloud function allows participants to follow narrated lessons. If desired, participants received automatic motivational text messages and small exercises on their mobile phones. These messages aim to support participants in transferring the exercises of the training into their daily lives (eg, short relaxation exercises: “Relax your muscles in your hands and arms for 3 seconds now. Follow your breathing and each time you breathe out, relax a little more.”). The participants had the opportunity to choose between “light coach” (one text message every other day) and “intensive coach” (2–3 text messages every day) options. Participants were supported by an e-coach holding a degree in psychology who followed a standardized manual for writing feedback. The e-coaches provided personalized feedback throughout the intervention. Participants completed on average 5.7 sessions (SD 2.32), used the intervention for a mean of 8.27 weeks (SD 8.54, range 0–56), and 93/132 (70.5%) completed all seven sessions. Ten participants (7.6%) did not start the intervention due to lack of time or changes in personal circumstances. Every participant received a maximum of 30 minutes of feedback per session of an e-coach, resulting in a maximum total of 3 hours (rounded) per participant over the whole course.

Outcome measures

Clinical outcome. The primary outcome of the CEA was the number of participants who achieved symptom-free status as measured by the PSS-10 (49). The PSS consists of 10 items rated on a 5-point Likert scale (0=never; 1=almost never; 2=sometimes; 3=fairly often; 4=very often; range 0–40) referring to the past week. Cronbach alphas ranged from 0.70–0.91 over different measurement points in this study (32). The symptom-free status was achieved when participants scored >2 SD below the mean (T1) of the stressed population (25.52, SD 3.91) as according to Jacobson & Truax (30, 50).

Cost outcome. Only costs that were directly relevant for the employer were considered: (i) intervention costs, (ii) costs, due to absenteeism, and (iii) costs due to presentism. Data on productivity losses (absenteeism and presentism) due to health problems were assessed at baseline and 6-month follow-up using the relevant module of the “Trimbos Institute and Institute of Medical Technology Questionnaire for Costs Associated with Psychiatric Illness” (TiC-P) adapted to the German context (51). The TiC-P is a self-report questionnaire with a 3-month recall period in the current study. It is a widely used (24, 29, 32, 33, 52) instrument for collecting data on healthcare utilization and productivity losses in patients with mild-to-moderate mental health conditions

(53). The TiC-P has shown a good test–retest reliability indicated by a satisfactory Cohen’s kappa for most items related to healthcare consumption and items absent from work as well as presenteeism (54). Cumulated costs over the six months of the trial were estimated using the area under curve (AUC) method to linearly interpolate 3-month costs as measured at each measurement point to cover the full follow-up period of six months (54).

$$AUC = \left(\frac{\frac{\text{Costs Baseline}}{3} + \frac{\text{Costs 6 months}}{3}}{2} \right) * 3 + \text{Costs 6 months}$$

All costs were calculated in euros (€) and indexed for the reference year 2013 (index factor 1.04) based on the German consumer price index (55).

Absenteeism

In order to measure absenteeism, participants reported how many days they had been absent from work during the previous three months (lost work days). Absenteeism cost were calculated by applying the human capital approach (53). In doing so, the number of lost work days was multiplied by the participant’s average gross daily wage based on their reported monthly salary.

Presenteeism

Presenteeism occurs when people feel ill and therefore are less efficient during work. In order to quantify the costs due to presenteeism, the participants reported both the number of days that they felt ill but continued to work and their personal inefficiency score on a scale from 0–10, where 0 reflects total inefficiency and 10 reflects that they were as efficient as when in good health. Lost work days due to presenteeism were computed by multiplying the number of work days with reduced functioning by the corresponding inefficiency score. This method is called the Osterhaus method (56). Subsequently, based on self-reported monthly salary, their gross wages per hour were calculated and used to calculate the costs that occurred due to presenteeism.

Costs of the intervention

The provider of GET.ON Stress estimated the average price of the intervention to be €299 per participant. This flat tariff contains all costs for developing, providing, website hosting and maintenance, as well as guidance and 19% VAT. Opportunity costs were not included since we assumed that participants would want to use the intervention at home and not at work.

Statistical analysis

This health economic evaluation was conducted alongside a randomized trial powered to detect a mean standardized difference of $d=0.35$ in the primary outcome (PSS-10) between the groups at post-measurement. However, it is worth noting that cost data usually have a large variance, which would require very large samples to test economic hypotheses. Such large samples are most likely prohibitive from a financial, logistical and medical-ethics point of view. Therefore, the study was not powered to statistically test differences in economic outcomes. Instead, a probabilistic decision-making approach for making health-economic inferences was used (57). This procedure takes the uncertainty about key parameters (in costs and effects) into account (58, 59) and informs decision-makers on probabilities rather than statistical significance. Perfect information is only available in very few decision-making situations. Probabilities account for the amount of risk each decision carries.

All analyses were conducted in accordance with the intention-to-treat (ITT) principle. All 264 participants completed the PSS and the cost measures at baseline. The missing data was assumed to be missing completely at random as indicated by the Little’s overall test of randomness. Missing cost data at T3 (wage, weekly working hours, presenteeism days, absenteeism days) were imputed using the regression imputation procedure in Stata version 13 to obtain predicted values of missing data. Predictors of outcome and dropout were identified by (logistic) regression. Baseline costs of presenteeism and absenteeism, wage, working hours as well as demographic variables, including age, and gender were found to be significant predictors for both categories and were thus included as predictors in the imputation model.

Cost–benefit analysis (CBA)

This study proceeds on the assumption that the employer is the one who pays for the intervention and benefits from cost savings due to decreased presenteeism and absenteeism. Benefits were defined as the difference in total monetarized outcome measures (ie, absenteeism and presenteeism) between study conditions. Positive benefits point to reduced spending. Costs were defined as the intervention costs, which are assumed to be paid by the employer for each employee (estimated at €299). This amount represents the current market price for the complete intervention, which includes access to seven modules + one booster session for 12 months, one individual written feedback per completed lesson and replies to individual questions from the participant on demand until completion of the booster session.

In the present study, the following metrics for costs and benefits are reported: (i) the net benefits (NB =

benefits - costs), the money saved after cost deduction; (ii) the benefit-to-cost ratio ($BCR = \text{benefit/cost}$) to measure the benefits against the costs; and (iii) the return-on-investment ($ROI = \text{NB/costs}$), the amount in euro returned per one euro invested (40, 60).

Statistical uncertainty is taken into account by bootstrapped 95% confidence intervals (95% CI) which are estimated around these measures with 5000 replications using the bias-corrected and accelerated bootstrap procedure as implemented in Stata version 13 (40, 61). Cost savings are indicated when following criteria are met: $NB > 0$, $BCR > 1$, and $ROI > 1$ (40, 62).

Cost-effectiveness analysis

For the CEA, costs and cost offsets (presenteeism + absenteeism + intervention costs) and effects (number of participants with symptom-free status) were calculated for the 6-month period. Subsequently, costs and outcomes were combined into the incremental cost-effectiveness ratio (ICER) using the following formula:

$$ICER = \frac{\text{Cost iSMI} - \text{Cost WLC}}{\text{Effect iSMI} - \text{Effect WLC}}$$

Since cost data tend to be skewed to the right (58), we used non-parametric bias-corrected accelerated bootstrapping resampling techniques (with 5000 replications) to assess uncertainties surrounding the ICER. The bootstrap analyses were undertaken in the context of Stata's procedure for seemingly unrelated regression equations (SURE) to simultaneously assess incremental costs and incremental effects. The generated ICER were plotted in a cost-effectiveness plane. On the plane, incremental effects between intervention group (IG) and control group (CG) are depicted on the x-axis and the incremental costs between IG and CG are depicted on the y-axis.

If the majority of simulated ICER fall in the south-east quadrant, the intervention is considered to "dominate" the active control groups because better effects are obtained for lower costs. In contrast, in the north-west quadrant the intervention is considered "inferior" to the control group since it is associated with higher costs and worse health outcomes and therefore not considered to be cost-effective (41). ICER in the south-west quadrant point to an intervention being less effective but also less costly than the control condition. Finally, ICER in the north-east quadrant reflect an intervention being more effective and more costly than control condition. In this case, the amount of money a decision-maker is willing to pay for one additional positive outcome is crucial for a new intervention to be adopted or not. The probability

that the intervention is cost-effective compared to CG at various willingness-to-pay (WTP) ceilings for one treatment response gained can be shown by the means of the cost-effectiveness acceptability curve (63).

Sensitivity analysis

The robustness of the findings was assessed by performing sensitivity analyses over a range of intervention costs. In the main analysis, we used intervention costs of €299. However, there is uncertainty concerning these costs, as intervention costs may differ once implemented into occupational routine care, for example due to changing demand, supply in the health sector, and preferences of employer and employees. Therefore, two additional sensitivity analyses were conducted assuming higher and lower intervention costs (with $\pm \text{€}100$).

Results

Sample

Table 1 presents the baseline characteristics; a comprehensive description can be found in the paper describing the primary efficacy results (30).

Study drop-outs

The study attrition rate was low: 10.6% (28/264) of participants did not complete the 6-month follow-up questionnaires [$N=17/132$ (12.8%) for the iSMI and $N=11/132$ (8.33%) in the WLC; $\chi^2=1.16$; $P>0.05$]. Persons who dropped out did not differ in a meaningful way from those who provided data, neither on baseline stress scores nor any other baseline outcomes, with the exception of worrying ($P<0.05$).

Clinical outcome

The effect outcome for the CEA was symptom-free status defined at <17.70 on the PSS-10. Therefore, at T3 79/132 (59.8%) in the iSMI were significantly more participants classified as symptom-free compared to 31/132 (23.5%) in the WLC [NNT (number needed to treat)=2.73; $\chi^2=15.0$; $P<0.001$]. On average, the iSMI improved by 9.75 (SD 6) PSS scores between pre- and 6-month follow-up whereas the WLC improved by 3.0 (SD 6) PSS scores. The analysis of covariance (ANCOVA) confirmed lower scores on the PSS-10 (relative to the WLC) were found in the iSMI group at T3 [$F_{1,261}=80.17$, $P<0.001$, with large between-group effect sizes ($d=1.02$; 95% CI 0.76–1.27)] (50).

Table 1. Demographic characteristics. [SD=standard deviations; iSMI=internet/mobile-supported stress management intervention; %=percentages at baseline; WLC=waitlist control; PSS=perceived stress scale.]

Characteristics	All (N=264)				iSMI (N=132)				WLC (N=132)			
	N	%	Mean	SD	N	%	Mean	SD	N	%	Mean	SD
Age			43.3	10.2			42.4	10.7			44.2	9.6
Gender (female),	193	73.1			97	73.5			96	72.7		
Married/partnership	160	60.6			80	60.6			80	60.6		
Experience												
Experience with health trainings	34	12.9			17	12.9			17	12.9		
Previous psychotherapy	95	36.0			52	39.4			43	32.6		
Current psychotherapy	16	6.1			5	3.8			11	8.3		
Work characteristics												
Years of work experience			18.1	11.1			17.2	10.8			18.9	11.2
Full-time employed	204	77.3			105	79.5			99	75.0		
Part-time employed	57	21.6			25	18.9			32	24.2		
On sick leave	3	1.1			2	1.5			1	0.8		
Effect and costs												
PSS-10			25.5	3.9			25.9	4			25.2	4
Presenteeism (euros)			1185	1455			1136	1363			1234	1545
Presenteeism (days)			16.6	16			16.0	14.9			17.3	17
Absenteeism (euros)			814	1756			837	1634			790	1876
Absenteeism (days)			4.7	9			5	8.7			4.4	9.3

Costs outcome

Table 2 shows mean presenteeism and absenteeism days, and similarly hourly wage for the 6-month follow-up assessment. In addition, cumulated costs for the intervention, presenteeism, and absenteeism as well as total cost per group and between-group cost differences are reported. Presenteeism cost (iSMI: €1346; WLC: €1655) caused slightly lower costs compared to absenteeism cost (iSMI: €1578.2; WLC: €1756.6) in both groups. The mean difference in indirect costs (iSMI–WLC) at the 6-month follow-up was €488 per person favoring the intervention group. In the intervention group (iSMI), each participant produced costs of €299 (intervention costs)

and therefore saved €189 (SD 434) on average compared to the control group (WLC) in the first six months.

Cost benefit analyses (CBA)

The results of the conducted CBA and sensitivity analyses are reported in table 3. Calculations, based on the intervention costs at €299 and costs per person of €3412 in the WLC and €3180 in the iSMI, yielded a net benefit of €181 and a benefit-to-cost ratio [BCR (benefit/costs)] of 1.6 (95% CI -1.2–4.5) after 6 months. Moreover, the [(benefits - costs)/(costs × 100)] was 0.6 (95% CI 2.2–3.5) meaning the employer gains €0.60 for every euro invested within the time span of six months.

Cost-effectiveness analyses (CEA)

Table 4 reports the results of the CEA. ICER ($C_1 - C_0 / (E_1 - E_0)$) is defined as the incremental difference of mean costs between two groups (CG - IG) divided by the incremental difference in positive effects, ie, the amount of symptom-free persons. The IG produced less indirect costs compared to the CG (€3223 versus €3412). Hence, the ICER formula $(322 - 3412) / (0.6 - 0.2)$ yields an adjusted point estimate of -€521 (95% CI -3123–1900). The negative ICER falls in the south-east quadrant is therefore dominant.

Figure 1 shows the cost-effectiveness plane in which each dot (N=5000) represents one of 5000 bootstrapped replicated ICER. The bulk of ICER fall in the south-east quadrant, indicating that there is a 67% probability that the iSMI generates greater health effects at lower costs compared to the WLC.

Supplementary figure 2 (www.sjweh.fi/show_

Table 2. Hourly wage, absenteeism, presenteeism, and related costs categorized by condition at 6-month follow-up. [IG=intervention group; CG=control group, WLC=waitlist control, iSMI=internet-based stress-management intervention; SD=standard deviation.]

	IG (iSMI) (N=132)		CG (WLC) (N=132)		Mean	SD
	Mean	SD	Mean	SD		
Hourly wage (euros) ^a	21.54	6.7	21.68	7.9		
Absenteeism days ^{b, a}	3.64	6.7	5.23	12.2		
Presenteeism days ^{b, a}	11.32	12.9	11.47	11.9		
Indirect costs (euros) ^c						
Absenteeism (euros) ^a	1578.18	1471	1756.35	1848		
Presenteeism (euros) ^a	1345.79	2184	1655.16	3436		
Intervention costs (euros)	299					
Total costs (euros)	3223	2787	3412	4133	-189	434

^a Missing data imputed by multiple imputation.

^b Regarding the last 3 months.

^c Cumulative costs for each participant during the 6-month follow-up period calculated by the area under the curve (AUC) of linearly interpolated 1-month costs.

Table 3. Results of the main and sensitivity analyses (based on 5000 bootstrap simulations) with N=132 per group. [CI=95% confidence interval, SA=sensitivity analyses; NB=net benefit; BCR=benefit cost ratio; ROI=return on investment]

Analysis	Costs	Benefits	Financial return					
	Total (euros)	Total (euros)	NB ^a	95% CI	BCR ^b	95% CI	ROI ^c	95% CI
Main analyses	299	488	181	-643–1042	1.6	-1.2–4.5	0.61	-2.2–3.5
SA1 (-100 euros)	199	488	281	-543–1142	2.4	-1.7–6.7	1.4	-2.8–5.7
SA2(+100 euros)	399	488	81	-743–942	1.2	-0.8–3.4	0.2	-1.9–2.4


^a Indicating amount of money (euros) gained after costs are recovered.

^b Indicating the amount of money (euros) the employer gains for every euro invested.

^c Indicating the percent of profit per euro invested.

Table 4. Cost-effectiveness analyses based on 5000 replicates of the incremental cost-effectiveness ratio (mean differences in costs from the employer's perspective and in symptom-free status). [CI=confidence interval, SA=sensitivity analysis IG=intervention group, CG=control group, ΔC= incremental costs, ΔE= incremental effects, ICER= incremental cost–effectiveness ratio, NE= north-east quadrant, SE=south-east quadrant, SW=south-west quadrant, NW=north-west quadrant]

Analysis	Sample size		Outcome (symptom-free status)	ΔC (euros)	ΔE (symptom-free status)	Euros/symptom-free status		Distribution cost effectiveness plane (%)			
	IG	CG				ICER	95% CI	NE	SE	SW	NW
Main analyses	132	132	PSS-10 <17.70	-188	0.36	-521	-3123–1900	33	67	0.0	0.0
SA1 (-100 euros)	132	132	PSS-10 <17.70	-328	0.36	-804	-3416–1601	25	75	0.0	0.0
SA2 (+100 euros)	132	132	PSS-10 <17.70	-88	0.36	-238	-2836–2206	42	58	0.0	0.0

**Figure 1.** Scatter plot showing the mean differences in costs and effect outcome (symptom-free status) data using 5000 bootstrap replications.

[abstract.php?abstract_id=3691](#)) presents the cost-effectiveness acceptability curve, which helps decision-makers decide if the intervention provides “good” value for money given varying willingness-to-pay thresholds. If the employer is willing to pay zero, respectively €500, €1000, €2000 for one additional symptom-free person, there is an 80%, 90% and 98% probability that the IG is more cost-effective than the WLC.

Sensitivity analyses

Both tables 3 and 4 demonstrate results based on the sensitivity analyses. The ICER increased up to -€804

when assuming lower treatment costs (SA1) and the net-benefit rose to €281. When assuming higher intervention costs (SA2), the ICER drops to -€238 but the intervention still has a 58% probability of gaining greater health effects at lower costs compared to the control condition.

Discussion

Main results

The present study took the employer's perspective to evaluate the health economic impacts of an internet-based mobile-supported SMI. Both CEA and CBA indicated that providing the intervention to employees with elevated symptoms of stress is likely to present good value for money in occupational healthcare. The CEA demonstrated a 67% likelihood that the intervention was more cost-effective than no immediate intervention. Likewise, our CBA indicated net-savings of €181 on average per participant already in the first six months following the intervention, with a return on investment of €0.61 per euro invested in the intervention. These savings are mostly due to reductions in absenteeism and presenteeism costs. The estimates can be seen as conservative, as costs due to absenteeism and presenteeism were calculated based on the employee's gross wage only. Costs that occur as a consequence of absenteeism and presenteeism such as reduced productivity and loss in earnings for the employer due to presenteeism and absenteeism were not taken into account. Sensitivity

analyses confirmed the robustness of the findings. If the employer is only willing to pay nothing or €500 for an additional symptom-free person, then there is a 67% and 82%, respectively, probability that the intervention is deemed more cost-effective compared to no immediate intervention.

Comparison with previous findings

This study is, to the best of our knowledge, the first to investigate the cost-effectiveness of a SMI from the employer's perspective. Our results compare favorably to the average ROI [-0.91 versus -0.22 (SD 2.41)] found in 12 RCT in the latest review (39) on the costs and benefits of health promotion at the workplace. The CEA are in line with a recent study on an internet-based intervention targeting insomnia in workers that found a net-benefit of €418 (95% CI -593.03–1488.70) per participant and an ROI of 208% (95% CI -296.52–744.35) (64). Interestingly, the reduction in costs in both studies was mainly driven by the effects of the intervention on presenteeism and to a lesser degree by reduced absenteeism. These findings emphasize the importance of taking indirect cost due to presenteeism into account when investigating health economic effects of interventions in the work-setting. However, our follow-up period was limited to six months only. As chronic stress is associated especially with poor economic consequences in the long run, longer follow-up periods may help to shed light on the true potential and cost savings of the intervention. Such an assumption is supported by one of the few modelling studies that modelled the potential cost-benefit of implementing mental health care interventions in the workplace. They found a net benefit of US\$30 per participating worker in year 1 and US\$257 in year two following the treatment (65). However, future studies are needed to determine the long-term economic impact of occupational mental health interventions.

Limitations

Limitations of our study include the fact that our large-scale trial was still underpowered for economic analyses. This is a common problem in health-economic evaluations conducted alongside a clinical trial, as cost variables often have a higher variance and generally require greater sample sizes than clinical evaluations (59). Like most randomized trials, the present study was only powered to detect findings on the primary clinical outcome, and in such cases a probabilistic decision-making approach is usually used (58, 59). However, once more studies focusing on a similar target are published, pooling techniques such as individual participant data meta-analyses may be used to overcome this power problem (66–68). Second, only severely distressed participants with high

baseline scores (PSS >22) were included in the study. Hence, the results may not apply to populations with lower stress levels and future studies should explore the cost-effectiveness probability for employees with lower stress-levels. Third, self-selection of the participants restrict the generalizability of results to employees willing to utilize such an intervention, with females being more likely to participate. Future studies should focus on reaching men for occupational SMI. Fourth, the assessment of costs due to absenteeism and presenteeism was based on self-reported data. Although this procedure has exhibit reasonably good reliability and validity (53, 69–71) and is standard procedure in health economic outcome evaluations alongside RCT (72), subjective data are prone to certain biases and hence the total costs in both groups may be either an overestimation or underestimation of true overall costs. Fifth, other work-related costs such as staff turnover, early retirement, and somewhat intangible costs such as the firm's reputation as a bad/good place to work were not included and could lead to greater cost-savings. Moreover, we used wages as a basis for measuring productivity costs, which might be a good proxy for true productivity costs only under certain conditions, as the method used ignores, for example, cases in which additional efforts of co-workers compensate productivity lost due to absenteeism and presenteeism of colleagues. Sixth, a WLC design with access to treatment as usual has been chosen where treatment as usual without delayed access is regarded as favorable control conditions (73). Finally, the present evaluation focused solely on costs relevant for the employer and excluded other costs that are relevant for the society, such as direct medical costs. Although indirect costs are often the major cost driver in stress-related health problems accounting for 50–80% of all associated costs (6, 7), future studies should also evaluate the cost-effectiveness from other perspectives, such as the societal or healthcare perspective.

Concluding remarks

There is substantial evidence for the effectiveness of psychological interventions for improving mental and work-related outcomes. However, methodological sound studies on the health economic effects of such interventions are still limited. The present study is the first showing that investing in internet-based stress-management interventions is associated with a high probability for a positive financial return, even when only focusing on costs that are relevant for the employer. In conclusion, when future studies could replicate the current findings, investing in digital interventions aiming to improve mental health at the workplace may be beneficial for employers as seen from a corporate health-economic and business point of view.

Acknowledgements

German healthcare insurance firm BARMER and the European Commission funded this study. The trial is registered with the German Clinical Trials Register (DRKS): DRKS00004749; www.drks.de/drks_web/ (archived by WebCite)

Declaration of interest

DDE, DL, MB and BF are stakeholders of the “Institute for Online Health Trainings”, which aims to transfer scientific knowledge related to the present research into routine healthcare.

References

1. Hammen C. Stress and depression. *Annu Rev Clin Psychol*. 2005 Jan;1:293–319. <https://doi.org/10.1146/annurev.clinpsy.1.102803.143938>.
2. Richardson S, Shaffer JA, Falzon L, Krupka D, Davidson KW, Edmondson D. Meta-analysis of perceived stress and its association with incident coronary heart disease. *Am J Cardiol*. 2012 Dec;110(12):1711–6. <https://doi.org/10.1016/j.amjcard.2012.08.004>.
3. Steptoe A, Kivimäki M. Stress and cardiovascular disease. *Nat Rev Cardiol*. 2012 Jun;9(6):360–70. <https://doi.org/10.1038/nrcardio.2012.45>.
4. CIPD (Chartered Institute of Personnel and Development). Building the business case for managing stress in the workplace. London; 2008.
5. Kalia M. Assessing the economic impact of stress—the modern day hidden epidemic. *Metabolism*. 2002 Jun;51(6 Suppl 1):49–53. <https://doi.org/10.1053/meta.2002.33193>.
6. Hassard J, Teoh K, Cox T, Dewe P, Cosmar M, Gründler R, et al. Calculating the costs of work-related stress and psychosocial risk - a literature review. European Agency for Safety and Health at Work. 2014.
7. Uegaki K, Bakker I, de Bruijne M, van der Beek A, Terluin B, van Marwijk H, et al. Cost-effectiveness of a minimal intervention for stress-related sick leave in general practice: results of an economic evaluation alongside a pragmatic randomised control trial. *J Affect*. 2010;120(1–3):177–87. <https://doi.org/10.1016/j.jad.2009.04.012>.
8. Rebergen DS, Bruinvels DJ, van Tulder MW, van der Beek AJ, van Mechelen W. Cost-Effectiveness of Guideline-Based Care for Workers with Mental Health Problems. *J Occup Environ Med*. 2009;51(3):313–22. <https://doi.org/10.1097/JOM.0b013e3181990d8e>.
9. Richardson KM, Rothstein HR. Effects of occupational stress management intervention programs: a meta-analysis. *J Occup Health Psychol*. 2008 Jan;13(1):69–93. <https://doi.org/10.1037/1076-8998.13.1.69>.
10. van der Klink JJ, Blonk RW, Schene AH, van Dijk FJ. The benefits of interventions for work-related stress. *Am J Public Health*. 2001 Feb;91(2):270–6. <https://doi.org/10.2105/AJPH.91.2.270>.
11. Junge MN, Lehr D, Bockting CLH, Berking M, Riper H, Cuijpers P, et al. For whom are internet-based Occupational Mental Health Interventions effective? Moderators of internet-based Problem-Solving Training Outcome. *Internet Interv*. Elsevier; 2015;2(1):39–47.
12. Ebert DD, Berking M, Thiart H, Riper H, Laferton J, Lehr D. Restoring depleted resources: Efficacy and mechanisms of change of an internet-based unguided recovery training for better sleep and psychological detachment from work. *Heal Psychol*. 2015;34S:1240–51. <https://doi.org/10.1037/hea0000277>.
13. Ebert DD, Daele T, Nordgreen T, Karekla M, Compare TA, Zarbo C, et al. internet and mobile-based psychological interventions: applications, efficacy and potential for improving mental health. A report of the EFPA e-health taskforce. *BMJ Open*. 2017;7:e016696.
14. Hedman E, Ljótsson B, Lindefors N. Cognitive behavior therapy via the internet: a systematic review of applications, clinical efficacy and cost-effectiveness. *Expert Rev Pharmacoecon Outcomes Res*. 2012 Dec;12(6):745–64. <https://doi.org/10.1586/erp.12.67>.
15. Ebert DD, Zarski A-C, Christensen H, Stikkelbroek Y, Cuijpers P, Berking M, et al. internet and computer-based cognitive behavioral therapy for anxiety and depression in youth: a meta-analysis of randomized controlled outcome trials. *PLoS One*. 2015 Jan;10(3):e0119895. <https://doi.org/10.1371/journal.pone.0119895>.
16. Ebert DD, Donkin L, Andersson G, Andrews G, Berger T, Carlbring P, et al. Does internet-based guided-self-help for depression cause harm? An individual participant data meta-analysis on deterioration rates and its moderators in randomized controlled trials. *Psychol Med*. 2016 Sep;46(13):2679–93. <https://doi.org/10.1017/S0033291716001562>.
17. Königbauer J, Letsch J, Doebler P, Ebert DD, Baumeister H. internet- and Mobile-based Depression Interventions for People with Diagnosed Depression: A Systematic Review and Meta-analysis. *J Affect Disord*. 2017 Jul;178(0):131–41.
18. Mayo-Wilson E, Montgomery P. Media-delivered cognitive behavioural therapy and behavioural therapy (self-help) for anxiety disorders in adults. *Cochrane database Syst Rev*. 2013 Jan;9:CD005330. <https://doi.org/10.1002/14651858.CD005330.pub4>.
19. Riper H, Blankers M, Hadiwijaya H, Cunningham J, Clarke S, Wiers R, et al. Effectiveness of Guided and Unguided Low-Intensity internet Interventions for Adult Alcohol Misuse: A Meta-Analysis. Stewart R, editor. *PLoS One*. 2014 Jan;9(6):e99912. <https://doi.org/10.1371/journal.pone.0099912>.
20. Buntrock C, Ebert DD, Lehr D, Smit F, Riper H, Berking M,

- et al. Effect of a Web-Based Guided Self-help Intervention for Prevention of Major Depression in Adults With Subthreshold Depression. *JAMA*. 2016 May 3;315(17):1854–63. <https://doi.org/10.1001/jama.2016.4326>.
21. Ebert DD, Buntrock C, Cuijpers P, K van Z, P C, C B, et al. Online Intervention for Prevention of Major Depression — Reply. *JAMA*. 2016 Aug;316(8):881. <https://doi.org/10.1001/jama.2016.9586>.
22. Sander L, Rausch L, Baumeister H. Effectiveness of internet-Based Interventions for the Prevention of Mental Disorders: A Systematic Review and Meta-Analysis. *JMIR Ment Health*. 2016 Aug 17;3(3):e38. <https://doi.org/10.2196/mental.6061>.
23. Ebert DD, Cuijpers P, Mu-oz RF, Baumeister H. Prevention of Mental Health Disorders using internet and mobile-based Interventions: a narrative review and recommendations for future research. *Front Psychiatry Frontiers*; 2017;8:116. <https://doi.org/10.3389/fpsy.2017.00116>.
24. Thiart H, Lehr D, Ebert D, Sieland B, Berking M, Riper H. Log in and breathe out: efficacy and cost-effectiveness of an online sleep training for teachers affected by work-related strain - study protocol for a randomized controlled trial. *Trials [internet]* . Springer Nature; 2013;14(1):169.
25. Thiart H, Lehr D, Ebert DD, Berking M, Riper H. Log in and breathe out: internet-based recovery training for sleepless employees with work-related strain {textendash} results of a randomized controlled trial. *Scand J Work Environ*. 2015 Jan;41(2):164–74. <https://doi.org/10.5271/sjweh.3478>.
26. Ly KH, Asplund K, Andersson G. Stress management for middle managers via an acceptance and commitment-based smartphone application: A randomized controlled trial. *internet Interv*. 2014 Jul;1(3):95–101. <https://doi.org/10.1016/j.invent.2014.06.003>.
27. Ruwaard J, Lange A, Bouwman M, Broeksteeg J, Schrieken B. E-mailed standardized cognitive behavioural treatment of work-related stress: a randomized controlled trial. *Cogn Behav Ther*. 2007 Jan;36(3):179–92. <https://doi.org/10.1080/16506070701381863>.
28. Aikens KA, Astin J, Pelletier KR, Levanovich K, Baase CM, Park YY, et al. Mindfulness goes to work: impact of an online workplace intervention. *J Occup Environ Med*. 2014;56(7):721–31. <https://doi.org/10.1097/JOM.0000000000000209>.
29. Heber E, Ebert DD, Lehr D, Nobis S, Berking M, Riper H. Efficacy and cost-effectiveness of a web-based and mobile stress-management intervention for employees: design of a randomized controlled trial. *BMC Public Health* . 2013 Jan;13:655. <https://doi.org/10.1186/1471-2458-13-655>.
30. Heber E, Lehr D, Ebert D, Berking M, Riper H. Web-based and mobile stress management intervention for employees: results of a randomised controlled trial. *J Med internet Res*. 2016;18(1):e21. <https://doi.org/10.2196/jmir.5112>.
31. Ebert DD, Lehr D, Smit F, Zarski A-C, Riper H, Heber E, et al. Efficacy and cost-effectiveness of minimal guided and unguided internet-based mobile supported stress-management in employees with occupational stress: a three-armed randomised controlled trial. *BMC Public Health* . 2014 Aug;14(1):807. <https://doi.org/10.1186/1471-2458-14-807>.
32. Ebert DD, Lehr D, Heber E, Riper H, Cuijpers P, Berking M. internet- and mobile-based stress management for employees with adherence-focused guidance: efficacy and mechanism of change. *Scand J Work Environ Health* . 2016 Sep;42(5):382–94. <https://doi.org/10.5271/sjweh.3573>.
33. Ebert DD, Heber E, Berking M, Riper H, Cuijpers P, Funk B, et al. Self-guided internet-based and mobile-based stress management for employees: Results of a randomised controlled trial. *Occup Environ Med*. 2016;73(5). <https://doi.org/10.1136/oemed-2015-103269>.
34. Imamura K, Kawakami N, Furukawa TA, Matsuyama Y, Shimazu A, Umanodan R, et al. Does internet-based cognitive behavioral therapy (iCBT) prevent major depressive episode for workers? A 12-month follow-up of a randomized controlled trial. *Psychol Med*. 2015 Jul;45(9):1907–17. <https://doi.org/10.1017/S0033291714003006>.
35. Ebert DD, Lehr D, Boß L, Riper H, Cuijpers P, Andersson G, et al. Efficacy of an internet-based problem-solving training for teachers: Results of a randomized controlled trial. *Scand J Work Env Health*. 2014;40(6):582–96. <https://doi.org/10.5271/sjweh.3449>.
36. Geraedts AS, Kleiboer AM, Wiezer NM, Cuijpers P, van Mechelen W, Anema JR. Feasibility of a worker-directed web-based intervention for employees with depressive symptoms. *internet Interv*. 2014 Jul;1(3):132–40. <https://doi.org/10.1016/j.invent.2014.07.001>.
37. Geraedts AS, Kleiboer AM, Twisk J, Wiezer NM, van Mechelen W, Cuijpers P. Long-term results of a web-based guided self-help intervention for employees with depressive symptoms: randomized controlled trial. *J Med internet Res*. 2014 Jan;16(7):e168. <https://doi.org/10.2196/jmir.3539>.
38. Geraedts AS, Kleiboer AM, Wiezer NM, van Mechelen W, Cuijpers P. Short-term effects of a web-based guided self-help intervention for employees with depressive symptoms: randomized controlled trial. *J Med internet Res*. 2014 Jan;16(5):e121. <https://doi.org/10.2196/jmir.3185>.
39. Baxter S, Sanderson K, Venn AJ, Blizzard CL, Palmer AJ. The Relationship Between Return on Investment and Quality of Study Methodology in Workplace Health Promotion Programs. *Am J Heal Promot*. 2014 Jun;28(6):347–63. <https://doi.org/10.4278/ajhp.130731-LIT-395>.
40. van Dongen JM, van Wier MF, Tompa E, Bongers PM, van der Beek AJ, van Tulder MW, et al. Trial-Based Economic Evaluations in Occupational Health: Principles, Methods, and Recommendations. *J Occup Environ Med*. 2014;56(6):563–72. <https://doi.org/10.1097/JOM.0000000000000165>.
41. Drummond MF, Sculpher MJ, Torrence GW, O'Brien BJ, Stoddart GL. *Methods for the Economic Evaluation of health Care Programmes*: third ed. Oxford: Oxford University Press; 2005.
42. Husereau D, Drummond M, Petrou S, Carswell C, Moher D, Greenberg D, et al. Consolidated health economic evaluation reporting standards (CHEERS) statement. *BMJ*. 2013;346. <https://doi.org/10.1136/bmj.f1049>.

43. Cost-Effectiveness Analysis Alongside Clinical Trials II—An ISPOR Good Research Practices Task Force Report. *Value Heal.* 2015;18(2):161–72. <https://doi.org/10.1016/j.jval.2015.02.001>.
44. Lesage F-X, Berjot S, Deschamps F. Psychometric properties of the French versions of the Perceived Stress Scale. *Int J Occup Med Environ Health.* 2012;25(2):178–84. <https://doi.org/10.2478/s13382-012-0024-8>.
45. Lazarus RS, Folkman S. *Stress, appraisal, and coping* [Sutoresuno shinrigaku]. New York, NY: Springer; 1984.
46. Ebert DD, Lehr D, Smit F, Zarski A-C, Riper H, Heber E, et al. Efficacy and cost-effectiveness of minimal guided and unguided internet-based mobile supported stress-management in employees with occupational stress: A three-armed randomised controlled trial. *BMC Public Health.* 2014;14(1). <https://doi.org/10.1186/1471-2458-14-807>.
47. Malouff JM, Thorsteinsson EB, Schutte NS. The efficacy of problem solving therapy in reducing mental and physical health problems: a meta-analysis. *Clin Psychol Rev.* 2007 Jan;27(1):46–57. <https://doi.org/10.1016/j.cpr.2005.12.005>.
48. Berking M, Ebert DD, Cuijpers P, Hofmann SG. Emotion regulation skills training enhances the efficacy of inpatient cognitive behavioral therapy for major depressive disorder: a randomized controlled trial. *Psychother Psychosom.* 2013 Jan;82(4):234–45. <https://doi.org/10.1159/000348448>.
49. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Heal Soc Behav.* 1983;24. <https://doi.org/10.2307/2136404>.
50. Jacobson NS, Truax P. Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *J Consult Clin Psychol.* 1991 Feb;59(1):12–9. <https://doi.org/10.1037/0022-006X.59.1.12>.
51. Hakkaart-van Roijen L, Van Straten A, Donker M, Tiemens B. Trimbos/iMTA questionnaire for costs associated with psychiatric illness (TIC-P). Rotterdam Inst Med Technol Assess. 2002.
52. Buntrock C, Berking M, Smit F, Lehr D, Nobis S, Riper H, et al. Preventing Depression in Adults With Subthreshold Depression: Health-Economic Evaluation Alongside a Pragmatic Randomized Controlled Trial of a Web-Based Intervention. *J Med internet Res.* 2017 Jan;19(1):e5. <https://doi.org/10.2196/jmir.6587>.
53. Bouwmans C, De Jong K, Timman R, Zijlstra-Vlasveld M, Van der Feltz-Cornelis C, Tan Swan S, et al. Feasibility, reliability and validity of a questionnaire on healthcare consumption and productivity loss in patients with a psychiatric disorder (TiC-P). *BMC Health Serv Res.* 2013 Jun 15;13:217. <https://doi.org/10.1186/1472-6963-13-217>.
54. Matthews JN, Altman DG, Campbell MJ, Royston P. Analysis of serial measurements in medical research. *BMJ.* 1990;300(6719):230–5. <https://doi.org/10.1136/bmj.300.6719.230>.
55. Destatis. Consumer Price Indices. [internet]. 2015. Available from: <https://www.destatis.de/EN/FactsFigures/NationalEconomyEnvironment/>.
56. Osterhaus JT, Gutterman DL, Plachetka JR. Healthcare resource and lost labour costs of migraine headache in the US. *Pharmacoeconomics.* 1992;2(1):67–76. <https://doi.org/10.2165/00019053-199202010-00008>.
57. van Hout BA, Al MJ, Gordon GS, Rutten FF. Costs, effects and C/E-ratios alongside a clinical trial. *Health Econ.* 3(5):309–19. <https://doi.org/10.1002/hec.4730030505>.
58. Briggs AH, Gray AM. Handling uncertainty in economic evaluations of healthcare interventions. *BMJ.* 1999 Sep;319(7210):635–8. <https://doi.org/10.1136/bmj.319.7210.635>.
59. Briggs A. Economic evaluation and clinical trials: size matters. *BMJ.* 2000 Dec;321(7273):1362–3. <https://doi.org/10.1136/bmj.321.7273.1362>.
60. Van Dongen JM, Proper KI, Van Wier MF, Van der Beek AJ, Bongers PM, Van Mechelen W, et al. Systematic review on the financial return of worksite health promotion programmes aimed at improving nutrition and/or increasing physical activity. *Obes Rev.* 2011;12(12):1031–49. <https://doi.org/10.1111/j.1467-789X.2011.00925.x>.
61. Dr. Henry AG, Dr. Jalpa AD, Dr. Seema SS, Dr. Daniel P. Designing economic evaluations in clinical trials [internet]. Oxford, UK: “Oxford University Press”; Available from: <file://oxfordmedicine.com/10.1093/med/9780199685028.001.0001/med-9780199685028-chapter-2>
62. Leigh JP. Expanding research on the economics of occupational health. *Scand J Work Environ Health.* 2006;32(1):1–4. <https://doi.org/10.5271/sjweh.969>.
63. Fenwick E, O’Brien B, Briggs A. Cost-effectiveness acceptability curves: facts, fallacies and frequently asked questions. *Health Econ.* 2004;13. <https://doi.org/10.1002/hec.903>.
64. Thiart H, Ebert DD, Lehr D, Nobis S, Buntrock C, Berking M, et al. internet-Based Cognitive Behavioral Therapy for Insomnia: A Health Economic Evaluation. *Sleep.* 2016;39(10):1769–78. <https://doi.org/10.5665/sleep.6152>.
65. Lo Sasso AT, Rost K, Beck A. Modeling the Impact of Enhanced Depression Treatment on Workplace Functioning and Costs. *Med Care.* 2006 Apr;44(4):352–8. <https://doi.org/10.1097/01.mlr.0000204049.30620.1e>.
66. Clarke MJ. Individual patient data meta-analyses. *Best Pract Res Clin Obstet Gynaecol.* 2005 Feb;19(1):47–55. <https://doi.org/10.1016/j.bpobgyn.2004.10.011>.
67. Jones AP, Riley RD, Williamson PR, Whitehead A. Meta-analysis of individual patient data versus aggregate data from longitudinal clinical trials. *Clin Trials.* 2009 Feb;6(1):16–27. <https://doi.org/10.1177/1740774508100984>.
68. Riley RD, Lambert PC, Abo-Zaid G. Meta-analysis of individual participant data: rationale, conduct, and reporting. *BMJ.* 2010 Jan;340:c221. <https://doi.org/10.1136/bmj.c221>.
69. Johns G, Miraglia M. The reliability, validity, and accuracy of self-reported absenteeism from work: A meta-analysis. *J Occup Health Psychol.* 2015 Jan;20(1):1–14. <https://doi.org/10.1037/a0037754>.

70. Prasad M, Wahlqvist P, Shikhar R, Shih Y-CT. A Review of Self-Report Instruments Measuring Health-Related Work Productivity. *Pharmacoeconomics*. 2004;22(4):225–44. <https://doi.org/10.2165/00019053-200422040-00002>.
71. The Use and Performance of Productivity Scales to Evaluate Presenteeism in Mood Disorders. *Value Heal*. 2012 Dec;15(8):1148–61. <https://doi.org/10.1016/j.jval.2012.08.2206>.
72. Lensberg BR, Drummond MF, Danchenko N, Despiéglé N, François C. Challenges in measuring and valuing productivity costs, and their relevance in mood disorders. *Clinicoecon Outcomes Res*. 2013;5:565–73. <https://doi.org/10.2147/CEOR.S44866>.
73. Furukawa TA, Noma H, Caldwell DM, Honyashiki M, Shinohara K, Imai H, et al. Waiting list may be a placebo condition in psychotherapy trials: a contribution from network meta-analysis. *Acta Psychiatr Scand*. 2014 Sep;130(3):181–92. <https://doi.org/10.1111/acps.12275>.

Received for publication: 1 August 2017