

Local and landscape responses of biodiversity in calcareous grasslands

Loos, Jacqueline; Krauss, Jochen; Lyons, Ashley; Föst, Stephanie; Ohlendorf, Constanze; Racky, Severin; Röder, Marina; Hudel, Lennart; Herfert, Volker; Tscharntke, Teja

Published in: **Biodiversity and Conservation**

DOI: 10.1007/s10531-021-02201-y

Publication date: 2021

Document Version Publisher's PDF, also known as Version of record

Link to publication

Citation for pulished version (APA):

Loos, J., Krauss, J., Lyons, A., Föst, S., Ohlendorf, C., Racky, S., Röder, M., Hudel, L., Herfert, V., & Tscharntke, T. (2021). Local and landscape responses of biodiversity in calcareous grasslands. *Biodiversity and* Conservation, 30(8-9), 2415-2432. https://doi.org/10.1007/s10531-021-02201-y

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ORIGINAL PAPER



Local and landscape responses of biodiversity in calcareous grasslands

Jacqueline Loos^{1,2} · Jochen Krauss³ · Ashley Lyons^{4,5} · Stephanie Föst¹ · Constanze Ohlendorf¹ · Severin Racky¹ · Marina Röder¹ · Lennart Hudel¹ · Volker Herfert¹ · Teja Tscharntke¹

Received: 6 January 2021 / Revised: 10 May 2021 / Accepted: 17 May 2021 / Published online: 26 May 2021 © The Author(s) 2021

Abstract

Across Europe, calcareous grasslands become increasingly fragmented and their quality deteriorates through abandonment and land use intensification, both affecting biodiversity. Here, we investigated local and landscape effects on diversity patterns of several taxonomic groups in a landscape of highly fragmented calcareous grassland remnants. We surveyed 31 grassland fragments near Göttingen, Germany, in spring and summer 2017 for vascular plants, butterflies and birds, with sampling effort adapted to fragment area. Through regression modelling, we tested relationships between species richness and fragment size (from 314 to 51,395 m²), successional stage, habitat connectivity and the per cent cover of arable land in the landscape at several radii. We detected 283 plant species, 53 butterfly species and 70 bird species. Of these, 59 plant species, 19 butterfly species and 9 bird species were grassland specialists. Larger fragments supported twice the species richness of plants than small ones, and hosted more species of butterflies, but not of birds. Larger grassland fragments contained more grassland specialist plants, but not butterfly or bird specialists. Increasing amounts of arable land in the landscape from 20 to 90% was related to the loss of a third of species of plants, and less so, of butterflies, but not of birds. Per cent cover of arable land negatively correlated to richness of grassland specialist plants and butterflies, but positively to grassland specialist birds. We found no effect by successional stages and habitat connectivity. Our multi-taxa approach highlights the need for conservation management at the local scale, complemented by measures at the landscape scale.

Keywords Abandonment \cdot Birds \cdot Butterflies \cdot Land use intensification \cdot Nature conservation \cdot Vascular plants

Jacqueline Loos jacqueline.loos@leuphana.de

Communicated by Dirk Sven Schmeller.

Extended author information available on the last page of the article

Introduction

During the twentieth century, land use changes towards agricultural intensification or abandonment decreased biodiversity and fragmented and degraded semi-natural biotopes, such as grasslands, across Europe (Bauer and Albrecht 2020, Poschlod and Wallis de Vries 2002). Semi-natural grasslands comprise sown and non-sown rangelands that developed anthropogenically through extensive forms of livestock herding and/ or mowing (Poschlod and Bonn 1998, Wallis de Vries et al. 2002, Willems 2001). Grazing as well as clearing of shrubs and tree saplings on semi-natural grasslands is a fundamental disturbance that prevents succession to forests (Hansson and Fogelfors 2000), which is the climax ecosystem in many central European landscapes. However, decreases in economic viability of sheep, goat and cattle grazing for wool, milk and meat reduced the number of sheep kept in Germany from 30 million in 1860 to 1.57 million by 2018 (Eurostat 2019). The subsequent abandonment of land, in addition to land use changes and intensification diminished plant species richness (Jacquemyn et al. 2011) with negative cascading effects on other grassland species over the long term (Balmer and Erhardt 2000; Gossner et al. 2016; Haddad et al. 2009; Polus et al. 2007). Nowadays, the conservation of semi-natural grasslands through management or avoidance of intensification is of high priority in Europe (Kahmen et al. 2002; Poschlod and WallisDeVries 2002; Valkó et al. 2016).

Due to their extraordinary plant species richness with up to 40 species per m² (Ellenberg and Leuschner 2010) and their vast decline, calcareous grasslands are of particular conservation concern in Europe. They are listed in Annex I of the Natura 2000 Habitats Directive (92/43/EEC) and are endangered in Germany (Finck et al. 2017). Dry grasslands on calcareous soils are considered the most species-rich grasslands in Central Europe (Diekmann et al. 2014; Karlik and Poschlod 2009; WallisDeVries et al. 2002), holding the world record for vascular plant species richness at small scales (Wilson et al. 2012). Due to their nutrient-poor conditions on shallow soils, calcareous grasslands provide optimal grounds for many rare and endangered species (Leuschner and Ellenberg 2017). However, these conditions also make these ecosystems and their distinct species composition vulnerable to the effects of environmental degradation, for example through climate change (Basto et al. 2018), atmospheric nutrient deposition (Bobbink et al. 2010) and by changes in land management.

Semi-natural grasslands in central Europe decreased substantially during the last century, e.g. in some regions of Germany by up to 95% (Brückmann et al. 2010). In a Europewide study, per region 18–80% of calcareous grassland area present in the middle of the twentieth century has been lost, with a range between 0 and 99.8% of area loss per patch (Krauss et al. 2010). Similarly, connectivity of grasslands in some regions decreased by up to 90% (Cousins et al. 2007). The most recent estimate of the extent of calcareous grassland in Natura 2000 habitats with high orchid richness in Germany is 31,079 ha, scattered into 924 sites (Calaciura and Spinelli 2008). The small size of many of these grassland fragments (with many less than one hectare) exposes them to edge effects, succession through abandonment and reduced connectivity, which reduces their genetic diversity and ecological fitness (Matthies et al. 2004). Therefore, conservation management plays a crucial role in maintaining these fragments both as core habitats and as stepping stones in agricultural landscapes (Madeira et al. 2016).

Additionally, long-lasting flow-on effects from the use of agrochemicals (Evans and Sanderson 2018; Melts et al. 2018) and from atmospheric nitrogen deposition (Dupre et al. 2010; Field et al. 2014) put grassland species under pressure (Fischer

and Stocklin 1997). Specialist species might be more affected than generalist species, as specialist species are in many cases mono- or oligophagous and therefore decline more dramatically than polyphagous species (Ernst et al. 2017; Steffan-Dewenter and Tscharntke 2000; Wenzel et al. 2006). Generalist species often have greater dispersal ability and larger landscape-wide population pools, which might increase success of survival in fragmented grassland landscapes (Riibak et al. 2015). Declines in species richness and changes in species composition towards biotic homogenisation (McKinney and Lockwood 1999) may affect the multiple ecosystem functions that grasslands provide (Allan et al. 2015; Diekmann et al. 2014; Huber et al. 2017). To this end, generalist species provide most functions, but specialist species can play an important role in complementing functions and maintaining the resilience of the ecosystem.

To understand the conservation value of these remaining grassland fragments, it is thus important to observe multiple species groups (Zulka et al. 2014) and account for the surrounding landscape structures at various spatial scales (Diacon-Bolli et al. 2012; Soderstrom et al. 2001). Such multileveled evidence on species richness and community composition contributes to informed management recommendations for this highly fragmented ecosystem. Here, we investigated how different taxonomic groups respond to size, habitat connectivity and landscape context of calcareous grassland fragments. For this purpose, we conducted a multi-taxa survey on plants, butterflies and birds in 31 calcareous grassland fragments near Göttingen, Lower Saxony in 2017. We selected these species groups because we expected them to react differently to environmental changes, and because they contain both specialist and generalist species that differ in abilities to disperse across the landscape (Dormann et al. 2007; Habel et al. 2016). We asked (i) which environmental variables influence local biodiversity patterns of plants, butterflies and birds?; and (ii) which spatial scale of landscape composition affects local biodiversity patterns?

Material and methods

Study site selection

We sampled 31 calcareous grassland fragments in the vicinity of Göttingen, Lower Saxony (Germany), which were subject to earlier studies (Steffan-Dewenter and Tscharntke 2000; Krauss et al. 2003a, 2004). These sites represent a gradient in size and habitat connectivity and provide a sub-sample of a total of 285 calcareous grassland fragments in the landscape. The nearest distance of the 31 fragments to other calcareous grassland fragments was 55 m, the furthest distance was 1894 m. Management on the sites varied from no visible management actions to grazing with cattle, goats or donkeys. Mowing regime varied from one mowing event per year up to mowing twice per year. The calcareous grasslands belonged to the plant association *Gentiano-Koelerietum* (Krauss et al. 2003a, 2004), but succession towards the natural climax state of a woody vegetation community had completely transformed one of the small grassland patches into a young forest, while only two grassland patches showed no sign of succession taking place. Those two intact grassland patches were small and medium in size. The other 28 grassland patches showed natural succession taking place at least to some extent (Fig. 1).



Fig. 1 Location of the study area within Europe (inset map) and display of the 31 study sites near Göttingen (center of map), lower Saxony, Germany. Map data © ESRI, accessed through QGIS 3

Biodiversity sampling

On all 31 study sites, we conducted surveys of plants, butterflies and birds during spring and summer of 2017. We used an area-adapted sampling strategy similar to previous surveys in the years 1996 and 2000 for vegetation and butterflies (Krauss et al. 2003a, 2004).

Plant surveys

We surveyed vascular plants on two occasions in May/June and in August in a total of 188 5 m×5 m quadrats. We adjusted the sampling effort to the size of the study site and thus placed one plot on small sites (314–1326 m²), two plots on medium sites (1914–7887 m²) and three plots on large sites (11,528–51,395 m²) in each sampling round. In each quadrat, we recorded per cent cover of each vascular plant species. Additionally, we surveyed the grassland fragments for further vascular plant species occurring outside the plots. In the field, species were identified with Jäger (2011) and Jäger et al. (2013). Nomenclature followed GermanSL Version 1.3 (Jansen and Dengler 2008). We classified grassland specialist plant species following von Drachenfels (2016).

Butterfly surveys

We surveyed butterflies and burnet moths along a total of 155 timed transects between May and August on five occasions with an interval of approximately two weeks. We

2419

adapted the survey effort to the size of the study sites (Krauss et al. 2003a, b; Steffan-Dewenter and Tscharntke 2000). Transect time per walk was 20 min for small grasslands, 40 min for intermediate grasslands and 60 min for large grasslands. We conducted the surveys under suitable conditions (no rain, <90% cloud cover, >17 °C if clouded, >13 °C if sunny, no strong wind (Pollard 1977)). Species identification and nomenclature followed Settele et al. (2015) for diurnal butterflies, and Zub (1996) and Weidemann and Köhler (1996) for burnets. Species that could not be distinguished with the help of external characteristics were counted as species-complex. We defined butterflies as grassland specialists when their known occurrence is linked to open, nutrient poor grassland habitats (Bink 1992) such as calcareous grasslands in Lower Saxony.

Bird surveys

We sampled birds through point counts (Bibby et al. 2000) between March and June 2017, following Südbeck and Weick (2005). We sampled each site on four different occasions throughout the season with a minimum interval of 10 days between the counts. During a 10-min survey sequence, we recorded all birds heard or seen within a fixed radius of 50 m around the centroid of the study site. Surveys began shortly after sunrise and ended at 10 am by the latest. We classified species as forest or open land species by using habitat categories of the Red List (Krüger and Nipkow 2015; Storchová and Hořák 2018).

Environmental variables

For subsequent analyses of biodiversity patterns, we used the following environmental variables as predictors: The current size of the fragments ("Fragment size") was calculated by a combined approach through observations in the field using a handheld GPS, which were then digitized and analyzed based on latest Google Earth satellite imagery in ArcGIS 10.4. Remnants were categorized into small (n = 11; 314-1326 m²), medium (n = 12; 1914-7887 m²) and large (n = 8; 11,528-51,395 m²) sites based on Krauss et al. (2003a, b). A habitat isolation index *I* had been calculated for the study sites in the year 2000 with the following formula derived from Hanski et al. (1994):

$$I = \sum e^{-dij} A_j$$

with A_j as the size of the nearest calcareous grassland patch, and dij as distance to the study site *i* (Steffan-Dewenter and Tscharntke 2000). The index *I* served as a negative proxy for habitat connectivity ("Connectivity"). We calculated the per cent cover of arable land cover ("% arable") at different spatial scales around the centroid of the focal grassland fragments in buffers of 250 m steps from 250 to 3000 m based on ATKIS Data (LGLN 2017). Moreover, we estimated the stage of succession on a categorical scale from 1 (no succession) to 4 (transformed into a different ecosystem—young forest).

Statistical analysis

We calculated biodiversity patterns of all investigated taxonomic groups as a response to the above-mentioned environmental variables. For this, we estimated species richness for all taxonomic groups using Chao 1 estimator of asymptotic richness (Chao 1984) (function "estimateR"). Estimated species richness strongly correlated with observed species richness (r=0.93 for plants, r=0.81 for butterflies and r=0.57 for birds). Estimated species richness served as the response variable in generalised linear models with Poisson error distribution in which the natural logarithm of the current size of grassland fragments, habitat connectivity and per cent cover of arable land at different spatial scales served as explanatory variables. We tested all environmental variables for collinearity in advance, with the highest correlation between fragment size and connectivity (Pearson's r: 0.29). We created individual models for each taxon at each spatial scale and compared model performance by Akaike's Information Criterion (AIC). We also calculated generalised linear models with the number of grassland species per taxon as a response variable, using the same environmental explanatory variables. Through multivariate statistics, we investigated species composition per size class of each taxon with NMDS. For each ordination, we superimposed all environmental variables and tested for significant (p < 0.05) correlations on species composition (function "envfit"). All analyses were conducted using the package "vegan" (Oksanen et al. 2018) in the program R (R Core Team 2016).

Results

General findings

In total, we found 283 plant species, 53 butterfly species and 70 bird species. Of these, we identified 59 plant species, 19 butterfly species and 9 bird species as grassland specialists.

Local and landscape effects on different species groups

Species richness models

Our models revealed that species richness of plants significantly increased with increasing size of the grassland fragment (Table 1, Fig. 2a), while an increasing per cent cover of arable land up to a radius of 1500 m around the grassland fragment had significantly decreased plant species richness (Table 1, Fig. 2b). Butterfly species richness significantly decreased with an increasing amount of arable land at small spatial scales up to 1000 m. None of our models explained the estimated richness of birds on calcareous grassland fragments at any spatial scale. Neither landscape connectivity nor the successional stages were significant for our models on any investigated species group at any scale.

Grassland specialist models

Fragment size was a positive predictor for species richness of grassland specialist plants at all spatial scales (Table 2). The increasing per cent cover of arable land showed a significant decrease of grassland specialist plants at scales from 500 to 2250 m and for birds from 1000 to 1500 m.

Species composition

Across all taxonomic groups, there was more variability in the composition of small fragments (Fig. 3). Large and small fragments differed in their species composition most

Table 1 Resu	ilts of the gei	reralized linea	ar mixed mode	els for the esti	mated species	s richness of th	rree different	taxonomic gr	oups at differe	ent spatial sca	lles	
	Radius (m)											
	250	500	750	1000	1250	1500	1750	2000	2250	2500	2750	3000
Plants												
AIC	279.64	272.21	271.03	274.88	277.51	279.31	281.10	282.40	283.01	283.28	283.66	284.09
Intercept	139.19^{***}	141.00^{***}	139.80^{***}	139.86^{***}	139.43***	137.53***	135.07***	132.83***	131.38^{***}	130.92^{***}	130.02^{***}	128.14^{***}
Fragment size	0.0005	0.0006*	0.0007*	0.0006*	0.0006*	0.0006*	0.0006*	0.0006*	0.0006*	0.0006*	0.0006	0.0006
Succession	- 7.96	- 2.66	- 2.25	- 4.74	- 6.10	- 6.64	- 7.22	- 7.90	- 8.19	- 8.37	- 8.77	- 9.37
Connectiv- ity	- 0.0001	- 0.0001	- 0.0001	- 0.0001	- 0.0002	- 0.0002	- 0.0003	- 0.0003	- 0.0003	- 0.0003	- 0.0003	- 0.0003
% arable	-0.45*	-0.74^{**}	-0.78^{***}	-0.68^{**}	-0.60*	-0.53*	- 0.44	- 0.35	-0.31	- 0.29	- 0.25	-0.18
Butterflies												
AIC	212.46	215.31	216.72	216.93	217.68	218.75	219.35	219.54	219.72	219.74	219.76	219.60
Intercept	45.43***	42.54***	41.58^{***}	42.11***	42.31***	41.76^{***}	41.42***	41.37^{***}	41.26^{***}	41.40^{***}	41.64^{***}	42.01***
Fragment size	0.00015	0.00022*	0.00023*	0.00022*	0.00022*	0.00022*	0.00023*	0.00023*	0.00023*	0.00023*	0.00024*	0.00024*
Succession	- 2.96	- 2.05	- 2.25	- 2.67	- 2.96	- 3.11	- 3.17	- 3.20	- 3.16	- 3.14	- 3.14	- 3.10
Connectiv- ity	0.00001	- 0.00001	- 0.00002	- 0.00003	- 0.00005	- 0.00006	- 0.00006	- 0.0007	- 0.00006	- 0.00006	- 0.00006	- 0.00005
% arable	- 0.23**	-0.21*	- 0.19*	-0.19*	- 0.18	- 0.16	- 0.15	- 0.14	- 0.14	- 0.15	- 0.15	- 0.17
Birds												
AIC	246.32	246.39	246.28	245.76	244.81	243.57	242.92	242.42	242.43	242.82	243.12	243.38
Intercept	23.22*	21.13*	20.74*	19.26	17.41	15.97	15.31	14.74	14.70	14.75	14.59	14.48
Fragment size	- 0.00007	- 0.00006	- 0.00007	- 0.00007	- 0.00007	- 0.00009	- 0.0001	- 0.0001	- 0.00011	- 0.00011	- 0.00011	- 0.00011
Succession	- 1.13	- 1.68	- 1.92	- 2.29	- 2.62	- 3.04	- 3.29	- 3.44	- 3.57	- 3.53	- 3.44	- 3.36

_
ģ
n
÷E
Ē
્ઇ
<u> </u>
ς
e
ā
ص .

 $\underline{\textcircled{O}}$ Springer

	(noniiiii											
	Radius (m)											
	250	500	750	1000	1250	1500	1750	2000	2250	2500	2750	3000
Connectiv- ity	0.00016	0.00013	0.00012	0.00011	0.0001	0.0001	0.00011	0.0001	0.0001	0.0000	0.00009	0.0000
% arable	- 0.04	0.03	0.05	0.11	0.16	0.22	0.24	0.26	0.27	0.27	0.27	0.27
Model nerfo	rmance is co	nnared hv Ak	taike's inform	ation criterio	n (AIC. renr	esented in ita	lic). with the l	hest nerform	ing model ind	licated in hold	italics letters	Cell contain

Model performance is compared by Akaike's information criterion (AIC, represented in italic), with the best performing model indicestimates of the explanatory variable. Significances of the explanatory variables are displayed by ***P < 0.001; **P < 0.01; *P < 0.05

prominently for plants and butterflies, while large and small sites shared species composition for birds (Fig. 3). The community composition of plants and butterflies was significantly related to the percent cover of arable land at 250 m radius around the grassland fragment and to the successional stage of the grassland fragment. The amount of arable land in the landscape correlated with plant species composition also at a radius of 1750 m. Patch size emerged as predictor for species composition of birds, but was only marginally significant for butterflies (Table 3). Neither habitat connectivity nor the per cent of arable land at a 3000 m radius around the grassland fragments in the landscape influenced species composition.

Discussion

Our study shows the diverging importance of several environmental variables for diversity in three taxonomic groups both at the local and at the landscape scale. At the local scale, the size of the calcareous grassland fragment predicted species richness of plants and butterflies, both for all species and for specialist species. In contrast, species richness of various taxonomic groups was independent of current habitat area in other studies (Deák et al. 2016; Huber et al. 2017; Lengyel et al. 2016), which is counterintuitive to the often-confirmed expectation that larger habitat area increases species richness (Adriaens et al. 2006; Cousins et al. 2007; Krauss et al. 2004). The per cent cover of arable land was far more important for species richness and species composition of plants and butterflies than connectivity to other dry grassland patches. These findings emphasize the hostility of landscapes that are dominated by intensive and homogenous cropland for species richness (Hallmann et al. 2017; Kormann et al. 2015). However, this effect was not pronounced for birds, which might be less sensitive to landscape connectivity and might be more tied to the presence of woody vegetation because of their higher mobility (Duflot et al. 2018; Dorresteijn et al. 2018).

Our results also concur with previous studies that documented the relatively low importance of connectivity (Adriaens et al. 2006; Cousins et al. 2007; Huber et al. 2017), despite its often-demonstrated importance for the immigration of new species and individuals (Rösch et al. 2015). Thus, it remains a challenge to derive general assumptions on the role of habitat connectivity (Ibáñez et al. 2014), as different taxonomic groups may respond differently to variables on the landscape scale (Lengyel et al. 2016). Considerations of both habitat area and connectivity, as well as landscape composition and configuration, can be important for explaining species diversity patterns (Lindgren and Cousins 2017). The per cent cover of arable land negatively affected both the species richness of all plants and butterflies as well as grassland specialist plants and butterflies. Moreover, the per cent cover of arable land influenced the species composition of all investigated taxonomic groups (supporting Kormann et al. 2015). Biodiversity management, thus, would ideally target both the local and the landscape scale (Loos et al. 2019). As an example, the introduction of buffer zones might prove beneficial for invertebrates (Madeira et al. 2016). Moreover, a closer examination of management details over long periods might provide more insights into the effect of quality of the calcareous grassland fragments on biodiversity.

Our study also observed one complete loss and decrease in size of some fragments. This observation is alarming as it highlights general trend of habitat loss in agricultural land-scapes despite formal protection under Annex I of the EU habitats directive across EU member states (Calaciura and Spinelli 2008). The observed losses stem from abandonment



Fig. 2 Display of species richness for plants, butterflies, birds and **a** fragment size (natural logarithm); **b** per cent cover of arable land at 1000 m radius. Color code: plants = green, butterflies = red, birds = blue

and consequent succession on the one hand and transformation of land use, especially to margins of roads or to agricultural fields, on the other hand. At first, these small losses may appear insignificant, but in sum they constitute habitat loss at a larger scale. Such gradual changes are difficult to track and control, as the study area hosts almost 300 single fragments, of which approximately 70% are smaller than 1 ha (Rösch et al. 2015). The small size of these fragments and their isolated position in the landscape make it difficult and costly to maintain them (Gustavsson et al. 2007). No-action, however, poses a particular problem to calcareous grasslands, as they are naturally subject to succession. To halt this trend requires a lot of effort, particularly with the limited resources that are available at the implementing agency for nature conservation. Thus, prioritization of conservation efforts is necessary.

Whether it is more useful for conservation success to concentrate and improve the conservation value on few large sites or on many small sites is an old debate (e.g. Diamond 1975), which the responsible government authority alone cannot solve. From an ecological perspective, it has been shown that both small and large fragments are important for biodiversity at the landscape scale (Rösch et al. 2015; Tscharntke et al. 2002), as they play an important function as stepping stones (Lindborg et al. 2014; Wenzel et al. 2006) and create higher connectivity, which is a key factor for many ecological functions (Brückmann et al. 2010; Lindborg et al. 2014). Moreover, diverse landscapes are able to host higher species diversity (Mallinger et al. 2016). Thus, both large and small fragments are important to create a network of grassland patches (Deák et al. 2016; Rösch et al. 2015; Zulka et al. 2014). Nevertheless, maintenance of many small sites is linked to the practical challenge of affordable management regimes in the respective sites, which is key for their conservation (Huber et al. 2017).

For the conservation management of small grassland fragments, it might be useful to foster systematic collaborations with amateur animal keepers that use the grassland fragments for grazing of their horses (Köhler et al. 2016), donkeys, or small flocks of sheep, which through a rotational system might even promote seed dispersal between sites (Riibak et al. 2015). However, the type and stocking density of animals (Lyons et al. 2018, 2017), as well as the timing and duration of grazing affects calcareous grasslands differently (Goodenough and Sharp 2016; Römermann et al. 2009) and should thus be agreed upon and controlled. A viable alternative to grazing is mowing (Kahmen et al. 2002), even though mowing allows fewer seeds to be dispersed (Poschlod and Bonn 1998). Periodic

Table 2 Resu Scales	alts of the g	eneralised line	ear mixed mo	dels for the	species richne	ess of grassla	nd specialist	species of the	three differe	ent taxonomic	e groups at di	fferent spatial
	Radius (m)											
	250	500	750	1000	1250	1500	1750	2000	2250	2500	2750	3000
Grassland												
AIC	215.79	209.59	205.39	205.21	206.13	207.58	209.91	213.29	215.29	216.13	217.21	218.19
Intercept	41.15^{***}	41.59^{***}	41.74***	42.72***	43.38^{***}	43.14^{***}	42.60^{***}	41.65***	40.85***	40.63^{***}	40.27***	39.73***
Fragment size	0.00026*	0.00032**	0.00034***	• 0.00032**	*0.00032***	0.00033**	* 0.00034**	0.00034**	0.00034**	0.00034**	0.00033**	0.00033**
Succession	-2.01	-0.25	0.18	-0.40	-0.75	-0.82	-0.90	-1.19	-1.34	- 1.43	-1.63	-1.84
Connectiv- ity	0	0.00002	0.00002	0.00001	0	-0.00002	-0.00003	- 0.00004	-0.00004	- 0.00004	-0.00003	- 0.00004
% arable	-0.16	-0.25^{**}	-0.29^{***}	-0.29^{***}	-0.29^{***}	-0.28^{**}	-0.26^{**}	-0.22*	-0.20*	-0.19	-0.18	-0.15
Grassland bu	tterflies											
AIC	143.71	142.78	142.56	141.74	141.38	141.6	142.49	143.96	144.61	144.72	145	145.27
Intercept	6.58^{**}	6.49**	6.43**	6.69**	6.89^{***}	6.87^{***}	6.72^{**}	6.41^{**}	6.19^{**}	6.17^{**}	6.06^{**}	5.87**
Fragment size	0	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001
Succession	-0.8229	-0.5741	-0.5519	-0.5902	-0.6199	-0.6256	-0.6549	-0.7506	-0.8021	-0.8106	-0.8537	-0.9143
Connectiv- ity	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001
% arable	-0.03	- 0.04	- 0.04	-0.05	-0.05	- 0.05	- 0.04	-0.03	-0.02	-0.02	-0.02	-0.01
Grassland bir	sp											
AIC	87.86	86.7	85.66	83.93	83.87	84.52	85.01	85.56	86.34	87.17	87.76	88.11
Intercept	0.86	0.88	0.86	0.71	0.64	0.67	0.69	0.71	0.76	0.81	0.85	0.87
Fragment size	0.00002	0.00002	0.00002	0.00002	0.00002	0.00002	0.00001	0.00001	0.00001	0.00001	0.00001	0.00002
Succession	-0.35	-0.46	-0.49	-0.48	- 0.46	-0.45	-0.45	- 0.44	- 0.44	-0.42	-0.40	-0.39

2425

(continued)	
Table 2	

 $\underline{\textcircled{O}}$ Springer

	Radius (m)											
	250	500	750	1000	1250	1500	1750	2000	2250	2500	2750	3000
Connectiv- (ity	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001
% arable	0.01	0.02	0.02	0.02^{*}	0.02*	0.02*	0.02	0.02	0.02	0.02	0.02	0.01
Model perforn estimates of th	mance is cc 1e explanat	mpared by Al ory variable. S	kaike's inform Significances c	nation criterio of the explane	on (AIC, repr atory variable	esented in itali s are displayed	ics), with the 1 by ***P<0	best perform .001; **P < 0	ing model inc .01; *P<0.05	licated in Bo	ld italics lette	rs. Cell contain



Fig.3 Non-metric multi-dimensional scaling (NMDS) ordinations of species compositions in the three different taxonomic groups. Displayed are the first two axes of the ordination, with NMDS Axis 1 at the x axis and NMDS Axis 2 at the y axis, respectively. Blue dots indicate sites, Ordihull areas indicate size classes: grey=small, green=medium, pink=large

	Plants	Butterflies	Birds
Patch size	0.113	0.069	0.039*
Connectivity	0.474	0.222	0.718
% Arable land 250 m	0.005**	0.022*	0.131
% Arable land 1750 m	0.032*	0.377	0.520
% Arable land 3000 m	0.311	0.987	0.930
Successional stage	0.006**	0.017*	0.932

Displayed are P-values of environmental variables from an environmental Fit function onto an ordination space of species composition for each respective taxon. ***<0.001; **<0.01; *<0.05;<0.1; *n.s.* not significant

mowing, for example once in three years, seems helpful for the maintenance of species richness, as this intervention is able to avert the process of natural succession and is able to prevent vegetation change (Hansson and Fogelfors 2000). In general, a more diversified grassland management towards less intensity is favorable for semi-natural elements in the landscape (Diekmann et al. 2014; Gustavsson et al. 2007). Given the challenging conditions to translate diversification into practice, stewardship approaches and increased involvement of local people might pose a viable alternative to the bureaucratic formal options that agri-environmental schemes have offered so far.

Conclusion

Table 3 Influence ofenvironmental variables onspecies composition

Calcareous grasslands are facing serious conservation concerns, and the status of these habitats in Germany depends on many small-scale interventions to maintain their connectivity and quality. In particular, small grassland fragments (<1.5 ha) are prone to shrub encroachment due to abandonment. Both local and landscape variables showed some impact on species richness and composition of some taxonomic groups, in particularly on habitat specialists. Given the difficulties to manage the many scattered fragments and the

diverse conservation targets within the landscapes, novel and less formal approaches of maintenance of calcareous grassland might prove beneficial for conservation in the future.

Acknowledgements We are grateful towards Bertram Preuschhoff and Sebastian Barthold of the nature conservation agencies Göttingen and Northeim for information and permits; towards the land managers of the calcareous grassland fragments for their support and towards Ingolf Steffan-Dewenter and Erwin Bergmeier for discussions. JL was partly funded through a Robert-Bosch Junior Professorship for Research into the Sustainable Use of Natural Resources; AL was funded through a Stapledon Travel Fellowship. We thank two anonymous reviewers for their constructive feedback on an earlier version of the manuscript.

Funding Open Access funding enabled and organized by Projekt DEAL.

Data availability Data and code can be requested directly from the authors.

Declarations

Conflict of interest We declare no conflict of interests.

Ethical approval The study was permitted by the administrative district environmental councils of Northeim and Göttingen in April 2017.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Adriaens D, Honnay O, Hermy M (2006) No evidence of a plant extinction debt in highly fragmented calcareous grasslands in Belgium. Biol Conserv 133:212–224
- Allan E, Manning P, Alt F, Binkenstein J, Blaser S, Bluethgen N, Bohm S, Grassein F, Holzel N, Klaus VH, Kleinebecker T, Morris EK, Oelmann Y, Prati D, Renner SC, Rillig MC, Schaefer M, Schloter M, Schmitt B, Schoning I, Schrumpf M, Solly E, Sorkau E, Steckel J, Steffen-Dewenter I, Stempfhuber B, Tschapka M, Weiner CN, Weisser WW, Werner M, Westphal C, Wilcke W, Fischer M (2015) Land use intensification alters ecosystem multifunctionality via loss of biodiversity and changes to functional composition. Ecol Lett 18:834–843
- Balmer O, Erhardt A (2000) Consequences of succession on extensively grazed grasslands for central European butterfly communities: rethinking conservation practices. Conserv Biol 14:746–757
- Basto S, Thompson K, Grime JP, Fridley JD, Calhim S, Askew AP, Rees M (2018) Severe effects of longterm drought on calcareous grassland seed banks. NPJ Clim Atmosph Sci 1:1–7
- Bauer M, Albrecht H (2020) Vegetation monitoring in a 100-year-old calcareous grassland reserve in Germany. Basic Appl Ecol 42:15–26
- Bibby CJ, Burgess ND, Hill DA, Mustoe S (2000) Bird census techniques. Elsevier, London
- Bink FA (1992) Ecologische Atlas van de dagvlinders van Noordwest-Europa. Schuyt, Haarlem.
- Brückmann SV, Krauss J, Steffan-Dewenter I (2010) Butterfly and plant specialists suffer from reduced connectivity in fragmented landscapes. J Appl Ecol 47:799–809
- Calaciura B, Spinelli O (2008) Management of Natura 2000 habitats. 6210 Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia)(* important orchid sites). European Commission.

Chao A (1984) Nonparametric estimation of the number of classes in a population. Scand J Stat 11:265-270

- Deák B, Valkó O, Török P, Tóthmérész B (2016) Factors threatening grassland specialist plants-A multiproxy study on the vegetation of isolated grasslands. Biol Conserv 204:255–262
- Diacon-Bolli J, Dalang T, Holderegger R, Bürgi M (2012) Heterogeneity fosters biodiversity: linking history and ecology of dry calcareous grasslands. Basic Appl Ecol 13:641–653
- Diamond JM (1975) The island dilemma: Lessons of modern biogeographic studies for the design of natural reserves. Biol Conserv 7:129–145
- Diekmann M, Jandt U, Alard D, Bleeker A, Corcket E, Gowing DJG, Stevens CJ, Dupre C (2014) Longterm changes in calcareous grassland vegetation in North-western Germany—no decline in species richness, but a shift in species composition. Biol Conserv 172:170–179
- Dormann CF, Schweiger O, Augenstein I, Bailey D, Billeter R, de Blust G, DeFilippi R, Frenzel M, Hendrickx F, Herzog F, Klotz S, Liira J, Maelfait JP, Schmidt T, Speelmans M, van Wingerden WKRE, Zobel M (2007) Effects of landscape structure and land-use intensity on similarity of plant and animal communities. Glob Ecol Biogeogr 16:774–787
- Dorresteijn I, Nimmo DG, Loos J, Hanspach J, Moga CI, David A, Ernst LM, Fischer J (2018) A new world for old landscapes: Land-use intensification and bird conservation in a traditional farming landscape. N West J Zool 14:199–207
- Duflot R, Daniel H, Aviron S, Alignier A, Beaujouan V, Burel F, Cochard A, Ernoult A, Pain G, Pithon JA (2018) Adjacent woodlands rather than habitat connectivity influence grassland plant, carabid and bird assemblages in farmland landscapes. Biodivers Conserv 27:1925–1942
- Dupre C, Stevens CJ, Ranke T, Bleeker A, Peppler-Lisbach C, Gowing DJG, Dise NB, Dorland E, Bobbink R, Diekmann M (2010) Changes in species richness and composition in European acidic grasslands over the past 70 years: the contribution of cumulative atmospheric nitrogen deposition. Glob Change Biol 16:344–357
- Ellenberg H, Leuschner C (2010) Vegetation Mitteleuropas mit den Alpen: in ökologischer, dynamischer und historischer Sicht. UTB, Stuttgart
- Ernst LM, Tscharntke T, Batáry P (2017) Grassland management in agricultural vs. forested landscapes drives butterfly and bird diversity. Biol Conserv 216:51–59
- Eurostat (2019) Number of sheep. European Commission. https://ec.europa.eu/eurostat/web/products-datas ets/product?code=tag00017. Accessed 29 Sept 2020
- Evans EC, Sanderson RA (2018) Long-term fertilizer regimes have both direct and indirect effects on arthropod community composition and feeding guilds. J Appl Entomol 142:230–240
- Field CD, Dise NB, Payne RJ, Britton AJ, Emmett BA, Helliwell RC, Hughes S, Jones L, Lees S, Leake JR, Leith ID, Phoenix GK, Power SA, Sheppard LJ, Southon GE, Stevens CJ, Caporn SJM (2014) The role of nitrogen deposition in widespread plant community change across semi-natural habitats. Ecosystems 17:864–877
- Finck P, Heinze S, Raths U, Riecken U, Ssymank A (2017) Rote Liste der gefährdeten Biotoptypen Deutschlands: dritte fortgeschriebene Fassung 2017. Bundesamt für Naturschutz, Bonn.
- Fischer M, Stocklin J (1997) Local extinctions of plants in remnants of extensively used calcareous grasslands 1950–1985. Conserv Biol 11:727–737
- Goodenough AE, Sharp MH (2016) Managing calcareous grassland for the declining Duke of Burgundy Hamearis lucina butterfly: effects of grazing management on Primula host plants. J Insect Conserv 20:1087–1098
- Gossner MM, Lewinsohn TM, Kahl T, Grassein F, Boch S, Prati D, Birkhofer K, Renner SC, Sikorski J, Wubet T (2016) Land-use intensification causes multitrophic homogenization of grassland communities. Nature 540:266–269
- Gustavsson E, Lennartsson T, Emanuelsson M (2007) Land use more than 200 years ago explains current grassland plant diversity in a Swedish agricultural landscape. Biol Conserv 138:47–59
- Habel JC, Segerer A, Ulrich W, Torchyk O, Weisser WW, Schmitt T (2016) Butterfly community shifts over two centuries. Conserv Biol 30:754–762
- Haddad NM, Crutsinger GM, Gross K, Haarstad J, Knops JM, Tilman D (2009) Plant species loss decreases arthropod diversity and shifts trophic structure. Ecol Lett 12:1029–1039
- Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, Stenmans W, Müller A, Sumser H, Hörren T (2017) More than 75 percent decline over 27 years in total flying insect biomass in protected areas. PLoS ONE 12:e0185809
- Hanski I, Kuussaari M, Nieminen M (1994) Metapopulation structure and migration in the butterfly Melitaea cinxia. Ecology 75:747–762
- Hansson M, Fogelfors H (2000) Management of a semi-natural grassland; results from a 15-year-old experiment in southern Sweden. J Veg Sci 11:31–38

- Huber S, Huber B, Stahl S, Schmid C, Reisch C (2017) Species diversity of remnant calcareous grasslands in south eastern Germany depends on litter cover and landscape structure. Acta Oecolog 83:48–55
- Ibáñez I, Katz DS, Peltier D, Wolf SM, Barrie C, Benjamin T (2014) Assessing the integrated effects of landscape fragmentation on plants and plant communities: the challenge of multiprocess–multiresponse dynamics. J Ecol 102:882–895
- Jacquemyn H, van Mechelen C, Brys R, Honnay O (2011) Management effects on the vegetation and soil seed bank of calcareous grasslands: an 11-year experiment. Biol Conserv 144:416–422
- Jäger EJ (2011) Rothmaler. Exkursionsflora von Deutschland: Gefäßpflanzen: Grundband. Spektrum Akademischer Verlag, Heidelberg
- Jäger EJ, Müller F, Ritz CM, Welk E, Wesche K (2013) Rothmaler. Exkursionsflora von Deutschland: Gefäßpflanzen: Atlasband. Springer-Verlag, Berlin Heidelberg
- Jansen F, Dengler J (2008) GermanSL Eine universelle taxonomische Referenzliste f
 ür Vegetationsdatenbanken in Deutschland. Tuexenia 28:239–253
- Kahmen S, Poschlod P, Schreiber KF (2002) Conservation management of calcareous grasslands. Changes in plant species composition and response of functional traits during 25 years. Biol Conserv 104:319–328
- Karlik P, Poschlod P (2009) History or abiotic filter: which is more important in determining the species composition of calcareous grasslands? Preslia 81:321–340
- Köhler M, Hiller G, Tischew S (2016) Year-round horse grazing supports typical vascular plant species, orchids and rare bird communities in a dry calcareous grassland. Agric Ecosyst Environ 234:48–57
- Kormann U, Rösch V, Batáry P, Tscharntke T, Orci KM, Samu F, Scherber C (2015) Local and landscape management drive trait-mediated biodiversity of nine taxa on small grassland fragments. Divers Distrib 21:1204–1217
- Krauss J, Bommarco R, Guardiola M, Heikkinen RK, Helm A, Kuussaari M, Lindborg R, Öckinger E, Pärtel M, Pino J (2010) Habitat fragmentation causes immediate and time-delayed biodiversity loss at different trophic levels. Ecol Lett 13:597–605
- Krauss J, Klein AM, Steffan-Dewenter I, Tscharntke T (2004) Effects of habitat area, isolation, and landscape diversity on plant species richness of calcareous grasslands. Biodivers Conserv 13:1427–1439
- Krauss J, Steffan-Dewenter I, Tscharntke T (2003a) How does landscape context contribute to effects of habitat fragmentation on diversity and population density of butterflies? J Biogeogr 30:889–900
- Krauss J, Steffan-Dewenter I, Tscharntke T (2003b) Local species immigration, extinction, and turnover of butterflies in relation to habitat area and habitat isolation. Oecologia 137:591–602
- Krüger T, Nipkow M (2015) Rote Liste der in Niedersachsen und Bremen gefährdeten Brutvögel. Informationsdienst Naturschutz Niedersachsen, 182–255.
- Lengyel S, Déri E, Magura T (2016) Species richness responses to structural or compositional habitat diversity between and within grassland patches: a multi-taxon approach. PLoS ONE 11:e0149662
- Leuschner C, Ellenberg H (2017) Ecology of central european non-forest vegetation: coastal to alpine, natural to man-made habitats: vegetation ecology of Central Europe. Springer International Publishing, New York
- Lindborg R, Plue J, Andersson K, Cousins SA (2014) Function of small habitat elements for enhancing plant diversity in different agricultural landscapes. Biol Conserv 169:206–213
- Lindgren JP, Cousins SA (2017) Island biogeography theory outweighs habitat amount hypothesis in predicting plant species richness in small grassland remnants. Landscape Ecol 32:1895–1906
- Loos J, Batáry P, Grass I, Westphal C, Bänsch S, Baillod AB, Hass AL, Rosa J, Tscharntke T (2019) Vulnerability of ecosystem services in farmland depends on landscape management. In: Atlas of ecosystem services. Springer, New York, pp 91–96
- Lyons A, Ashton PA, Powell I, Oxbrough A (2018) Epigeal spider assemblage responses to vegetation structure under contrasting grazing management in upland calcareous grasslands. Insect Conserv Diversity 11:383–395
- Lyons A, Ashton PA, Powell I, Oxbrough A (2017) Impacts of contrasting grazing management on plants and carabid beetles in upland calcareous grasslands. Agr Ecosyst Environ 244:22–34
- Madeira F, Tscharntke T, Elek Z, Kormann UG, Pons X, Rösch V, Samu F, Scherber C, Batáry P (2016) Spillover of arthropods from cropland to protected calcareous grassland-the neighbouring habitat matters. Agr Ecosyst Environ 235:127–133
- Mallinger RE, Gibbs J, Gratton C (2016) Diverse landscapes have a higher abundance and species richness of spring wild bees by providing complementary floral resources over bees' foraging periods. Landscape Ecol 31:1523–1535
- Matthies D, Brauer I, Maibom W, Tscharntke T (2004) Population size and the risk of local extinction: empirical evidence from rare plants. Oikos 105:481–488

- McKinney ML, Lockwood JL (1999) Biotic homogenization: a few winners replacing many losers in the next mass extinction. Trends Ecol Evol 14:450–453
- Melts I, Lanno K, Sammul M, Uchida K, Heinsoo K, Kull T, Laanisto L (2018) Fertilising semi-natural grasslands may cause long-term negative effects on both biodiversity and ecosystem stability. J Appl Ecol 55:1951–1955
- Oksanen J, Blanchet F, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin P, O'Hara R, Simpson G, Solymos P (2018) vegan: community ecology package. R package version 2.4–4
- Pollard E (1977) A method for assessing changes in the abundance of butterflies. Biol Conserv 12:115–134
- Polus E, Vandewoestijne S, Choutt J, Baguette M (2007) Tracking the effects of one century of habitat loss and fragmentation on calcareous grassland butterfly communities. Biodivers Conserv 16:3423–3436
- Poschlod P, Bonn S (1998) Changing dispersal processes in the central European landscape since the last ice age: an explanation for the actual decrease of plant species richness in different habitats? Acta Botanica Neerlandica 47:27–44
- Poschlod P, WallisDeVries MF (2002) The historical and socioeconomic perspective of calcareous grasslands—lessons from the distant and recent past. Biol Conserv 104:361–376
- Riibak K, Reitalu T, Tamme R, Helm A, Gerhold P, Znamenskiy S, Bengtsson K, Rosén E, Prentice HC, Pärtel M (2015) Dark diversity in dry calcareous grasslands is determined by dispersal ability and stress-tolerance. Ecography 38:713–721
- Römermann C, Bernhardt-Römermann M, Kleyer M, Poschlod P (2009) Substitutes for grazing in semi-natural grasslands–do mowing or mulching represent valuable alternatives to maintain vegetation structure? J Veg Sci 20:1086–1098
- Rösch V, Tscharntke T, Scherber C, Batáry P (2015) Biodiversity conservation across taxa and landscapes requires many small as well as single large habitat fragments. Oecologia 179:209–222
- Settele J, Steiner R, Reinhardt R, Feldmann R, Hermann G (2015) Schmetterlinge. Die Tagfalter Deutschlands. Ulmer, Stuttgart
- Soderstrom B, Svensson B, Vessby K, Glimskar A (2001) Plants, insects and birds in semi-natural pastures in relation to local habitat and landscape factors. Biodivers Conserv 10:1839–1863
- Steffan-Dewenter I, Tscharntke T (2000) Butterfly community structure in fragmented habitats. Ecol Lett 3:449–456
- Storchová L, Hořák D (2018) Life-history characteristics of European birds. Glob Ecol Biogeogr 27:400–406
- Südbeck P, Weick F (2005) Methodenstandards zur Erfassung der Brutvögel Deutschlands. Mugler, Radolfzell
- Tscharntke T, Steffan-Dewenter I, Kruess A, Thies C (2002) Contribution of small habitat fragments to conservation of insect communities of grassland–cropland landscapes. Ecol Appl 12:354–363
- Valkó O, Zmihorski M, Biurrun I, Loos J, Labadessa R, Venn S (2016) Ecology and conservation of steppes and semi-natural grasslands. Hacquetia 15:5–14
- von Drachenfels O (2016) Kartierschlüssel für Biotoptypen in Niedersachsen unter besonderer Berücksichtigung der gesetzlich geschützten Biotope sowie der Lebensraumtypen von Anhang I der FFH-Richtlinie, Hannover.
- WallisDeVries MF, Poschlod P, Willems JH (2002) Challenges for the conservation of calcareous grasslands in Northwestern Europe: integrating the requirements of flora and fauna. Biol Conserv 104:265–273
- Weidemann HJ, Köhler J (1996) Nachtfalter: Spinner und Schwärmer. Naturbuch-Verlag, Augsburg
- Wenzel M, Schmitt T, Weitzel M, Seitz A (2006) The severe decline of butterflies on western German calcareous grasslands during the last 30 years: a conservation problem. Biol Conserv 128:542–552
- Willems JH (2001) Problems, approaches, and results in restoration of Dutch calcareous grassland during the last 30 years. Restor Ecol 9:147–154
- Wilson JB, Peet RK, Dengler J, Partel M (2012) Plant species richness: the world records. J Veg Sci 23:796–802
- Zub P (1996) Die Widderchen Hessens. Mitteilungen des Internationalen Entomologischen Vereins [Suppl] IV, 1–122.
- Zulka KP, Abensperg-Traun M, Milasowszky N, Bieringer G, Gereben-Krenn BA, Holzinger W, Hölzler G, Rabitsch W, Reischütz A, Querner P (2014) Species richness in dry grassland patches of eastern Austria: a multi-taxon study on the role of local, landscape and habitat quality variables. Agr Ecosyst Environ 182:25–36

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Jacqueline Loos^{1,2} Jochen Krauss³ · Ashley Lyons^{4,5} · Stephanie Föst¹ · Constanze Ohlendorf¹ · Severin Racky¹ · Marina Röder¹ · Lennart Hudel¹ · Volker Herfert¹ · Teja Tscharntke¹

- ¹ Agroecology Group, Department of Crop Science, Georg-August University Göttingen, Göttingen, Germany
- ² Institute of Ecology, Faculty of Sustainability Science, Leuphana University Lüneburg, Lüneburg, Germany
- ³ Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Würzburg, Germany
- ⁴ Department of Geography and Environmental Science, Liverpool Hope University, Liverpool, UK
- ⁵ Centre for Conservation Science, RSPB, Haweswater, UK