

Legacy effects of pre-crop plant functional group on fungal root symbionts of barley

Roy, Julien; van Duijnen, Richard; Leifheit, Eva F.; Mbedi, Susan; Temperton, Victoria Martine; Rillig, Matthias C.

Published in: **Ecological Applications**

DOI: 10.1002/eap.2378

Publication date: 2021

Document Version Publisher's PDF, also known as Version of record

Link to publication

Citation for pulished version (APA): Roy, J., van Duijnen, R., Leifheit, E. F., Mbedi, S., Temperton, V. M., & Rillig, M. C. (2021). Legacy effects of pre-crop plant functional group on fungal root symbionts of barley. *Ecological Applications*, *31*(6), Article e02378. https://doi.org/10.1002/eap.2378

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Legacy effects of pre-crop plant functional group on fungal root symbionts of barley

JULIEN ROY D,^{1,2,6} RICHARD VAN DUIJNEN,³ EVA F. LEIFHEIT,^{1,2} SUSAN MBEDI,^{4,5} VICKY M. TEMPERTON,³ AND MATTHIAS C. RILLIG D^{1,2}

¹Institut für Biologie, Freie Universität Berlin, Berlin 14195 Germany ²Berlin–Brandenburg Institute of Advanced Biodiversity Research (BBIB), Berlin 14195 Germany ³Institute of Ecology, Leuphana University of Lüneburg, Lüneburg, Germany ⁴Naturkundemuseum Berlin, Berlin 10115 Germany ⁵Berlin Center for Genomics in Biodiversity Reseach, Berlin 14195 Germany

Citation: Roy, J., R. van Duijnen, E. F. Leifheit, S. Mbedi, V. M. Temperton, and M. C. Rillig. 2021. Legacy effects of pre-crop plant functional group on fungal root symbionts of barley. Ecological Applications 31(6):e02378. 10.1002/eap.2378

Abstract. Arbuscular mycorrhizal (AM) fungi, a group of widespread fungal symbionts of crops, could be important in driving crop yield across crop rotations through plant-soil feedbacks (PSF). However, whether preceding crops have a legacy effect on the AM fungi of the subsequent crop is poorly known. We set up an outdoor mesocosm crop rotation experiment that consisted of a first phase growing either one of four pre-crops establishing AM and/or rhizobial symbiosis or not (spring barley, faba bean, lupine, canola), followed by an AM crop, winter barley. After the pre-crop harvest, carbon-rich organic substrates were applied to test whether it attenuated, accentuated or modified the effect of pre-crops. The pre-crop mycorrhizal status, but not its rhizobial status, affected the richness and composition of AM fungi, and this difference, in particular community composition, persisted and increased in the roots of winter barley. The effect of a pre-crop was driven by its single symbiotic group, not its mixed symbiotic group and/or by a crop-species-specific effect. This demonstrates that the pre-crop symbiotic group has lasting legacy effects on the AM fungal communities and may steer the AM fungal community succession across rotation phases. This effect was accentuated by sawdust amendment, but not wheat straw. Based on the previous observation of decreased crop yield after AM pre-crops, our findings suggest negative PSF at the level of the plant symbiotic group driven by a legacy effect of crop rotation history on AM fungal communities, and that a focus on crop symbiotic group offers additional understanding of PSF.

Key words: agro-ecosystems; amplicon sequence variants; arbuscular mycorrhizal fungi; crop rotation; mesocosm experiment; phylogenetic scale; plant–soil feedback; sustainable agriculture.

INTRODUCTION

Plants influence abiotic and biotic soil properties, which in turn affect plants, defining plant–soil feedbacks (PSF). Over time and plant generations, soils accumulate species-specific pathogens or symbionts, which consequently colonize surrounding plants and their offspring. PSF are positive when plants of a certain species grow better in soil conditioned by individuals of the same species than in soil conditioned by individuals of a different species, and PSF are negative in the reverse case. PSF scale to the ecosystem level to drive plant diversity in natural ecosystems (Van der Putten et al. 2013, Bennet et al. 2017, Teste et al. 2017) and affect crop yield in agricultural systems (Huang et al. 2013). In agriculture, PSF occur particularly during continuous cropping (Mariotte

⁶E-mail: roy.julien@live.fr

et al. 2018). In fact, crop rotations have probably been implemented from empirical observations of decreasing yield (i.e., negative PSF) over succeeding monoculture crops. Understanding PSF mechanisms is crucial because they may steer ecological restoration of degraded ecosystems or improve crop yield (Mariotte et al. 2018).

As symbionts, arbuscular mycorrhizal (AM) fungi occur in the roots of most plants including crops (Brundrett and Tedersoo 2018) and are strong drivers of plant diversity and productivity (Van Der Heijden et al. 2008). AM fungi provide soil nutrients (mostly nitrogen and phosphorus) to the plant in exchange for sugars and lipids (Smith and Read 2008). AM fungi show important functional variation such as the degree of soil and root colonization, investment in spores, propagule sources, phenology, nutrient preferential uptake and ecosystem specialization (Hart and Reader 2002, Hart et al. 2002). Different lineages within AM fungi drive positive or negative PSF (Koziol and Bever 2019). Therefore, the

Manuscript received 14 February 2020; revised 23 November 2020; accepted 14 January 2021; final version received 10 May 2021. Corresponding Editor: Hinsby Cadillo-Quiroz.

community composition of AM fungi is of importance in affecting plant growth (van der Heijden et al. 1998, Maherali and Klironomos 2007, Hoeksema et al. 2010). Cover cropping with AM crops can increase root colonization by AM fungi in the subsequent crop (Thompson 1987, Lekberg and Koide 2005, Bowles et al. 2016). However, whether preceding crops (hereafter referred to as pre-crops) have a legacy effect on the AM fungal community associated with the following crop remains poorly understood (Berruti et al. 2018, Campos et al. 2018).

Designing crop rotation schemes that acknowledge the role of AM fungi requires understanding the factors affecting AM fungal communities. Climate, edaphic factors such as fertility or pH, and vegetation characteristics all simultaneously affect AM fungal communities (Davison et al. 2015). Host-specificity of AM fungi is generally low (Davison et al. 2015, Lekberg and Waller 2016), although preference has been observed (Vandenkoornhuyse et al. 2002, Torrecillas et al. 2012). Rather than plant taxonomic identity, plant traits and functional group (such as herbs, forbs, and grasses or ecological strategies such as resource acquisitive or conservative) drive the recruitment of AM fungi (Chagnon et al. 2015, Davison et al. 2015, López-García et al. 2017) to form an association where plant and fungal ecological strategies are tightly coupled (Chagnon et al. 2013). A focus on plant functional group in rotation schemes (Lekberg and Koide 2005) may offer a predictive and mechanistic understanding of PSF beyond metrics of plant evolutionary relatedness and soil physicochemical parameters (Mariotte et al. 2018).

Fertilization changes the nutritional demand of plants and therefore its investment into mycorrhiza (Johnson 2010, Treseder et al. 2018). While mineral N and P fertilization has mostly been studied, few studies have addressed the influence of organic amendment on AM fungal communities and their symbiosis with plants (Yang et al. 2020). In parallel with building soil organic matter stocks and increasing soil C storage, organic amendments are promising options for decreasing N leaching over winter and to supply the leftover N to the following crop by stimulating microbial nitrogen immobilization. Undoubtedly, AM fungi (indirectly) contribute to decomposition and the uptake of newly available nutrients (Leifheit et al. 2015, Yang et al. 2020). As a consequence, organic amendment probably affects the AM fungal community through changes in soil stoichiometry and plant investment into mycorrhiza (Johnson 2010, Treseder et al. 2018). It is unknown whether extrinsic inputs of nutrients, particularly in the form of organic carbon-rich substrate (high carbon amendment, HCA), affect AM fungal community composition independently or in interaction with the crop rotation strategy.

It is still uncertain if ecological or functional specialization of AM fungi occurs below or above the species level, or even at higher taxonomic ranks (Sanders and Rodriguez 2016). On the one hand, a turnover of distantly related AM fungi is observed along broad environmental gradients suggesting deep phylogenetic signal in the ecology of AM fungi (Powell and Sikes 2014, Roy et al. 2019). On the other hand, a high variability for certain traits is observed within AM fungal species (Munkvold et al. 2004, Mensah et al. 2015). For example, differences in functional consequences (plant performance and nutrient uptake) can be large among plants colonized by genetically different populations or individuals of one AM fungal species (Koch et al. 2006). Testing whether dissimilarity in community composition among and between treatments is due to the recruitment of relatively closely or distantly related AM fungi (Roy et al. 2019) is important to understand the level of phylogenetic differentiation in AM fungi that induce different functions in soil or in their effects on plants. This will further help identify the ecologically relevant level of diversity to be conserved or engineered and to define bioindicators (i.e., clades that correlate with plant yield).

We set up an outdoor mesocosm crop rotation experiment that consisted of a first phase growing AM or non-AM and/or rhizobial or non-rhizobial crops followed by an AM crop (Appendix S1: Fig. S1). After the pre-crop harvest, carbon-rich organic substrates were applied. Yield results suggested a negative PSF at the level of plant symbiotic group (continuous cropping of AM crops), independently of organic fertilization (van Duijnen et al. 2018). Here, we characterized root and soil AM fungal communities using molecular techniques and, coupled with recent bioinformatic and phylogenetic sequence analyses and microscopy techniques, we asked the following questions: (1) Does the crop symbiotic functional group imprint (i) composition or (ii) richness of AM fungi associated with a following AM crop, and is it observable in roots and soil? (2) If a pre-crop legacy effect is observed, is this legacy attenuated, accentuated or modified by organic amendment? (3) Do the traits of AM fungi reflect community richness or composition? (4) Which community attribute (richness, composition, and traits) best correlates with crop yield?

MATERIAL AND METHODS

Mesocosm set-up

The experiment was conducted outside at the Leuphana University garden facilities, Lüneburg, Germany $(53^{\circ}14'23.8" \text{ N}, 10^{\circ}24'45.5" \text{ E})$. Mesocosms (rectangular pots; edge length of 37.5 cm at the top; edge length of 26.5 cm at the bottom; height of 37 cm; resulting volume of 38 L) were filled with soil originating from the top 0–30 cm of the experimental farm Hohenschulen of the Christian-Albrechts-University in Kiel $(54^{\circ}19'05.6" \text{ N}, 95^{\circ}8'38.8" \text{ E})$. The soil is sandy loam (Cambic Luvisol) and has a history of agricultural practice, including cropping with a mixture of catch crops (such as clover and lupine) without fertilization in the

growing season before the experiment started, and with maize the season before that, fertilized with slurry and triple superphosphate. The soil had a total of 1.26% C, 0.14% N, a C:N ratio of 9.2, and a pH of 6.0 at the start of the experiment.

Crop species choices

The experiment consisted of two phases: during the first, conditioning phase (hereafter referred to as t1), mesocosms were cultivated with one of four different pre-crops that varied in their symbiotic associations with root microbes (i.e., plant symbiotic groups). Plant symbiotic groups focused on two plant-microbe symbioses: the rhizobial and AM symbioses. Four plant species were used: spring canola (Brassica napus cv. Medicus, NPZ; non-rhizobial/non-AM), white lupine (Lupinus albus cv. Energy, Feldsaaten Freudenberger; rhizobial/ non-AM), spring barley (Hordeum vulgare cv. Barke, Saatzucht Breun; non rhizobial/AM), and faba bean (Vicia faba cv. Tiffany, NPZ; rhizobial/AM). Thus, each plant species represents a combination of the two symbiotic groups, and the AM or rhizobial plant symbiotic group is replicated with two plant species. The plant AM status is not correlated with the plant rhizobial status. This allows to test the independent effect of each symbiotic group. The statistical interaction between the AM and rhizobial status therefore represents the pre-crop species-specific effect (see Statistical analyses of AM fungal community richness and composition). The pre-crop mixed symbiotic group is confounded with speciesspecific additional evolutionary history and ecophysiological properties. Therefore, no inference can be drawn on the legacy effect of the pre-crop's mixed symbiotic group. However, inferences can be drawn on the legacy effect of each single symbiotic status (mycorrhizal or rhizobial).

The succeeding crop was winter barley (Hordeum vulgare, cv. Antonella, Nordsaat Saatzucht, non-rhizobial/ AM). Winter barley was chosen because it is widely used in agriculture in Germany. Winter barley is a close relative to spring barley (they are the same species). Withinspecies (Semchenko et al. 2017) and even withinpopulation specialization of soil microbes to different plant genotypes (Eck et al. 2019) may induce PSF. In this study, growing winter barley after spring barley represents a treatment of a plant species being grown in a soil conditioned by the same species (conspecific). Growing a second distantly related crop to spring or winter barley (for instance, faba bean) but of similar symbiotic functional group (for instance, AM) allows us to test legacy effect at the plant symbiotic functional group (i.e., growing an AM crop in soils conditioned by AM or non-AM crops, e.g., Teste et al. 2017). If not observed, having two distantly related pre-crops would inform whether there is stronger legacy effect between conspecifics than heterospecifics, which could be due to a degree of phylogenetic signal in the AM symbiosis.

Crop rotation and organic amendment timing

Pre-crops (n = 15 pots per pre-crops) were sown on May 2016 along with a mineral fertilization. Fertilization followed recommendations specific for each crop species and standard agricultural German practices. Following pre-crop harvest (one week during late August/ early September 2016), three types of carbon-rich substrates (high carbon amendment, HCA) were applied (n = 5 pots per pre-crops): wheat straw (W), sawdust (S), or no amendment (No). The HCA treatment was applied one week after pre-crop harvest by mixing the HCA material within the top 10 cm of the soil. The soil in the no-amendment treatment received the soil disturbance as well. Winter barley was sown on October 2016. Soil of winter barley was fertilized with a total of 160 kg N/ha, separated in equal additions on three time points. A control with no pre-crop and no HCA treatment was also conducted in three additional pots (hereafter referred to as Control), sown with winter barley after pre-crop harvest time. More details of the experiment, climatic data and level of fertilization are reported in van Duijnen et al. (2018).

Soil sampling

Soil samples were taken at three time points: at the beginning of the experiment before sowing any plants (t0, May 2016), after the pre-crop harvest (t1, one week during late August/early September 2016), and at crop harvest (t2, one day during late June 2017). The t0 samples (n = 6) were randomly cored from the soil pile collected to set up the mesocosms. At t1, five soil cores of 1 cm width and 10 cm depth were randomly cored per pot after the removal of the precrops, and pooled to give a composite sample for each pot, yielding 60 t1 soil samples and the additional three samples with no pre-crop. At t2, five soil cores of 5 cm width and 10 cm depth were randomly cored after the removal of the crop plants, and pooled to give a composite sample for each pot, yielding 60 t2 soil samples and the additional three samples with no pre-crop and no HCA. Soil samples for assessing soil AM fungal abundance were air-dried. Soil samples devoted to molecular analyses were kept in liquid nitrogen and at -80°C before lab processing.

Root sampling

Root sampling was performed during the second phase of the experiment (t2, crop cultivation), yielding 60 t2 root samples and the additional three root samples of winter barley with no previous pre-crop and no HCA. The entire root system per pot was sampled. Roots were thoroughly washed with tap water. Root samples for assessing root colonization by AM fungi were air-dried. Root samples devoted to molecular analyses were kept in liquid nitrogen and at -80° C before lab processing.

Molecular analyses of AM fungal communities

Molecular analyses of AM fungi were conducted on a total of 195 samples: six t0 soil samples, 60 t1 soil samples (four pre-crops \times 15 replicates), and three control pots, 60 t2 soil samples (four pre-crops \times three HCA treatments \times five replicates) and the three control pots, 60 t2 root samples (four pre-crops \times 3 HCA treatments \times five replicates) and the three control pots.

DNA extraction, PCR amplification, and Illumina sequencing library preparation.—For the soil samples, DNA was extracted from 250 mg of the pooled soil cores using the PowerSoil DNA isolation kit (MoBio Laboratories, Carlsbad, California, USA), following the manufacturer's instructions. For the root samples, we use the same procedure as for the soil samples except that we used 100 mg of crushed freeze-dried roots. The protocol for DNA extraction, PCR amplification, and Illumina sequencing library preparation followed Roy et al. (2017). Briefly, using a nested-PCR approach, we amplified the D1-D2 region of the large subunit (LSU) of rDNA using universal fungal primers bound to an 8 nt long index for sample multiplexing, first targeting a region spanning the small subunit-internal transcribed spacer-large subunit (SSU-ITS-LSU) region using AM fungi-specific primer mixtures (Krüger et al. 2009). The purified final PCR products were pooled on an equimolar basis and sequenced on an Illumina MiSeq platform at the Berlin Center for Genomics in Biodiversity Research (BeGenDiv), Berlin, using 2×300 base pair paired-end sequencing. Raw reads are available at ENA under study accession number PRJEB36419.

Bioinformatics sequence processing.—Amplicon reads were analyzed using DADA2 (Callahan et al. 2016) in R (R Core Team 2017) to obtain denoised, chimera-free, nonsingleton, AM fungal exact sequence variants (ESVs) as implemented in Roy et al. (2019). AM fungal ESVs were identified using BLAST search against reference AM fungal rDNA sequences (Krüger et al. 2012) with a minimum of 90% coverage and of 90% to our targeted region. Sequences without this requirement were discarded. Additional putative PCR errors were identified using LULU in R (Frøslev et al. 2017) and were discarded. ESV sequences are available at ENA under accession numbers LR761341-LR761569. The ESV contingency table and ESV taxonomy and related sample metadata are freely *available online*.^{7,8}

Statistical analyses of AM fungal community richness and composition.—All statistical analyses were conducted in R 3.4.1 (R Core Team 2017). Before community analyses, ESV read counts per sample were normalized by rarefying sample read count to the lowest number of reads.

⁷ https://doi.org/10.20387/bonares-sea4-n5sp

8 https://doi.org/10.20387/bonares-xnhz-xw6e

The ESV richness per sample before and after rarefaction was highly correlated (Pearson's R = 0.97). Similarly, pairwise sample Bray-Curtis dissimilarities before (ESV read count per sample normalized by sample read count) and after rarefaction were highly correlated (Mantel Pearson's R = 0.99).

AM fungal richness.—AM fungal richness was calculated by the number of ESVs and analyzed using ANOVA. We tested the effect of crop rotation phase, pre-crop symbiotic group, HCA, and plant compartment (roots vs. soil). All factors could not be tested in one single model given some factors were restricted to a particular rotation phase (e.g., HCA and plant compartment factors are only relevant at t2). Therefore, we ran separate models using different sample sets to test the effects of these factors. We used all samples to test whether there was a difference in AM fungal richness between crop rotation phases; we used the t1 samples (conditioning phase) to test whether there was a difference in AM fungal richness between pre-crop symbiotic groups at t1; we used the t2 samples (PSF phase) to test whether there was a difference in AM fungal richness between pre-crop symbiotic groups, HCA treatments or plant compartments at t2. In addition, mixed linear models implemented in lme4 (Bates et al. 2015) were used to test the differences of richness between t1 and t2 crop rotation phases, and to test the effect of plant compartment at t2, setting the pot identity as random factor. The significance of these factors was tested compared to a model without fixed factors using log-likelihood ratio tests.

AM fungal community composition.—The change in the composition of the AM fungal community composition was measured using Bray-Curtis dissimilarity. We used permutational multivariate ANOVA (PERMANOVA; Anderson 2001) to estimate the part of variance explained by each factor. Statistical departure from a null model of random treatment assignation of samples was assessed using 999 Monte-Carlo permutations of samples. As for AM fungal richness, different sets of samples were used to run separate models in order to test for effects of crop rotation phases (all samples), of pre-crop symbiotic group (t1 samples), and of pre-crop symbiotic group, HCA, and plant compartment (t2 samples). In addition, a distance-based redundancy analysis (Legendre and Anderson 1999) was used to test for differences of community composition between crop rotation phases (t1 and t2 samples) and to test for differences between plant compartments (t2 samples) while partialing out the variance of pot identity. Similarly to richness analyses, the control pots were removed from the community composition analyses because of low sample size (n = 3) and to focus on treatment differences. Bray-Curtis dissimilarities among samples were visualized using non-metric multidimensional scaling (NMDS) ordination. Control pots were included along with the other samples to visualize their genetic compositional similarity to other treatments. The direction and strength of the correlation of phylotypes defined at different phylogenetic distances (see Phylogenetic dissimilarity of AM fungal communities) to the community dissimilarities were measured using multivariate linear regression between phylotype relative abundance and sample coordinates in the NMDS space (Oksanen et al. 2016). Multivariate community analyses were conducted in vegan (Oksanen et al. 2016).

Phylogenetic dissimilarity of AM fungal communities.— We analyzed the dissimilarity of AM fungal communities across phylogenetic depths in order to test whether dissimilarity in community composition among and between treatments is due to the recruitment of relatively closely or distantly related AM fungi. We screened community dissimilarities across phylogenetic depths from the ESV level to the order level and tested at which phylogenetic depth the correlation between community composition and treatments was the highest. To infer the phylogenetic relatedness between ESVs, reference partial SSU-ITS-LSU rDNA sequences (Krüger et al. 2012) were first aligned, followed with the alignment of ESVs to the reference backbone alignment using MAFFT (Katoh et al. 2002). A phylogenetic tree was built using all ESVs and all reference sequences, by conducting a bootstrap analysis (100 bootstraps) and searching for the best-scoring maximum-likelihood tree under a GTRGAMMA model of nucleotide substitution in RAxML (Stamatakis 2014). Reference sequences were pruned and ESVs were clustered into phylotypes at phylogenetic distances (in substitution per site, hereafter referred to as subs/site) spanning the entire phylogenetic tree and keeping cluster monophyly using BDTT in R (Groussin et al. 2017). A phylotype-by-sample count matrix was generated at each phylotype resolution and the PERMANOVA models were fitted as described in AM fungal community composition. We generated a null distribution of PERMANOVA R^2 across phylogenetic depths by randomizing ESV phylogenetic relatedness while keeping constant community composition and repeating the analysis. Uncertainty in phylogenetic relatedness between ESVs was accounted for by repeating the analysis for the 100 bootstrapped phylogenetic trees. We accepted a genuine phylogenetic signal when explained variance was higher than the 5-95% null distribution obtained from phylogenetic randomizations and that the PERMANOVA P value was below 0.05 for more than 90% of the bootstrapped phylogenetic trees.

Analyses of AM fungal traits

Root colonization and structural investment.—Different intra-radical AM fungal structures were recorded separately, including arbuscules, hyphae, vesicles and spores. Dried root systems were cut into 1–2 cm fragments, placed in 10% KOH and left in a water bath at 80°C for

30 minutes. Bleached roots were rinsed with tap water and acidified with 1% HCl for 10 minutes and stained with 0.05% Trypan Blue (Phillips and Hayman 1970). Approximately 20 root fragments were mounted on glass slides. The percentage of roots colonized by AM fungi, as well as separate AM fungal structures, was quantified at 200 magnification using the magnified root intersections method (McGonigle et al. 1990). The percentage of microsclerotia and dark-septate pigmented hyphae, reflecting the colonization by non-mycorrhizal rootcolonizing fungi, was also reported.

Soil hyphal colonization.—Soil colonization by AM fungi was determined as hyphal length in m/g soil (modified from Leifheit et al. 2015). Briefly, hyphae were extracted from 4.0 g of soil, stained with 0.05% Trypan Blue and quantified following the line-intersect method at $200 \times$ magnification (Jakobsen et al. 1992). Hyphae were identified as AM fungi if they were aseptate, darkto light-blue stained and with characteristic unilateral angular projections according to Mosse (1959). All other hyphae were counted as non-AM fungi. Very short pieces of hyphae were discarded because a reliable morphological identification was not possible. The effects of pre-crop symbiotic group and HCA were tested using ANOVA.

Statistical analyses of community traits.—The effects of pre-crop symbiotic group and HCA on hyphal root colonization, other intra-radical structures and soil hyphal length were tested using ANOVA. Furthermore, the direction and strength of the correlation of the different intra-radical AM fungal structures and soil hyphal length to the community dissimilarities at different phylogenetic distances (see Phylogenetic dissimilarity of AM fungal communities) was measured using multivariate linear regression between the AM fungal structures and sample coordinates in the NMDS space.

Correlation of AM fungal community attributes with winter barley yield.-Winter barley yield data were originally described in van Duijnen et al. (2018), from which we extracted total grain yield (mass) and total grain N uptake (or yield, the N concentration of the seeds multiplied by grain yield). We used multivariate and univariate analyses to identify which community attributes (richness, composition and traits) best correlate with winter barley yield. The direction and strength of the correlation of yield to the community dissimilarities at different phylogenetic distances (see Phylogenetic dissimilarity of AM fungal communities) was measured using multivariate linear regression between yield and sample coordinates in the NMDS space. We also summarized covariation among all community attributes and winter barley yield (1) using Principal Component Analysis (PCA) of centered and scaled data implemented in ade4 (Dray and Dufour 2007) and (2) using pairwise Pearson's correlations. The analyses were assessed at phase

t2 (where the AM symbiosis takes place, although one could argue that the abundance of phylotypes associated to pre-crops in the conditioning t1 phase can affect crop biomass at t2), and to root compartment (to focus on a likely active symbiosis).

RESULTS

General description of the sequencing results and AM fungal diversity

AM fungal ESVs were assigned to three orders (Archaeosporales, Diversisporales, Glomerales), five families (Archaeosporaceae, Acaulosporaceae, Diversisporaceae, Claroideoglomeraceae, Glomeraceae), and six genera (*Archaeospora*, *Acaulospora*, *Diversispora*,

Claroideoglomus, Rhizophagus, Funneliformis; Fig. 1a). The number of phylotypes strongly decreased from 229 ESVs to 7 phylotypes at 0.15-0.20 subs/site (substitutions per site). These phylotypes were composed of ESVs always annotated to the same species, or to the same genus (Fig. 1b). Four clades were composed of several lineages annotated to different species. For instance, this was the case for Rhizophagus node 338 (R. irregularis and a second unknown lineage), Funneliformis node 392 (F. caledonius, F. mosseae, and F. constrictus), Claroideoglomus node 306 (C. claroideum, C. etunicatum, C. luteum, and an unknown lineage), Diversispora node 276 (D. eburnea, D. celata, D. epigaea, and two unknown lineages). The Rhizophagus node 324, the Acaulospora node 267, and the Archaeospora node 231 were represented by ESVs annotated to a single species,



FIG. 1. Overview of arbuscular mycorrhizal (AM) fungal diversity. (a) Maximum likelihood phylogenetic tree of AM fungal exact sequence variants (ESVs). Clades delineated at a 0.2 genetic distance (mean substitution/site, subs/site) are highlighted with alternating gray and blue rectangles. The taxonomic annotation of subclades within clades at 0.2 subs/site is the majority consensus annotation based on the taxonomic identity at the species level. Scale bar indicates 0.1 mean subs/site. (b) Number of clades (operational taxonomic units, OTUs) across phylogenetic distances. (c) Overall number of ESVs and relative abundance of clades at 0.2 subs/site.

respectively, *Rhizophagus cerebriformis, Acaulospora cavernata*, and *Archaeospora schenckii*. All phylotypes at 0.2 subs/site were composed of a minimum of 10 ESVs (*Diversispora* node 276) to a maximum of 67 ESVs (*Funneliformis* node 392). At 0.4 subs/site, all Glomerales EVS grouped into one phylotype while the other phylotypes (*Diversispora, Acaulospora cavernata*, and *Archaeospora schenckii*) remained delineated. Overall, *Acaulospora cavernata* node 267 was the most abundant phylotype in terms of read number, whereas *Rhizophagus* node 338 and *Funneliformis* node 292 were the most ESV-rich phylotypes (Fig. 1c).

Does the crop symbiotic group imprint the succeeding crop's AM fungal composition?

Overall, community variance was better explained at the species to genus level (0.15–0.20 subs/site) than at the within-species level (ESVs) for all factors investigated (Fig. 2; Appendix S1: Table S1). In the following, we therefore focused on the dynamics of clades delineated at this phylogenetic level.

The original soil was mostly composed of *Diversispora* node 276 (66% of the reads) and *Acaulospora cavernata* node 267 (33%). *Archaeospora schenckii* node 231 (0.4%), *Rhizophagus* node 338 (0.1%), and *Funneliformis* node 392 (0.1%) were present but rare, while *Rhizophagus cerebriformis* node 324 and *Claroideglomus* node 306 were not detected (Fig. 3; Appendix S1: Fig. S2, Table S2). Overall, we observed differences in

community composition among rotation phases (0.2 subs/site: $R^2 = 0.095$, P = 0.001) due to the selection of different phylotypes by different pre-crops and the persistence and accentuation of this effect, even when accounting for pot autocorrelation (0.2 subs/site: $R^2 = 0.095\%$, P = 0.001).

At t1, after pre-crop conditioning, we observed an effect of the pre-crop mycorrhizal group on the AM fungal community composition left in soil (ESV, $R^2 = 0.04$, P = 0.01; 0.2 subs/site: $R^2 = 0.1, P = 0.002$), but not of its rhizobial group (P > 0.05 in all cases) (Fig. 2b; Appendix S1: Table S1). At the species to genus level (0.2 subs/site), community composition strongly converged between the two pre-crops with mycorrhizal associations, and diverged from the two pre-crops with no mycorrhizal associations (Fig. 4a, b). Rhizophagus node 338, Funneliformis node 392, and Acaulospora cavernata node 267 were the main clades driving the differences in community composition after the conditioning by AM or non-AM pre-crops (Fig. 4b; Appendix S1: Table S3). Compared to t0, Rhizophagus node 338 and Funneliformis node 392 strongly increased in abundance, especially in pots conditioned with AM pre-crops (Fig. 3a; Appendix S1: Table S2). Conversely, Acaulospora cavernata node 267 was slightly depleted in soil conditioned by AM pre-crops. Claroideoglomus node 306 sporadically appeared after AM pre-crops (Fig. 3a). Archaeospora schenckii node 231 was only abundant in control pots (i.e., without pre-crop) and depleted from the other soils (Figs. 3a and 4b; Appendix S1: Table S2).



FIG. 2. Variance partitioning of AM fungal community dissimilarities across phylogenetic depth. (a) Variance explained (R^2) by mycorrhizal (AM) and rhizobial (N2) plant symbiotic groups at t1 (conditioning phase of experiment), at each phylogenetic depth used to cluster ESVs into broader OTUs. (b) Variance explained (R^2) by plant mycorrhizal (AM) and rhizobial (N2) groups, high carbon amendment (HCA) and plant compartment at t2 (harvest), at each phylogenetic depth used to cluster ESVs into OTUs. Filled and open symbols, respectively, indicate P < 0.05 or P > 0.05 after 1,000 sample permutations for >90% of the 100 phylogenetic rees obtained with bootstrapping. The gray area represents the 95% percentiles interval of the R^2 for 100 randomizations of ESVs phylogenetic relatedness while keeping community composition constant.



FIG. 3. Distribution of AM fungal clades across rotation phases, pre-crops, high carbon amendment, and plant compartment. Distribution across (a) pre-crop species and (b) high carbon amendment treatments is emphasized. The size of the point represents clade relative read abundance. The color of the points represents clade relative occurrence. Communities are t0, original soil communities; t1, soil communities after pre-crop conditioning phase; and t2, root communities at crop harvest. Abbreviations are Can, canola; Lup, white lupine; Bar, spring barley; Fab, faba bean; Control, control; No, no amendment; W, wheat straw; S, sawdust.

At t2 (PSF phase), the recruitment of AM fungi of different species and genera between AM and non-AM pre-crops persisted and was accentuated in the roots of winter barley (0.2 subs/site, $R^2 = 0.07$, P = 0.004, Appendix S1: Table S1). Root communities grouped according to the mycorrhizal group of the conditioning pre-crop, with a particularly strong convergence of the winter barley root community in the pots conditioned by the two AM pre-crops (Fig. 4d). Rhizophagus node 338 and Acaulospora cavernata node 267 were again the main drivers of these differences (Fig. 4d; Appendix S1: Table S3). Rhizophagus node 338 strongly persisted and even increased in abundance in the roots of winter barley in soils conditioned by AM pre-crops whereas Acaulospora cavernata node 267 further decreased (Fig. 3a; Appendix S1: Table S2). Other distantly related fungi from different families and even orders were more abundant in the roots of winter barley when grown in soils conditioned by non-AM than

AM pre-crops, e.g., *Claroideoglomus* node 306 and *Diversispora* node 276 (Figs. 3a and 4d; Appendix S1: Table S2).

In soil, the imprint of pre-crops was less evident (at 0.2 subs/site, $R^2 = 0.03$, P = 0.136) and soil communities largely retained clades abundant in the original soil (Fig. 3a). We observed differences between root and soil communities from the ESV level ($R^2 = 0.267, P = 0.001$) to the selection of fungi from different families or orders $R^2 = 0.137$, P = 0.001) (Fig. 2b; (0.4)subs/site, Appendix S1: Table S1). Acaulospora cavernata node 267 and Rhizophagus node 338 were the main drivers of the differences between root and soil communities (Fig. 3; Appendix S1: Fig. S2; Table S3). Acaulospora cavernata node 267 was relatively more abundant in soil than roots (Fig. 3a; Appendix S1: Table S2). This clade was the dominant clade in all samples, both in the original soil and throughout the experiment irrespective of the treatments, including in the control pots and has opposite



FIG. 4. Unconstrained ordination analysis of AM fungal community dissimilarities among pre-crops and high carbon amendment at 11 and t2. Dissimilarities (Bray-Curtis) are split among rotation timing, (a, b) t1, pre-crop harvest, and (c–f) t2, winter barley harvest, prior to ordination analysis. Dissimilarities were computed at two phylogenetic depths, (a, c, e) ESV level and (b, c, f) 0.2 subs/site. The centroid of sample distribution per treatment is shown with bigger dots that connect to the respective samples. For t2 ordinations, centroids for (c, d) pre-crops and (e, f) high carbon amendment (HCA) are shown separately. At t2 (c, f), the analysis presents root communities. Supported correlations of ESVs (P = 0.001) or clades at 0.2 subs/site (P = 0.01) to dissimilarities are overlayed as vectors. Statistical differences among treatments are presented in Appendix S1: Table S1. All correlations of clades at 0.2 subs/site to dissimilarities are presented in Appendix S1: Table S3.

trends to *Rhizophagus* node 338 and *Funneliformis* node 392 (Fig. 3; Appendix S1: Fig. S1). Significant community dissimilarities remained when accounting for pot autocorrelation indicating consistent differences across

pots ($R^2 = 0.152$, P = 0.001 at 0.2 subs/site) as observed by the distinction in the centroids of soil and root samples due to the non-overlap of numerous soil and root samples (Appendix S1: Fig. S3).

JULIEN ROY ET AL.

Generally, the impact of the plant mixed symbiotic group and/or the crop species-specific effect was statistically absent after accounting for a plant single symbiotic group (i.e., rhizobial or mycorrhizal), at any phylogenetic level, time (t1, t2), and plant compartment (root, soil) (Appendix S1: Table S1). Nonetheless, for t1, a separation of community composition for each pre-crop was observed at the ESV level (Fig. 4a). At t2, root communities conditioned by faba bean (AM/rhizobial) differed from all others (i.e., conditioned by spring barley, canola, or lupine, and the control) (Fig. 4c).

Does the crop symbiotic functional group imprint on the richness of AM fungi associated with winter barley?

On average there were $12.6 \pm 8 \pmod{\pm SD}$ ESVs per sample. ESV richness increased with time $(F_{2,187} = 11.08, P = 2.83 \times 10^{-5};$ Appendix S1: Table S4) from t0 (4.7 ± 2.9 ESVs [mean \pm SD]) to t1 (9.5 ± 5.8 ESVs) to t2 (14.4 ± 8.3 ESVs) due to the recruitment of AM fungi barely detected at t0 (Fig. 5). This effect remained when accounting for pot autocorrelation ($\chi^2 = 16.368, P = 5.217 \times 10^{-5}$). At t1, ESV



FIG. 5. ESV richness of AM fungal communities across rotation phases (t0, t1, t2), pre-crops, high carbon amendment, and plant compartment. Richness across (a) pre-crop species and (b) high carbon amendment treatments is emphasized. The box plots represent the 25th and 75th percentiles (box edges), the median (mid line), and outlying points (whiskers). Data points are shown on the boxplots. The *y*-axes of all panels are on the same scale for comparability among rotation phases and treatments. Statistical comparisons are presented in Appendix S1: Table S4. Abbreviations are Can, canola; Lup, white lupine; Bar, spring barley; Fab, faba bean; No, no amendment; W, wheat straw; S, sawdust; Control, control.

richness in soil after pre-crop harvest was affected by the pre-crop mycorrhizal group ($F_{1,55} = 33.484, P = 3.55 \times$ 10^{-7}), but not its rhizobial group ($F_{1.55} = 1.362$, P = 0.248) (Appendix S1: Table S4): ESV richness was higher after AM pre-crops (spring barley and faba bean, 13 ± 5.6 ESVs) than non-AM pre-crops (lupine and canola, 6 ± 3.4 ESVs; Fig. 5a). This was driven by increasing Rhizophagus node 338 and Funneliformis node 392 ESV richness, whereas Acaulospora node 276 and Archaeospora node 231 showed opposite patterns (Appendix S1: Fig. S2). This observation did not hold at t2, after crop (winter barley) harvest ($F_{1.95} = 1.980$, P = 0.162; Appendix S1: Table S4): we observed an increase in richness after non-AM pre-crops to a similar level as after AM pre-crops (Fig. 5a), due to the selection of Claroideoglomus node 306 and Diversispora node 267 (Appendix S1: Fig. S2). In general, root communities (16.5 \pm 10.4 ESVs) had higher ESV richness than soil communities $(12.4 \pm 4.8 \text{ ESVs})$ (Appendix S1: Table S4); this effect remained when accounting for pot autocorrelation ($\chi^2 = 10.157, P = 0.001437$).

Is the pre-crop legacy effect attenuated, accentuated or modified by organic amendment?

High carbon amendment (HCA) explained a higher part of variance than the mycorrhizal group of the precrops in the community dissimilarities at the harvest of winter barley at t2 (Fig. 2b; Appendix S1: Table S1). We observed strong community differences between the HCA treatments from the ESV level ($R^2 = 0.034$, P = 0.003) to the selection of fungi from different families or orders (0.4 subs/site: $R^2 = 0.175$, P = 0.003; Fig. 2b; Appendix S1: Table S1). The HCA effect was mostly driven by the selection of a *Rhizophagus cerebriformis* clade (node 324) but also by the above-mentioned *Funneliformis* (node 392) and *Rhizophagus* (node 338) clades after sawdust amendment (Fig. 3b; Fig. 4e, f; Appendix S1: Table S3). These three clades, *Rhizophagus cerebriformis* node 324, *Rhizophagus* node 338, and *Funneliformis* node 392 were more abundant in roots (Fig. 3b; Appendix S1: Fig. S4; Table S2), so that the soil community was less effected by HCA than the root community (Appendix S1: Table S1).

At t2, HCA strongly affected ESV richness $(F_{2,95} = 21.032, P = 2.74 \times 10^{-8};$ Appendix S1: Table S4): ESV richness increased after sawdust amendment (20.3 ± 11.7 ESVs) compared to wheat straw (10.3 ± 3.8 ESVs) and no amendment (13.1 ± 4.4 ESVs; Fig. 5b). This effect was more pronounced in roots compared to soil ($F_{2,95} = 3.096, P = 0.049;$ Appendix S1: Table S4), with a higher richness in roots than soil (Fig. 5).

Do the traits of AM fungi reflect community richness or composition?

None of the community traits correlated with community richness or composition (Fig. 6; Appendix S1: Fig. S4; Table S5). Accordingly, neither AM hypha in roots (root colonization: $18.6\% \pm 12.5\%$) or in soil (AM hyphal length: 1.4 ± 0.6 m/g soil), nor other fitnessrelated features such as vesicle ($4.7\% \pm 7.7\%$) and spore production ($8.6\% \pm 10.9\%$) varied with pre-crop mycorrhizal (all P > 0.05) or rhizobial (all P > 0.05) group, or with HCA (all P > 0.05) (Appendix S1: Fig. S5, S6, Table S6, S7). The soil colonization by non-AM fungi did not vary among treatments either (3.2 ± 1.1 m/g soil), but the percentage of microsclerotia in roots was slightly higher in the control ($24.4\% \pm 12.7\%$) than wheat straw ($15.3\% \pm 9.4\%$) or sawdust ($21.3\% \pm 12.4\%$) treatments (Appendix S1: Fig. S5, S6, Table S6, S7).



FIG. 6. Correlation overview between AM fungal community richness, composition, traits and winter barley yield at winter barley harvest t2 phase. (a, b) Principal Component Analysis of AM fungal community richness, composition, traits and winter barley yield. The centroid of sample distribution per treatment is shown with bigger dots that connect to the respective samples. Centroids of treatments are shown separately for (a) pre-crops and (b) high carbon amendment (HCA). (c) Pairwise Pearson's correlations. Only significant correlations at P < 0.05 are shown. Community composition (clade relative abundance and NMDS axes) and richness variables are those calculated at t2 in roots of winter barley, similarly to community traits and winter barley yield.

Which community attribute (richness, composition, and traits) best correlates with crop yield?

Winter barley yield showed a correlation with the community composition in roots, decreasing with communities conditioned with AM pre-crops (Appendix S1: Fig. S4). The correlation was weak and only significant for N yield, not grain yield (Appendix S1: Table S5). The PCA and pairwise correlations revealed overall weak correlations between any of the AM fungal community attributes in winter barley and its yield (Fig. 6). The only correlation was a negative correlation between the abundance of *Rhizophagus* node 338 and winter barley grain yield (Pearson's r = -0.317, P = 0.014) and total N yield (Pearson's r = -0.36, P = 0.005; Fig. 6c). For instance, mean grain yield was 659.4 ± 113.1 and $733.1 \pm 107.5 \text{ g/m}^2$ for presence or absence of *Rhizoph*agus node 338. In contrast, the presence/absence of two other clades, Claroideoglomus node 306 (correlation with N yield, Pearson's r = 0.255, P = 0.052) and Diversispora node 276 (correlation with N yield, Pearson's r = 0.255, P = 0.051), correlated with higher crop yield (Appendix S1: Fig. S7, Table S8).

DISCUSSION

Does the crop symbiotic functional group imprint on the composition or richness of AM fungi associated with a succeeding AM crop and is it observable in roots and soil?

The pre-crop mycorrhizal status affected the richness and composition of AM fungi available to the succeeding crop. This difference, in particular community composition, persisted and increased in the roots of the succeeding crop. The effect of the pre-crop on AM fungal communities was driven by its single symbiotic group (for instance, mycorrhizal), not mixed symbiotic group and/or crop species-specific or conspecific effect. This demonstrates that pre-crop symbiotic group has lasting legacy effects on the AM fungal communities associated with the succeeding crops and may steer the AM fungal community succession across rotation phases. Our previous findings showed a 23% decrease in winter barley vield when grown in soil conditioned by AM pre-crops, N-fixing or not (van Duijnen et al. 2018). Our findings suggest negative PSF at the level of the plant symbiotic group, here of a continuous AM cropping, driven by a legacy effect of crop rotation history on AM fungal communities.

Our result may appear surprising, as it is well established that AM fungi often increase plant growth (Smith and Read 2008). Microbial symbionts do not necessarily imply positive PSF, even when the presence of these symbionts increases plant growth (Bever 2002, Bennet et al. 2017). In fact, it has long been recognized by farmers that the colonization of crop roots by AM fungi can be detrimental to yield. This was experimentally demonstrated (Hoeksema et al. 2010, Koch et al. 2017), pointing to parasitism by AM fungi (Kirkegaard et al. 2008). Farmers implement crop rotation schemes that may involve a breaking crop, i.e., a non-AM host such as from the Brassicaceae family, to decrease the load of AM fungi to the next AM crop, and this has been shown to benefit the following crop in terms of yield (e.g., wheat, Kirkegaard et al. 2008, Angus et al. 2015). Legumes, which establish an atmospheric N-fixing rhizobial symbiosis and most of which are mycorrhizal (e.g., faba bean), are also used during crop rotation with yield benefits to the next crop (Angus et al. 2015). This was not the case here when legumes were the AM host. While several crop species, including cereals (Zhang et al. 2019), generally respond positively to inoculation with AM fungi, this is often less so for cereals (Eo and Eom 2009). Indeed, cereal responses to AM fungi are generally positive but weak, highly crop-specific, and tend to be negative for crops released after the 1950s (Zhang et al. 2019).

High N content, such as in our study, could favor mutualistic interactions but not under high P content (Johnson 2010). We likely had high soil P here, since pre-crops were fertilized with triple superphosphate and because soils used here had long history of agriculture. Even in P-limited soils, AM fungi with tolerance to high N enrichment may not be able to intensively forage for P, since plants can reduce the investment into AM fungi (Treseder et al. 2018). In this study, we do not have AM fungal data under low N input. Unsurprisingly, winter barley yield was strongly reduced in low compared to high N fertilization (van Duijnen et al. 2018). However, in the low N fertilization treatment there was no difference in yield between plants grown after AM or non-AM pre-crops (van Duijnen et al. 2018). Therefore, it is unlikely that the AM fungal community parasitized winter barley under low N input. This is a crucial point for further research, to test whether AM fungi tend to form parasitic associations with crop roots under higher N, but mainly have positive interactions under low N.

Specifically, we identified one Rhizophagus clade (node 338) in particular and one Funneliformis clade (node 392) that were selected after AM pre-crops. However, the original soil, which has a long history of agricultural practice, was mostly composed of an Acaulospora cavernata clade, a Diversispora clade, and an Archaeospora schenckii clade, as observed in 50-yrold re-cultivated barren soils (Roy et al. 2017). Rhizophagus and Funneliformis were almost undetected in the original soil, indicating the strong colonization ability of these fungi. Rhizophagus fungi possess life-history traits of ruderal fungi (Hart and Reader 2002, Chagnon et al. 2013) and are often observed in early stages of ecosystem succession (Nielsen et al. 2016, Roy et al. 2017). Ruderal AM fungi may colonize roots faster after a first crop is sown, giving them an advantage over slowgrowing fungi to colonize the succeeding crop (Verbruggen and Kiers 2010).

Nonetheless, the legacy effect was stronger in roots than in soil. Soil retained members of fungal clades abundant in the original soil (Acaulospora cavernata, Diversispora, and Archaeospora schenckii). Root colonization is intrinsically determined by ability to germinate, propagule sources, growth rate, competitive ability, host and habitat preference, and phenology. It is possible that the time scale of agricultural practices (e.g., rotations) and the time scale of fungal root colonization are not matched. The spore bank and/or outcompeted AM fungi could better colonize roots if conditions are favorable or if time to colonize roots is sufficiently long. Indeed, AM fungal communities are highly dynamic in agricultural systems (Roy et al. 2017) with seasonal succession being observed in natural (Dumbrell et al. 2011) and agricultural ecosystems (Berutti et al. 2018).

If a pre-crop legacy effect is observed, is this legacy attenuated, accentuated, or modified by organic amendment?

We did not observe an interaction between HCA and the pre-crop symbiotic group, indicating that their respective influences on AM fungal communities were independent. HCA induced the most important shifts in AM fungal communities, including strong change in community composition and richness, especially in roots, which was mostly due to sawdust amendment. Sawdust increased AM fungal richness, especially of ruderal fungi. It strongly positively selected Rhizophagus node 338, Funneliformis node 392, and Rhizophagus node 324, and negatively selected Acaulospora cavernata node 267. A 6.3% decrease in yield was observed after sawdust, but not wheat straw (van Duijnen et al. 2018). Therefore, HCA may have an additive effect with crop mycorrhizal group, either positive in the case of sawdust amendment following AM pre-crops, or antagonistic in the case of wheat straw. It is interesting to note that the strongest effect on AM fungal communities (sawdust) was not related to the strongest effect on yield (pre-crop mycorrhizal group).

Do the traits of AM fungi reflect community richness or composition?

The dissimilarities between AM fungal communities were due to recruitment of distantly related lineages, not variants within species. This suggests that relatively ancient evolutionary divergences in AM fungi, at the species level and above, are related to observable ecological and/or functional specialization (Lekberg et al. 2014, Roy et al. 2019) with consequences for crop yield. Given phylogenetic conservatism of morphology and growth traits in AM fungi (Hart and Reader 2002, Powell et al. 2009, Koch et al. 2017) and the congruent deep phylogenetic dissimilarities between AM fungal communities of different treatments (e.g., AM vs. non-AM pre-crops), we expected soil and root colonization to be lower and higher after AM pre-crops, respectively. Higher soil and lower root colonization should induce a low C cost to the plant to support AM fungi while increasing nutrient foraging by AM fungi (Chagnon et al. 2013). This was not observed, and trait data did not reflect community composition. Similarly, a recent study showed that host performance cannot be predicted from AM fungal morphology and growth traits (Koch et al. 2017). They concluded that divergent effects on plant growth among isolates within an AM fungal species may be caused by coevolution between co-occurring fungal and plant populations. Contrary to sterile conditions and single inoculum addition (Hart et al. 2002), competition among the microbial community may blur the correlation between AM fungal lifehistory traits and community composition in natural settings (Maherali and Klironomos 2007). Soil colonization rates by AM fungi were also comparatively low in this experiment (e.g., Rillig et al. 2002), probably due to the overall high nutrient levels in the collected soil and the N fertilization (Treseder 2004). Enhanced phenotyping of AM fungi and of the symbiosis would help understand why the communities that are favored after AM pre-crops negatively affect the succeeding crop yield (Montero et al. 2019).

Which community attribute (richness, composition, and traits) best correlates with crop yield?

The correlation between winter barley yield and the abundance of Rhizophagus node 338 was the best and only one. Rhizophagus fungi, but also Funneliformis, poorly colonize soil but mostly roots (Hart and Reader 2002, Chagnon et al. 2013, Koch et al. 2017). They may not provide nutritional benefits given the poor nutrient foraging and transfer capacity (Chagnon et al. 2013). It is possible that competition and low functional complementarity between coexisting individuals of Rhizophagus and Funneliformis may have reduced the benefit of the symbiosis: while the coexistence of AM fungi from different families can increase plant growth due to functional complementarity (Maherali and Klironomos 2007), coexisting AM fungi within species, genus, or family may have opposite effects (Roger et al. 2013, Yang et al. 2017). The relatively low phylogenetic divergence of coexisting Rhizophagus and Funneliformis fungi might have led to enhanced competition and reduced functional complementarity, with less benefit to the plants (Maherali and Klironomos 2007).

CONCLUSION

Our findings suggest negative PSF at the level of the plant symbiotic group driven by a legacy effect of crop rotation history on AM fungal communities, suggesting that a focus on crop symbiotic group offers additional understanding of PSF. Differences in AM fungal community composition may have functional consequences that should not be neglected in agriculture (Rillig et al. 2019), although this debate is open (Ryan and Graham 2018). Crop rotation design, but also nutrient input strategies, could foster the AM symbiosis towards targeted services. Overall, differences in AM fungal communities among treatments were due to recruitment of relatively phylogenetically distant AM fungi, with *Rhi-zophagus* correlated with decreasing crop yield. These findings suggest that knowing how and when certain AM fungal phylotypes are favored could provide inroads towards better managing the symbiosis in agroecosystems. More experiments with a range of crops of different symbiotic groups, of different succeeding crops and with various farming regimes are now needed to expand the generality of our findings.

ACKNOWLEDGMENTS

This work was supported by the German Federal Ministry of Education and Research (BMBF) initiative 'BonaRes-Soil as a sustainable resource for the bioeconomy' through the project INPLAMINT (grant number: 031B0508B). E. F. Leifheit acknowledges funding from the Deutsche Forschungsgemeinschaft (LE 3859/1-1). We thank Max Fussan and Tobias Wiesner for help with measuring soil and root colonization. We thank Paul Gentil for help with root DNA extraction. We thank the DAAD AIESTE and RISE programmes for funding student helpers and Christoph Schmidt, Arthur César Coares, Douglas Henrique, Farida Samad-Zada, and Phong Hong in helping to set up, manage, and harvest the mesocosm experiment. Author contributions: R. van Duijnen, J. Roy, V. Temperton, and M. C. Rillig designed the experiment. R. van Duijnen performed the mesocosm experiment. R. van Duijnen and J. Roy performed soil and root sampling. J. Roy performed molecular analyses and analyzed data, M. C. Rillig and S. Mbedi contributed new reagents or analytical tools. J. Roy wrote the manuscript, which was revised by all authors.

LITERATURE CITED

- Anderson, M. J. 2001. A new method for non parametric multivariate analysis of variance. Austral Ecology 26:32–46.
- Angus, J. F., J. A. Kirkegaard, J. R. Hunt, M. H. Ryan, L. Ohlander, and M. B. Peoples. 2015. Break crops and rotations for wheat. Crop and Pasture Science 66:523–552.
- Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models using lme4. Journal of Statistical Software 67:1–48.
- Bennett, J. A., H. Maherali, K. O. Reinhart, Y. Lekberg, M. M. Hart, and J. Klironomos. 2017. Plant-soil feedbacks and mycorrhizal type influence temperate forest population dynamics. Science 355:181–184.
- Berruti, A., V. Bianciotto, and E. Lumini. 2018. Seasonal variation in winter wheat field soil arbuscular mycorrhizal fungus communities after non-mycorrhizal crop cultivation. Mycorrhiza 28:535–548.
- Bever, J. D. 2002. Reduces Plant Benefit Linked Negative feedback within a mutualism: host-specific growth of mycorrhizal fungi reduces plant benefit. Proceedings of The Royal Society of Biological Sciences 269:2595–2601.
- Bowles, T. M., L. E. Jackson, M. Loeher, and T. R. Cavagnaro. 2016. Ecological intensification and arbuscular mycorrhizas: a meta-analysis of tillage and cover crop effects. Journal of Applied Ecology 54:1785–1793.

- Brundrett, M. C., and L. Tedersoo. 2018. Evolutionary history of mycorrhizal symbioses and global host plant diversity. New Phytologist 220:1108–1115.
- Callahan, B. J., P. J. Mcmurdie, M. J. Rosen, A. W. Han, A. J. Johnson, and S. P. Holmes. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods 13:581.
- Campos, C., M. Carvalho, C. Brígido, M. J. Goss, T. Nobre, and M. C. Pagano. 2018. Symbiosis specificity of the preceding host plant can dominate but not obliterate the association between wheat and its arbuscular mycorrhizal fungal partners. Frontiers in Microbiology 9:1–14.
- Chagnon, P., R. L. Bradley, and J. N. Klironomos. 2015. Traitbased partner selection drives mycorrhizal network assembly. Oikos 124:1609–1616.
- Chagnon, P. L., R. L. Bradley, H. Maherali, and J. N. Klironomos. 2013. A trait-based framework to understand life history of mycorrhizal fungi. Trends in Plant Science 18:484– 491.
- Davison, J., et al. 2015. Global assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism. Science 127:970–973.
- Dray, S., and A. B. Dufour. 2007. The ade4 Package: implementing the duality diagram for ecologists. Journal of Statistical Software 22:1–20.
- Dumbrell, A. J., P. D. Ashton, N. Aziz, G. u. Feng, M. Nelson, C. Dytham, A. H. Fitter, and T. Helgason. 2011. Distinct seasonal assemblages of arbuscular mycorrhizal fungi revealed by massively parallel pyrosequencing. New Phytologist 190:794–804.
- Eck, J. L., S. M. Stump, C. S. Delavaux, S. A. Mangan, and L. S. Comita. 2019. Evidence of within-species specialization by soil microbes and the implications for plant community diversity. Proceedings of the National Academy of Sciences USA 116:7371–7376.
- Eo, J., and A. Eom. 2009. Differential growth response of various crop species to arbuscular mycorrhizal inoculation. Mycobiology 37:72–76.
- Frøslev, T. G., R. Kjøller, H. H. Bruun, R. Ejrnæs, A. K. Brunbjerg, C. Pietroni, and A. J. Hansen. 2017. Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. Nature Communications 8:1188.
- Groussin, M., F. Mazel, J. G. Sanders, C. S. Smillie, S. Lavergne, W. Thuiller, and E. J. Alm. 2017. Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. Nature Communications 8:1–12.
- Hart, M. M., and R. J. Reader. 2002. Taxonomic basis for variation in the colonization strategy of arbuscular mycorrhizal fungi. New Phytologist 153:335–344.
- Hart, M. M., R. J. Reader, and J. N. Klironomos. 2002. Lifehistory strategies of arbuscular mycorrhizal fungi in relation to their successional dynamics. Mycologia 93:1186–1194.
- Hoeksema, J. D., et al. 2010. A meta-analysis of contextdependency in plant response to inoculation with mycorrhizal fungi. Ecology Letters 13:394–407.
- Huang, L., L. Song, X. Xia, W.-H. Mao, K. Shi, Y.-H. Zhou, and J.-Q. Yu. 2013. Plant-soil feedbacks and soil sickness: from mechanisms to application in agriculture. Journal of Chemical Ecology 39:232–242.
- Jakobsen, I., L. K. Abbott, and A. D. Robson. 1992. External hyphae of vesicular-arbuscular mycorrhizal fungi associated with *Trifolium subterraneum* L. New Phytologist 120:371– 380.
- Johnson, N. C. 2010. Resource stoichiometry elucidates the structure and function of arbuscular mycorrhizas across scales. New Phytologist 185:631–647.

- Katoh, K., K. Misawa, K. Kuma, and T. Miyata. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research 30:3059–3066.
- Kirkegaard, J., O. Christen, J. Krupinsky, and D. Layzell. 2008. Break crop benefits in temperate wheat production. Field Crops Research 107:185–195.
- Koch, A. M., P. M. Antunes, H. Maherali, M. M. Hart, and J. N. Klironomos. 2017. Evolutionary asymmetry in the arbuscular mycorrhizal symbiosis: conservatism in fungal morphology does not predict host plant growth. New Phytologist 214:1330–1337.
- Koch, A. M., D. Croll, and I. R. Sanders. 2006. Genetic variability in a population of arbuscular mycorrhizal fungi causes variation in plant growth. Ecology Letters 9:103–110.
- Koziol, L., and J. D. Bever. 2019. Mycorrhizal feedbacks generate positive frequency dependence accelerating grassland succession. Journal of Ecology 107:622–632.
- Krüger, M., C. Krüger, C. Walker, H. Stockinger, and A. Schüßler. 2012. Phylogenetic reference data for systematics and phylotaxonomy of arbuscular mycorrhizal fungi from phylum to species level. New Phytologist 193:970–984.
- Krüger, M., H. Stockinger, C. Krüger, and A. Schüssler. 2009. DNA-based species level detection of *Glomeromycota*: one PCR primer set for all arbuscular mycorrhizal fungi. New phytologist 183:212–223.
- Legendre, P., and M. J. Anderson. 1999. Distance-based redundancy analysis: testing multispecies responses in multifactorial ecological experiments. Ecological Monographs 69:1–24.
- Leifheit, E. F., E. Verbruggen, and M. C. Rillig. 2015. Arbuscular mycorrhizal fungi reduce decomposition of woody plant litter while increasing soil aggregation. Soil Biology and Biochemistry 81:323–328.
- Lekberg, Y., S. M. Gibbons, and S. Rosendahl. 2014. Will different OTU delineation methods change interpretation of arbuscular mycorrhizal fungal community patterns? New Phytologist 202:1101–1104.
- Lekberg, Y., and R. T. Koide. 2005. Is plant performance limited by abundance of arbuscular mycorrhizal fungi? A metaanalysis of studies published between 1988 and 2003. New Phytologist 168:189–204.
- Lekberg, Y., and L. P. Waller. 2016. What drives differences in arbuscular mycorrhizal fungal communities among plant species? Fungal Ecology 24:135–138.
- López-García, A., S. Varela-cervero, M. Vasar, M. Öpik, J. M. Barea, and C. Azcón-Aguilar. 2017. Plant traits determine the phylogenetic structure of arbuscular mycorrhizal fungal communities. Molecular Ecology 26:6948–6959.
- Maherali, H., and J. N. Klironomos. 2007. Influence of phylogeny on fungal community assembly and ecosystem functioning. Science 316:1746–1748.
- Mariotte, P., Z. Mehrabi, T. M. Bezemer, D. G. B. De, A. Kulmatiski, B. Drigo, G. F. C. Veen, H. M. G. A. Van Der, and P. Kardol. 2018. Plant – soil feedback: bridging natural and agricultural sciences. Trends in Ecology & Evolution 33:129– 142.
- McGonigle, T., M. Millers, D. Evans, G. Fairchild, and J. Swan. 1990. A new method which gives an objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. New Phytologist 115:495–501.
- Mensah, J. A., A. M. Koch, P. M. Antunes, E. T. Kiers, M. Hart, and H. Bücking. 2015. High functional diversity within species of arbuscular mycorrhizal fungi is associated with differences in phosphate and nitrogen uptake and fungal phosphate metabolism. Mycorrhiza 25:533–546.

- Montero, H., J. Choi, and U. Paszkowski. 2019. Arbuscular mycorrhizal phenotyping: the dos and don'ts. New Phytologist 221:1182–1186.
- Mosse, B. 1959. Observations on the extra-matrical mycelium of a vesicular-arbuscular endophyte. Transactions of the British Mycological Society 42:439–448.
- Munkvold, L., R. Kjøller, M. Vestberg, S. Rosendahl, and I. Jakobsen. 2004. High functional diversity within species of arbuscular mycorrhizal fungi. New Phytologist 164:357–364.
- Nielsen, K., R. Kjøller, H. Bruun, T. Schnoor, and S. Rosendahl. 2016. Colonization of new land by arbuscular mycorrhizal fungi. Fungal Ecology 20:22–29.
- Oksanen, J. F., et al. 2016. vegan: Community Ecology Package. https://cran.r-project.org/package=vegan
- Phillips, J., and D. Hayman. 1970. Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. Transactions of the British Mycological Society 55:158–161.
- Powell, J. R., J. L. Parrent, M. M. Hart, J. N. Klironomos, M. C. Rillig, and H. Maherali. 2009. Phylogenetic trait conservatism and the evolution of functional trade-offs in arbuscular mycorrhizal fungi. Proceedings of The Royal Society of Biological Sciences 276:4237–4245.
- Powell, J. R., and B. A. Sikes. 2014. Method or madness: Does OTU delineation bias our perceptions of fungal ecology? New Phytologist 202:1095–1097.
- R Core Team. 2017. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/
- Rillig, M. C., et al. 2019. Why farmers should manage the arbuscular mycorrhizal symbiosis. New Phytologist 222:1–5.
- Rillig, M. C., S. F. Wright, and V. T. Eviner. 2002. The role of arbuscular mycorrhizal fungi and glomalin in soil aggregation: comparing effects of five plant species. Plant and Soil 238:325–333.
- Roger, A., A. Colard, C. Angelard, and I. R. Sanders. 2013. Relatedness among arbuscular mycorrhizal fungi drives plant growth and intraspecific fungal coexistence. ISME Journal 7:2137–2146.
- Roy, J., F. Mazel, A. Sosa-Hernandez, J. F. Duenas, S. Hempel, L. Zinger, and M. Rillig. 2019. The relative importance of ecological drivers of arbuscular mycorrhizal fungal distribution varies with taxon phylogenetic resolution. New Phytologist 224:936–948.
- Roy, J., R. Reichel, N. Brüggemann, S. Hempel, and M. C. Rillig. 2017. Succession of arbuscular mycorrhizal fungi along a 52-year agricultural recultivation chronosequence. FEMS Microbiology Ecology 93:1–13.
- Ryan, M. H., and J. H. Graham. 2018. Little evidence that farmers should consider abundance or diversity of arbuscular mycorrhizal fungi when managing crops. New Phytologist 220:1092–1107.
- Sanders, I. R., and A. Rodriguez. 2016. Aligning molecular studies of mycorrhizal fungal diversity with ecologically important levels of diversity in ecosystems. ISME Journal 10:2780–2786.
- Semchenko, M., S. Saar, and A. Lepik. 2017. Intraspecific genetic diversity modulates plant–soil feedback and nutrient cycling. New Phytologist 216:90–98.
- Smith, S. E., and D. Read. 2008. Mycorrhizal symbiosis. Academic Press, San Diego, California, USA.
- Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30:1312–1313.
- Teste, F. P., P. Kardol, B. L. Turner, D. A. Wardle, G. Zemunik, M. Renton, and E. Laliberté. 2017. Plant-soil feedback and

the maintenance of diversity in Mediterranean-climate shrublands. Science 355:173–176.

- Thompson, J. P. 1987. Decline of vesicular-arbuscular mycorrhizas in long fallow disorder of field crops and its expression in phosphorus deficiency in sunflower. Australian Journal of Agricultural Research 38:847–867.
- Torrecillas, E., M. M. Alguacil, and A. Roldán. 2012. Host preferences of arbuscular mycorrhizal fungi colonizing annual herbaceous plant species in semiarid Mediterranean prairies. Applied and Environmental Microbiology 78:6180–6186.
- Treseder, K. K. 2004. A meta-analysis of mycorrhizal responses to nitrogen, phosphorus, and atmospheric CO 2 in field studies. Ecology and Evolution 164:347–355.
- Treseder, K. K., M. M. Hart, E. B. A. L. M. Egertonwarburton, J. N. Klironomos, H. Maherali, and L. Tedersoo. 2018. Arbuscular mycorrhizal fungi as mediators of ecosystem responses to nitrogen deposition: A trait-based predictive framework. Journal of Ecology 106:480–489.
- Van Der Heijden, M. G. A., R. D. Bardgett, and N. M. Van Straalen. 2008. The unseen majority: Soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecology Letters 11:296–310.
- van der Heijden, M. G. A., J. N. Klironomos, M. Ursic, P. Moutoglis, R. Streitwolf-engel, T. Boller, A. Wiemken, and I. R. Sanders. 1998. Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. Nature 74:69–72.

- Van Der Putten, P. W. H., et al. 2013. Plant soil feedbacks: the past, the present and future challenges. Journal of Ecology 101:265–276.
- van Duijnen, R., J. Roy, W. Härdtle, and V. M. Temperton. 2018. Precrop functional group identity affects yield of winter barley but less so high carbon amendments in a mesocosm experiment. Frontiers in Plant Science 9:1–12.
- Vandenkoornhuyse, P., R. Husband, T. J. Daniell, I. J. Watson, M. Duck, A. H. Fitter, and J. P. W. Young. 2002. Arbuscular mycorrhizal community composition associated with two plant species in a grassland ecosystem. Molecular Ecology 11:1555–1564.
- Verbruggen, E., and T. E. Kiers. 2010. Evolutionary ecology of mycorrhizal functional diversity in agricultural systems. Evolutionary Applications 3:547–560.
- Yang, H., Q. Zhang, R. T. Koide, J. D. Hoeksema, J. Tang, X. Bian, S. Hu, and X. Chen. 2017. Taxonomic resolution is a determinant of biodiversity effects in arbuscular mycorrhizal fungal communities. Journal of Ecology 105:219–228.
- Yang, H., J. Zhou, M. Weih, Y. Li, S. Zhai, Q. Zhang, W. Chen, J. Liu, L. Liu, and S. Hu. 2020. Mycorrhizal nitrogen uptake of wheat is increased by earthworm activity only under no-till and straw removal conditions. Applied Soil Ecology 155:103672.
- Zhang, S., A. Lehmann, W. Zheng, Z. You, and M. C. Rillig. 2019. Arbuscular mycorrhizal fungi increase grain yields: a meta-analysis. New Phytologist 222:543–555.

SUPPORTING INFORMATION

Additional supporting information may be found online at: http://onlinelibrary.wiley.com/doi/10.1002/eap.2378/full

OPEN RESEARCH

Raw Illumina sequencing reads of LSU amplicons are available on the European Nucleotide Archive under accession no. PRJEB36419. Amplicon sequence variants obtained after bioinformatics are available on the European Nucleotide Archive under accession nos. LR761341 through LR761569. The sequence contingency matrix and the taxonomy of the sequences (https://doi.org/10.20387/bonares-sea4-n5sp), and the sample metadata (https://doi.org/10.20387/bonares-xnhz-xw6e) are freely available through the BONARES data portal.