

What shapes ground beetle assemblages in a tree species-rich subtropical forest?

Zumstein, Pascale; Bruelheide, Helge; Fichtner, Andreas; Schuldt, Andreas; Staab, Michael; Härdtle, Werner; Zhou, Hongzhang; Aßmann, Thorsten

Published in:
ZooKeys

DOI:
[10.3897/zookeys.1044.63803](https://doi.org/10.3897/zookeys.1044.63803)

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for pulished version (APA):
Zumstein, P., Bruelheide, H., Fichtner, A., Schuldt, A., Staab, M., Härdtle, W., Zhou, H., & Aßmann, T. (2021). What shapes ground beetle assemblages in a tree species-rich subtropical forest? *ZooKeys*, 2021(1044), 907-927. <https://doi.org/10.3897/zookeys.1044.63803>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

What shapes ground beetle assemblages in a tree species-rich subtropical forest?

Pascale Zumstein¹, Helge Bruelheide^{2,3}, Andreas Fichtner¹, Andreas Schuldt⁴,
Michael Staab⁵, Werner Härdtle¹, Hongzhang Zhou⁶, Thorsten Assmann¹

¹ Institute of Ecology, Leuphana University Lüneburg, Universitätsstraße 1, 21335, Lüneburg, Germany

² Institute of Biology/Geobotany, Martin Luther University Halle-Wittenberg, 06108 Halle (Saale), Germany

³ German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, 04103 Leipzig, Germany

⁴ Forest Nature Conservation, Georg-August-University Göttingen, Büsgenweg 3, 37077 Göttingen, Germany

⁵ Ecological Networks, Technical University Darmstadt, Schnitzpahnstraße 3, 64287 Darmstadt, Germany

⁶ Key Laboratory of Zoological Systematics and Evolution, Chinese Academy of Sciences, 1 Beichen West Road, Chaoyang District, Beijing 100101, China

Corresponding author: Pascale Zumstein (zumstein@leuphana.de)

Academic editor: John Spence | Received 31 January 2021 | Accepted 8 April 2021 | Published 16 June 2021

<http://zoobank.org/A8ED7174-2412-46B4-9A5D-0345ADE42D07>

Citation: Zumstein P, Bruelheide H, Fichtner A, Schuldt A, Staab M, Härdtle W, Zhou H, Assmann T (2021) What shapes ground beetle assemblages in a tree species-rich subtropical forest? In: Spence J, Casale A, Assmann T, Liebherr JK, Penev L (Eds) Systematic Zoology and Biodiversity Science: A tribute to Terry Erwin (1940–2020). ZooKeys 1044: 907–927. <https://doi.org/10.3897/zookeys.1044.63803>

Abstract

As woody plants provide much of the trophic basis for food webs in forests their species richness, but also stand age and numerous further variables such as vegetation structure, soil properties and elevation can shape assemblages of ground beetles (Coleoptera: Carabidae). However, the combined impact of these numerous variables on ground beetle diversity and community structure has rarely been studied simultaneously. Therefore, ground beetles were studied in 27 plots in a highly diverse and structurally heterogeneous subtropical forest ecosystem, the Gutianshan National Park (southeast China) using pitfall traps and flight interception traps. Both trapping methods collected partly overlapping species spectra. The arboreal fauna was dominated by lebiines and to a smaller extent by tiger beetles and platynines; the epigeic fauna comprised mostly representatives of the genus *Carabus* and numerous tribes, especially anisodactylines, pterostichines, and sphodrines. Ground beetle species richness, abundance, and biomass of the pitfall trap catches were analyzed with generalized linear mixed models (GLMMs), fitted with seven environmental variables. Four of these variables influenced the ground beetle assemblages: Canopy cover, herb cover, pH-value of the topsoil and elevation. Contrary to our expectations, woody plant species richness and stand age did not significantly affect ground beetle assemblages. Thus, ground beetles seem to respond differently

to environmental variables than ants and spiders, two other predominantly predatory arthropod groups that were studied on the same plots in our study area and which showed distinct relationships with woody plant richness. Our results highlight the need to study a wider range of taxa to achieve a better understanding of how environmental changes affect species assemblages and their functioning in forest ecosystems.

Keywords

Abundance, BEF-China, biomass, canopy cover, Carabidae, elevational gradient, herb cover, pH-value, species richness

Introduction

Tropical and subtropical forests are among the world's ecosystems with the highest biodiversity. A substantial part of this biodiversity consists of arthropods, which are involved in many ecosystem processes (Basset et al. 2012; Schuldt et al. 2015b; Stork 2018). Numerous biotic and abiotic variables can influence their diversity, but also the abundance and biomass of these arthropods (Rainio and Niemela 2003; Mottl et al. 2020; Marrec et al. 2021). Trees structure forest ecosystems and, as they are the most important primary producers of biomass, they provide the basis for much of the food web consisting of decomposers, herbivores, and predators. As many biodiversity patterns in forest ecosystems seem to be influenced by bottom-up processes (Gruner 2004; Wu et al. 2013; Korboulewsky et al. 2016), woody plant species richness can have a strong impact on higher trophic levels (Schuldt et al. 2017). As a consequence, the diversity of herbivores may be strongly determined by the diversity of trees (Riihimäki et al. 2005; Wang et al. 2019). Also at higher trophic levels (predators, parasitoids), there seem to be bottom-up effects from woody plant diversity (Strong et al. 1984; Lefcheck et al. 2015; Schuldt et al. 2015a). However, such relationships between plant diversity and arthropods are not always present or apparent (Vehviläinen et al. 2008).

To analyze the effects of woody plant species richness on arthropod diversity in a real-world forest ecosystem, it is important to consider that environmental variables can also influence communities or assemblages of arthropods (Thiele 1977; Bonn and Schröder 2001). In experiments with manipulated woody plant species richness, possible biodiversity effects can be better determined (Loreau et al. 2001; Duffy 2009); however, conclusions from such manipulative experimental approaches cannot simply be transferred to forests with long-lasting maturation processes, in particular because experiments are usually not much older than a few decades. Therefore, real-world ecosystems are of particular importance for the study of arthropod communities (or assemblages) in forests of different stand ages.

Ground beetles (Coleoptera: Carabidae) represent a highly diverse predatory taxon and are typical of forest ecosystems across latitudinal gradients (Zou et al. 2013; Nolte et al. 2017; Garcia-Tejero et al. 2018). Moreover, ground beetles in temperate and boreal regions are known to be sensitive to numerous edaphic variables (e.g., pH-value: Paje and Mossakowski 1984; moisture: Antvogel and Bonn 2001). Even effects of

land-use legacy, tree age, vegetation structure and succession are known (Terrell-Nield 1990; Assmann 1999; Jelaska et al. 2011; Marrec et al. 2021). In general, ground beetles are considered as a well-known taxon for bio-indication since numerous species react sensitively to small changes in environmental variables (Koivula 2011).

The shaping variables of ground beetle assemblages in global biodiversity forest hotspots are poorly understood. In particular, the ground beetles of the southeast Chinese subtropics with summer rainfalls, which host particularly species-rich forests (Liu et al. 2003; Tang 2006), provide an interesting opportunity for study. We worked in Gutianshan National Park (Zhejiang, southeast China) that covers an area with remarkable subtropical tree and shrub species diversity. Relationships between biodiversity and ecosystem functions and the effects of environmental variables have been studied extensively in this area (e.g., Schuldt et al. 2010; Lang et al. 2012; Staab et al. 2014; Eichenberg et al. 2015; Zou et al. 2015; O'Brien et al. 2016; Brezzi et al. 2017). However, while the main abiotic and biotic drivers potentially shaping herbivore, decomposer and predator diversities have been studied (e.g., Schuldt et al. 2011; Staab et al. 2014; Schuldt et al. 2015a; Binkenstein et al. 2018; Staab et al. 2021) studies on ground beetles are still lacking for this National Park.

A study of ground beetles would reveal not only how this arthropod taxon responds to environmental variables, but the comparison with other important taxa (esp. spiders and ants) could provide a better understanding of how environmental variables affect the abundance and species diversity of predatory arthropods. Thus, since environmental variables can potentially influence ground beetle assemblages, we hypothesize that abundance, richness, and biomass of ground beetles increases with (H1) woody plant species richness and (H2) stand age in a subtropical forest in China. Moreover, we expect that (H3) the structural richness of vegetation and abiotic variables affect ground beetle abundance, richness, and biomass. Specifically, we investigate whether woody plant species richness, stand age of forests plots, structural richness of vegetation (cover of canopy and herb layer), and abiotic variables (elevation and soil pH) influence abundance, species richness, and biomass of ground beetles.

Material and methods

Study area and plots

Gutianshan National Park (Gutianshan NP), formerly Gutianshan National Nature Reserve, is located in the western part of Zhejiang Province in southeast China (29°14'N, 118°07'E). The park is approximately 81 square kilometers in size and was established in 1975, first as a National Forest Reserve, to preserve parts of the old-growth evergreen broad-leaved forest in the region. The climate is typical for subtropical areas with an annual mean temperature of 15.3 °C and ~2000 mm mean precipitation per year, occurring mostly between March and September (Yu et al. 2001). Further meteorological information is provided by Bruelheide et al. (2011) or cited therein. The park

is characterized by a mixed broad-leaved forest composed of deciduous and evergreen tree species, and in which *Castanopsis eyrei* (Champion & Bentham, 1845) and *Schima superba* (Gardner & Champion, 1849) are the prevailing tree species (Hu and Yu 2008; Legendre et al. 2009). Most stands consist of secondary forest, with maximum ages of trees ≤ 180 years. The park is located in a mountain range with elevations varying between 300 and 1260 m above sea level. Our study focused on stands ranging from 251 to 903 m a.s.l. The local rock is granitic and thus the topsoil pH-values ranges from 5.5 to 6.5 (Bruelheide et al. 2011).

In 2008, 27 study plots, each measuring 30 m \times 30 m, were established by a research consortium of Chinese and European scientists (Bruelheide et al. 2011). The plots were randomly distributed throughout the park, excluding sites with slopes $> 55^\circ$. The plots included gradients of woody plant species richness (25–69 trees and shrub species per plot) and stand age (21–115 years). Further information on plot characteristics and study design is provided by Bruelheide et al. (2011).

Pitfall trapping

We sampled ground beetles using four pitfall traps in each plot, installed in the corners of the central 10 m \times 10 m square (resulting in 108 traps). Traps were plastic cups with a diameter of 8.5 cm and a capacity of ~ 550 ml filled with 150 ml of preserving solution (40% ethanol, 20% glycerol, 10% acetic acid, 30% water). They were open continuously during the main vegetation period in 2009 (end of March to the beginning of September). Traps were emptied fortnightly (in total ten collections per trap) and catches were preserved in 70% ethanol until identification (Schuldt et al. 2011). Flight interception traps were installed close to the pitfall traps in 2010 (Schuldt et al. 2015a; Staab et al. 2021). Ground beetles were sorted and body length was measured for biomass calculation (see section *Statistical analysis*). Specimens were identified to species level or classified to morphospecies by carabid taxonomists (David W. Wrase, Thierry Deuve, and Thorsten Assmann). Results from such a “morphospecies approach” show high correlation to true species diversity (Oliver and Beattie 1996). Hereafter, we use the term species for both diagnosed species and morphospecies to characterize our catches.

Environmental data

During plot establishment, a comprehensive set of environmental variables of biotic and abiotic habitat characteristics was collected (Bruelheide et al. 2011). We asked if seven variables, known to influence ground beetles assemblages (e.g., Assmann 1999; Antvogel and Bonn 2001; Marrec et al. 2021) could predict the abundance, species richness and biomass of ground beetles. The age of the forest included in each plot was estimated using stem core measurements of the tree with the fifth largest diameter at breast height within each plot. Woody plant species richness was determined as the number of all tree and shrub species represented by individuals larger than 1 m in the

plot (Bruelheide et al. 2011). Species richness and abundance of the herbaceous layer were estimated for all herb species smaller than 1 m height in the inner 10 m × 10 m subplots (Both et al. 2011). Canopy cover (%), herb cover (%), elevation (meters a.s.l.), and pH-value of the topsoil (0–5 cm; taken from nine dried and sieved soil samples per plot, pooled and measured potentiometrically) were additionally assessed as plot characteristics (see Bruelheide et al. (2011) for details).

Statistical analysis

All statistical analyses were conducted using the packages *glmmTMB* (Brooks et al. 2017) and *DHARMa* (Hartig 2020) in R 3.6.3 (<http://www.R-project.org>). We applied generalized linear mixed models (GLMMs) to assess the effects of biotic (woody species richness, stand age, canopy and herb cover, herb species richness) and abiotic (topsoil pH-value, elevation) stand conditions (i.e., plot characteristics) on the abundance, species richness and biomass of ground beetles. Plot identity was used as a random factor to account for the nested data structure (traps nested in study plots). Abundance and species richness data were modelled with a Poisson distribution and log-link function. There was no indication of overdispersion in the data for either abundance ($p = 0.25$) or species richness ($p = 0.20$). For the biomass data, we applied a Gamma distribution with a log-link and added a value of 0.01 to the observed biomass prior to model fitting to ensure the convergence of the algorithm. Ground beetle biomass was calculated according to the following formula:

$$\ln y = -8.92804283 + 2.5554921 \ln x,$$

where: x is the measured body length of the specimen and y is the estimated body weight of the individual (Schwerk and Szyszko 2007). The calculated biomass estimates were then summed over all individuals per plot.

Model selection was based on likelihood-ratio tests starting with a fully saturated model that included all predictors and the interaction between woody species richness and stand age to test, if possible, species richness effects depended on the age of the forest plots. We sequentially removed non-significant ($p > 0.05$) terms and tested for assumptions of the best-fitting model following (Zuur et al. 2009). Prior to analysis, all predictor variables were standardized (mean = 0, SD = 1) and tested for critical correlations (all variance inflation factors were < 1.75).

Results

Number of species, abundance, and biomass

In total, we caught 258 ground beetle specimens in the pitfall traps, belonging to 22 species (Table 1). Catches ranged from two to 27 specimens per plot (mean

Table 1. Collected ground beetles from pitfall traps of 27 plots in Gutianshan NP. For classification, we followed the systematics of the Palearctic catalogue (Löbl and Löbl 2017). Abundance (No.) and body size (mean body size if more than one individual caught) is given for each (morpho-) species. The elevation data refer to the highest and lowest plots where the ground beetles were caught.

Tribe	(Morpho-) Species	No.	Size (mm)	Elevation (m a.s.l.)
Anisodactylini	undet. spec. 1	60	8.9	348–903
	undet. spec. 2	3	8	639
Carabini	<i>Carabus (Apotomopterus) davidis</i> Deyrolle, 1878	5	35	566–679
	<i>Carabus (Isiocarabus) kiukiangensis</i> Bates, 1888	20	30	348–903
	<i>Carabus (Damaster) lafossei</i> Feisthamel, 1845	5	42	566–679
Brachinini	<i>Pheropsochus (Stenaptinus) beckeri</i> Jedlicka, 1930	1	14	647
Harpalini	<i>Amara</i> spec. 1	1	8	542
	<i>Harpalus</i> spec. 1	1	12	617
Lebiini	<i>Calleida (Callidiula)</i> spec. 1	1	12	617
	<i>Lachnoderma asperum</i> Bates, 1883	1	8	880
	Pericalina, undet. 1	1	5	617
Pentagonici	<i>Pentagonica</i> spec. 1	10	4.5	251–679
	<i>Pentagonica</i> spec. 2	1	5	542
Perigonini	<i>Perigona</i> spec. 1	4	3	542–720
Pterostichini	<i>Lesticus</i> spec. 1	7	25.3	590–903
	<i>Lesticus</i> spec. 2	43	28.1	251–903
	<i>Prerostichus</i> spec. 1	47	24.8	251–903
	<i>Prerostichus</i> spec. 2	3	11	419–670
Sphodrini	<i>Synuchus</i> spec. 1	32	13.7	251–679
	<i>Synuchus</i> spec. 2	10	10.9	251–903
Cicindelini	<i>Cylindera (Ifasina) kaleea</i> Bates, 1863	1	9	880
Collyridini	<i>Tricondyla macrodera</i> Chaudoir, 1861	1	19	566

Table 2. Collected ground beetles from flight interception traps of 27 plots in Gutianshan NP. For classification, we followed Löbl and Löbl (2017). Abundance (No.) and body size (mean body size if more than one individual caught) given for each (morpho-) species.

Tribe	(Morpho-) Species	No.	Size (mm)
Lebiini	<i>Lachnoderma asperum</i> Bates, 1883	1	8
	<i>Lioptrera erotyloides</i> Bates, 1883	1	13
	<i>Coptodera (Coptoderina)</i> spec. 1	42	7.5
Pentagonici	<i>Pentagonica</i> spec. 1	2	4.5
Platynini	undet. spec. 1	1	8
Collyridini	<i>Tricondyla macrodera</i> Chaudoir, 1861	2	19

9.6 ± 1.19 SE), based on a relatively low number of individuals (2.4 ± 0.22) per trap. As expected, species richness was strongly correlated with abundance (Pearson correlation coefficient $r = 0.85$, $p \leq 0.001$). Biomass of captured ground beetles ranged from 0.15 to 10.63 g per plot. The average biomass per plot was 3.11 g (± 0.54 SE).

The number of ground beetle specimens in flight interception traps was lower than in pitfall traps. In total, we caught 49 individuals of six species (Table 2), three of which were recorded also by the pitfall traps on the forest floor. As we caught few ground beetles in the flight interception traps, we performed further analyses only with the catches of the pitfall traps. Although there was overlap in species spectra, the catches of the two trapping methods differed strongly. Species of the genus

Table 3. Results from mixed-effects models for ground beetle abundance, species richness, and biomass. P-values were obtained from likelihood-ratio tests starting with a full-saturated model and removing non-significant ($p > 0.05$) terms sequentially. Significant predictors ($p < 0.05$) are indicated in bold.

	Abundance		Species richness		Biomass	
	χ^2	p-value	χ^2	p-value	χ^2	p-value
Woody species richness (WSR)	0.07	0.787	0.04	0.849	1.44	0.230
Stand age	0.33	0.565	0.00	0.961	1.17	0.280
Canopy cover	4.28	0.039	3.90	0.048	4.98	0.026
Herb cover	4.67	0.031	5.98	0.014	0.60	0.438
Herb species richness	1.60	0.206	0.89	0.345	0.18	0.673
pH-value (topsoil)	5.30	0.021	3.64	0.056	0.97	0.324
Elevation	0.39	0.531	0.18	0.668	14.71	<0.001
WSR * stand age	0.01	0.941	0.19	0.664	2.41	0.120




Figure 1. Relationships between ground beetle abundance and canopy cover (A), herb cover (B) and pH-value of the soil (C). Black lines indicate significant relationships at $p < 0.05$ obtained from mixed-effects models (keeping other significant predictors fixed at their means) with grey areas indicating the 95% confidence intervals. Points represent observed values per trap. Note that some traps had similar abundance and predictor values. The fixed-effects explained 22% of the variation in ground beetle abundance.

Carabus and the tribes Anisodactylini and Pterostichini were caught exclusively in the pitfall traps; and lebiines were more abundant in the flight interception traps (Tables 1, 2; Fig. 4).

Ground beetle assemblages and environmental variables

Four of the seven environmental variables tested for effects on ground beetles in generalized linear mixed models were significantly related to the ground beetle assemblages. Canopy cover had a positive influence on species richness, abundance, and biomass of ground beetles (Table 3, Figs 1A, 2A, 3A); however, herb cover influenced ground beetle species richness and abundance negatively (Table 3, Figs 1B, 2B). In addition, ground beetle abundance decreased significantly with increasing soil pH (Table 3, Fig. 1C). Finally, ground beetle biomass significantly increased with elevation (Table 3, Fig. 3B). This is related to the presence of *Carabus* species, the ground beetles with the largest body lengths in our study, especially at higher elevations. The biomass of repre-




Figure 2. Relationships between ground beetle species richness and canopy cover (A) and herb cover (B). Black lines indicate significant relationships at $p < 0.05$ obtained from mixed-effects models (keeping other significant predictors fixed at their means) with grey areas indicating the 95% confidence intervals. Points represent observed values per trap. Note that some traps had similar richness and predictor values. The fixed-effects explained 12% of the variation in ground beetle species richness.




Figure 3. Relationships between ground beetle biomass and canopy cover (A) and herb cover (B). Black lines indicate significant relationships at $p < 0.05$ obtained from mixed-effects models (keeping other significant predictors fixed at their means) with grey areas indicating the 95% confidence intervals. Points (slightly jittered to improve visibility) represent observed values per trap. The fixed-effects explained 30% of the variation in ground beetle biomass.

representatives of the genus *Carabus* was strongly correlated with elevation (Pearson correlation coefficient $r = 0.65$, $p \leq 0.001$). The biomass of the representatives of the tribes Pterostichini, which included species with the second largest individuals, however, was




Figure 4. Representatives of ground beetles from pitfall traps and flight interception traps in Gutianshan NP **A** *Carabus kiukiangensis* **B** *Carabus davidis* **C** *Lioptera erotyloides* **D** *Tricondyla macrodera*.

not significantly correlated with elevation ($r = 0.26$, $p > 0.05$). The same was true for the Anisodactylini ($r = 0.005$, $p > 0.05$) which were numerous in terms of individuals ($n = 63$). Neither species richness of woody plants and of herbs nor stand age significantly influenced any characteristic of the ground beetle assemblages (Table 3).

Discussion

Our study revealed that four environmental variables impacted on ground beetle assemblages at our sample sites. Contrary to our expectations, woody plant species richness (H1) and stand age (H2) did not influence ground beetle assemblages in this study. We did corroborate H3, as higher canopy cover led to increased species richness, abundance, and biomass of ground beetles, and because ground beetle abundance and species richness decreased with higher herb cover. Moreover, soil pH negatively influenced ground beetle abundance, and greater biomass of beetles was found at higher elevations.

Vegetation effects and ground beetle assemblages

Classical ecological theory such as the “enemies” hypothesis (Root 1973; Staab and Schuldt 2020), predict higher predator abundance and diversity with increasing plant diversity caused by different mechanisms at herbivore and predator trophic levels. This theory has been corroborated in other studies, especially in non-forest ecosystems (Andow 1991; Haddad et al. 2009). Furthermore, Jouveau et al. (2020) demonstrated positive and additive effects of vegetation diversity (understory, canopy and surrounding scales) on the density of ground beetles in a tree diversity experiment with planted tree individuals. Nonetheless, with increasing diversity of an ecosystem, mixed results are often obtained. For example, some studies postulate positive relationships between plant diversity and ground beetles, while other studies show no or negative relationships (Vehvilainen et al. 2008; Worthen and Merriman 2013; Yeeles et al. 2017; Zou et al. 2019).

We found no relationship between woody plant species richness and species numbers, abundance, or biomass of ground beetles in Gutianshan NP. This result is consistent with those from numerous other studies on plants and arthropods, especially ground beetles (Wolters et al. 2006; Harry et al. 2019; Corcos et al. 2021). Ground-dwelling predators do not directly depend on vegetation, while most herbivores depend directly on host plant selection. This difference alone may result in different patterns for the two trophic groups. It is primarily the structural features of vegetation that increase with increasing woody plant species diversity (Bruehlheide et al. 2011; Schuldt et al. 2019) that would be expected to affect predator assemblages. Such structural patterns, in turn, influence microclimatic factors such as temperature, humidity, and light availability and thus the activity and distribution of species such as carabid beetles (Work et al. 2011). Nevertheless, the non-significant results of the mixed-effects model

provide no support for relationships between woody plant species richness and ground beetle assemblages in Gutianshan NP. However, previous studies of epigeic arthropods in our study plots found mixed evidence of relationships with woody plant species richness. The relationship for spiders was negative but positive for predatory ants, with no relationship for omnivorous ants (Schuldt et al. 2011; Staab et al. 2014).

The positive relationship between woody plant and ground beetle diversity found by Zou et al. (2019) is limited to mature forests in a study area in temperate China. For secondary forests, no correlation was found, which the authors attributed to the lower forest age. The forest in Gutianshan NP is classified as secondary forest, because it was previously used agriculturally (Bruehlheide et al. 2011). Compared to the forests studied by Zou et al. (2019), the older plots in Gutianshan NP closely resemble mature stands. However, we found no relationship between ground beetle richness and woody plant species diversity in older stands in Gutianshan NP. Possibly, these forest plots are not old enough to have fostered a reasonably distinct natural ground beetle community.

Greater closure of the canopy layer was associated with more beetle species and specimens. However, our results contrast findings from numerous studies in forests of boreal, temperate and Mediterranean climate zones, according to which the number of ground beetle species increases with decreasing canopy cover (Koivula et al. 2002; Taboada et al. 2008; Thorn et al. 2016). These findings can be explained (at least partly) by the presence of more open habitat species of ground beetles in addition to the forest species in open or structural-rich forest sites (Heliölä et al. 2001; Magura et al. 2001). Nonetheless, if open habitat specialists are rare in the study region and simultaneously forest specialists avoid more open patches in Gutianshan NP, beetle catches would be lower in plots with less canopy cover. Although the hypothesis is tempting, habitat preferences of ground beetles in Chinese forests and other habitats are not as well studied as in the western Palearctic (Yu et al. 2009; Zou et al. 2015) and no data are available to test this idea.

In contrast, herb cover had a negative influence on ground beetle abundance and species richness. Studies from forests in both temperate and Mediterranean climate regions have documented both negative and positive influences of the forest herb layer on the species richness of ground beetles (Antvogel and Bonn 2001; Taboada et al. 2010; Liu et al. 2016). It is known that the cover of the herb layer can impede the movements of ground beetles on the forest floor (Taboada et al. 2010). Thus, a denser understory could hinder ground beetle activity within these stands. This, in turn, could negatively affect hunting processes in such stands. It might also reduce estimates of abundance and diversity that depend on activity-based trapping as in our study which uses pitfall traps. Contrastingly, a greater herb cover provides protection from potential predators. Additionally, the forest understory can modify the microclimate (temperature, humidity, sun-exposition), which is known to shape the ecological niches of many ground beetles (Thiele 1977; Bonn and Schröder 2001). Nonetheless, the influence of understory vegetation could be more complex and therefore requires further research with regard to ground beetles.

Abiotic effects and ground beetle assemblages

The guild of predatory arthropods (Araneae, Chilopoda, Formicidae, cavity-nesting wasps and their parasitoids) in Gutianshan NP showed a significant decrease of both abundance and species richness with increasing elevation (Binkenstein et al. 2018). In contrast, we found that the biomass of the predominately predatory ground beetles studied on the same plots increased along the same elevational gradient. Therefore, response of organisms to elevation depends not only on trophic guild (predators, herbivores, decomposers) in these forests (Binkenstein et al. 2018), but it can also vary within trophic guilds.

Our results for biomass indicate that beetle biomass within plots increases with elevation. This relationship is driven by the higher numbers of representatives of the genus *Carabus* (Fig. 4A, B), with specimens up to 42 mm in body length, the largest ground beetles in our study. The genus *Carabus* does not occur in the tropics and only a few species occur in the summer humid subtropics (such as in Gutianshan NP). The centers of species diversity are clearly in the temperate climatic zones (Breuning 1932; Meyer 2008). The general climate preference of *Carabus* species may therefore be reflected in our catches, where we found more individuals of these species at the cooler (and moister) higher elevations.

Ground beetles are able to adapt to a wide range of varying pH-values (Krogerus 1939, 1960; Paje and Mossakowski 1984); however, our data showed significantly negative relationship between ground beetle abundance and increasing topsoil pH. Evidence from field studies suggests that different ground beetle species prefer different pH-values in the soil (Mossakowski 1970; Matern et al. 2007). Explanations have been suggested only for a very limited number of species (e.g., species adapted to feed on snails, which prefer calcareous soils with higher pH-values: Assmann 2003). To our knowledge, no such studies are available for ground beetles from China. However, the pH-gradient we found in Gutianshan NP is relatively compressed, and also indirect effects, e.g., via plants, could influence ground beetle assemblages.

Low species number, abundance, and biomass of ground beetles

The overall number of ground beetle specimens, species and biomass was relatively low in Gutianshan NP. Given the fact, that our study took place in a subtropical hot-spot of vascular plant diversity, we expected higher numbers of ground beetles. For example, studies using the same sampling approach in Central European forests have reported 60 times more ground beetle specimens and 90 times more ground beetle biomass (Homburg et al. 2019; Hülsmann et al. 2019). Similarly, the number of species from temperate forests is higher (Nolte et al. 2017; Marrec et al. 2021). Further south, towards the European subtropical zone, Brandmayr et al. (1983) found similar low catches of both individuals and species, but in a less plant diverse forest (Mediterranean forest dominated by *Quercus ilex* (Linnaeus, 1753)). Comparable subtropical forest studies in China are largely lacking, but low catch rates of ground beetles have also been reported from another study conducted in Gutianshan NP (Yu et al. 2017).

Pitfall trap catches for carabids seem to be low in most tropical evergreen forests (Vennila 1999, 2000; Maveety et al. 2011; Qodri et al. 2016). In comparison to Gutianshan NP, the dry tropics of Colombia host more species and greater abundance (Ariza et al. 2021). Many of these tropical species occur mostly in the canopy. Terry Erwin was the pioneer of research on insect diversity of this forest stratum (Erwin 1982a, b, 2002), discovering many new species in the canopy of South American rainforests (Arndt et al. 2001; Erwin 2004a, b). However, we know from other studies that the ground strata can contribute to the overall biodiversity similar to that of higher strata, such as canopies (Stork and Grimbacher 2006).

Catches from flight interception traps

Although only a few species were recorded in our flight interception traps, they represent guilds or taxonomic entities that are well known from the tropics (Erwin and Erwin 1976; Erwin 1982b, 1983; Maveety et al. 2011), illustrating that the fauna of the subtropical Gutianshan NP is composed of temperate and tropical elements. At least some of these species have morphological adaptations to an arboreal life like pectinate claws and adhesive setae on the lower side of the tarsi (Erwin 1979; Stork 1980). All species from our flight interception traps show these morphological adaptations: lebiine species with their ectoparasitic larval development (e.g., *Lioptera erotyloides* (Fig. 4C), which mimics erotylids in coloration, its possible hosts which also occur in Gutianshan NP (cf. Erwin and Erwin 1976; Hunting and Yang 2019)), the collyridine *Tricondyla macrodera* (Fig. 4D), which not only resembles ants in their habitus, but also regularly hunts them (Shook and Wu 2007; own observations), and the platynine species (cf. Whitehead and Ball 1997).

The epigeic assemblages with their few species and individuals, but also the arboreal fauna with its specific morphological adaptations in Gutianshan NP resembles the ground beetle fauna typical for tropical forests. However, further study is required to achieve a better understanding of patterns of ground beetle species richness in subtropical forests.

Acknowledgements

We dedicate this work to the memory of the late Terry Erwin (1940–2020). He was one of the pioneers who studied how tree species diversity can affect the diversity of insects, especially ground beetles. One of his publications from a rainforest in Panama stimulated intensive research on tropical species diversity, particularly of arthropods. At the same time, Terry was one of the most dedicated entomologists we have ever met. His ecological work, as well as his contributions to natural history and conservation biology of carabids, has been very stimulating to our own studies. We are grateful for his influence and we will not forget him and his work.

We thank the administration of the Gutianshan National Park, and all BEF-China members for their support. We further thank Jiang Zaigen (Gutianshan) for help with

trap maintenance, Marianne Peters (Leuphana University Lüneburg) for sorting trap catches, David D. Wrase (Gusow-Platkow), and Thierry Deuve (Natural History Museum Paris) for help with identification. We gratefully acknowledge funding by the German Research Foundation (DFG FOR891/1-3).

We also thank John Spence, Nigel Stork, and an anonymous reviewer for useful comments that helped improve this work.

References

- Andow DA (1991) Vegetational diversity and arthropod population response. *Annual Review of Entomology* 36: 561–586. <https://doi.org/10.1146/annurev.ento.36.1.561>
- Antvogel H, Bonn A (2001) Environmental parameters and microspatial distribution of insects: a case study of carabids in an alluvial forest. *Ecography* 24: 470–482. <https://doi.org/10.1034/j.1600-0587.2001.d01-203.x>
- Ariza GM, Jácome J, Esquivel HE, Kotze DJ (2021) Early successional dynamics of ground beetles (Coleoptera, Carabidae) in the tropical dry forest ecosystem in Colombia. In: Spence J, Casale A, Assmann T, Liebherr JK, Penev L (Eds) *Systematic Zoology and Biodiversity Science: A tribute to Terry Erwin (1940–2020)*. *ZooKeys* 1044: 877–906. <https://doi.org/10.3897/zookeys.1044.59475>
- Arndt E, Kirmse S, Erwin TL (2001) Arboreal beetles of Neotropical forests: *Agra* Fabricius, larval descriptions with notes on natural history and behaviour (Coleoptera: Carabidae, Lebiini, Agrina). *Coleopterists Bulletin* 55: 297–310. [https://doi.org/10.1649/0010-065X\(2001\)055\[0297:ABONFA\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2001)055[0297:ABONFA]2.0.CO;2)
- Assmann T (1999) The ground beetle fauna of ancient and recent woodlands in the lowlands of north-west Germany (Coleoptera: Carabidae). *Biodiversity and Conservation* 8: 1499–1517. <https://doi.org/10.1023/A:1008974413376>
- Assmann T (2003) Biology and ecology. In: Turin H, Penev L, Casale A (Eds) *The Genus Carabus in Europe: a Synthesis*. Pensoft, Sofia-Moscow, 287–305.
- Basset Y, Cizek L, Cuenoud P, Didham RK, Guilhaumon F, Missa O, Novotny V, Odegaard F, Roslin T, Schmidl J, Tishechkin AK, Winchester NN, Roubik DW, Aberlenc HP, Bail J, Barrios H, Bridle JR, Castano-Meneses G, Corbara B, Curletti G, da Rocha WD, de Bakker D, Delabie JHC, Dejean A, Fagan LL, Floren A, Kitching RL, Medianero E, Miller SE, de Oliveira EG, Orivel J, Pollet M, Rapp M, Ribeiro SP, Roisin Y, Schmidt JB, Sorensen L, Leponce M (2012) Arthropod diversity in a tropical forest. *Science* 338: 1481–1484. <https://doi.org/10.1126/science.1226727>
- Binkenstein J, Klein AM, Assmann T, Buscot F, Erfmeier A, Ma K, Pietsch KA, Schmidt K, Scholten T, Wubet T, Bruelheide H, Schuldt A, Staab M (2018) Multi-trophic guilds respond differently to changing elevation in a subtropical forest. *Ecography* 41: 1013–1023. <https://doi.org/10.1111/ecog.03086>
- Bonn A, Schröder B (2001) Habitat models and their transfer for single and multi-species groups: a case study of carabids in an alluvial forest. *Ecography* 24: 483–496. <https://doi.org/10.1034/j.1600-0587.2001.d01-204.x>

- Both S, Fang T, Böhnke M, Bruelheide H, Geissler C, Kühn P, Scholten T, Trogisch S, Erfmeier A (2011) Lack of tree layer control on herb layer characteristics in a subtropical forest, China. *Journal of Vegetation Science* 22: 1120–1131. <https://doi.org/10.1111/j.1654-1103.2011.01324.x>
- Brandmayr P, Colombetta G, Polli S (1983) Waldcarabiden des Triester Karstes als Indikatoren des makroklimatischen Überganges vom kontinentalen Europa zur Mediterraneis (Coleoptera: Carabidae). *Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere* 110: 201–220.
- Breuning S (1932) Monographie der Gattung *Carabus*, L. Bestimmungs-Tabellen der europäischen Coleopteren. 104th–110th edn. Troppau, 1016 pp.
- Brezzi M, Schmid B, Niklaus PA, Schuldt A (2017) Tree diversity increases levels of herbivore damage in a subtropical forest canopy: evidence for dietary mixing by arthropods? *Journal of Plant Ecology* 10: 13–27. <https://doi.org/10.1093/jpe/rtw038>
- Brooks M, Kristensen K, van Benthem K, Magnusson A, Berg CW, Nielsen A, Skaug H, Mächler M, Bolker B (2017) glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *R Journal* 9: 378–400. <https://doi.org/10.32614/RJ-2017-066>
- Bruelheide H, Böhnke M, Both S, Fang T, Assmann T, Baruffol M, Bauhus J, Buscot F, Chen XY, Ding BY, Durka W, Erfmeier A, Fischer M, Geissler C, Guo D, Guo LD, Härdtle W, He JS, Hector A, Kröber W, Kühn P, Lang AC, Nadrowski K, Pei K, Scherer-Lorenzen M, Shi X, Scholten T, Schuldt A, Trogisch S, von Oheimb G, Welk E, Wirth C, Wu YT, Yang X, Zeng X, Zhang S, Zhou H, Ma K, Schmid B (2011) Community assembly during secondary forest succession in a Chinese subtropical forest. *Ecological Monographs* 81: 25–41. <https://doi.org/10.1890/09-2172.1>
- Corcos D, Lami F, Nardi D, Boscutti F, Sigura M, Giannone F, Pantini P, Tagliapietra A, Busato F, Sibella R, Marini L (2021) Cross-taxon congruence between predatory arthropods and plants across Mediterranean agricultural landscapes. *Ecological Indicators* 123: e107366. <https://doi.org/10.1016/j.ecolind.2021.107366>
- Duffy JE (2009) Why biodiversity is important to the functioning of real-world ecosystems. *Frontiers in Ecology and the Environment* 7: 437–444. <https://doi.org/10.1890/070195>
- Eichenberg D, Trogisch S, Huang YY, He JS, Bruelheide H (2015) Shifts in community leaf functional traits are related to litter decomposition along a secondary forest succession series in subtropical China. *Journal of Plant Ecology* 8: 401–410. <https://doi.org/10.1093/jpe/rtu021>
- Erwin TL (1979) Thoughts on the evolutionary history of ground beetles: hypotheses generated from comparative faunal analyses of lowland forest sites in temperate and tropical regions. In: Erwin TL, Ball GE, Whitehead DR, Halpern AL (Eds) *Carabid beetles: their evolution, natural history, and classification*. Junk Publishers, The Hague, 539–592. https://doi.org/10.1007/978-94-009-9628-1_30
- Erwin TL (1982a) *Agra*, arboreal beetles of neotropical forests: *erythropus* group systematics (Carabidae). *Systematic Entomology* 7: 39–71. <https://doi.org/10.1111/j.1365-3113.1982.tb00125.x>
- Erwin TL (1982b) Tropical forests: their richness in Coleoptera and other arthropod species. *The Coleopterists Bulletin* 36: 74–75. <http://europepmc.org/abstract/AGR/IND83098006>
- Erwin TL (1983) Tropical forest canopies: the last biotic frontier. *Bulletin of the Entomological Society of America* 29: 14–20. <https://doi.org/10.1093/besa/29.1.14>

- Erwin TL (2002) The beetle family Carabidae of Costa Rica: twenty-nine new species of *Agra* Fabricius 1801 (Coleoptera: Carabidae, Lebiini, Agrina). *Zootaxa* 119: 1–68. <https://doi.org/10.11646/zootaxa.119.1.1>
- Erwin TL (2004a) The beetle family Carabidae of Costa Rica: the genera of the Cryptobatida group of subtribe Agrina, tribe Lebiini, with new species and notes on their way of life (Insecta: Coleoptera). *Zootaxa* 662: 1–54. <https://doi.org/10.11646/zootaxa.662.1.1>
- Erwin TL (2004b) The beetle family Carabidae of Costa Rica and Panamá: descriptions of four new genera and six new species with notes on their way of life (Insecta: Coleoptera). *Zootaxa* 537: 1–18. <https://doi.org/10.11646/zootaxa.537.1.1>
- Erwin TL, Erwin LJM (1976) Relationships of predaceous beetles to tropical forest wood decay. Part II. The natural history of Neotropical *Eurycoleus macularis* Chevrolat (Carabidae: Lebiini) and its implications in the evolution of ectoparasitoidism. *Biotropica* 8: 215–224. <https://doi.org/10.2307/2989713>
- García-Tejero S, Spence JR, O'Halloran J, Bourassa S, Oxbrough A (2018) Natural succession and clearcutting as drivers of environmental heterogeneity and beta diversity in North American boreal forests. *PLoS ONE* 13: e0206931. <https://doi.org/10.1371/journal.pone.0206931>
- Gruner DS (2004) Attenuation of top-down and bottom-up forces in a complex terrestrial community. *Ecology* 85: 3010–3022. <https://doi.org/10.1890/04-0020>
- Haddad NM, Crutsinger GM, Gross K, Haarstad J, Knops JMH, Tilman D (2009) Plant species loss decreases arthropod diversity and shifts trophic structure. *Ecology Letters* 12: 1029–1039. <https://doi.org/10.1111/j.1461-0248.2009.01356.x>
- Harry I, Höfer H, Schielzeth H, Assmann T (2019) Protected habitats of Natura 2000 do not coincide with important diversity hotspots of arthropods in mountain grasslands. *Insect Conservation and Diversity* 12: 329–338. <https://doi.org/10.1111/icad.12349>
- Hartig F (2020) DHARMA: Residual diagnostics for hierarchical (multi-level/mixed) regression models. R package version 0.3.2.0. <https://CRAN.R-project.org/package=DHARMA>
- Heliölä J, Koivula M, Niemelä J (2001) Distribution of carabid beetles (Coleoptera: Carabidae) across a boreal forest–clearcut ecotone. *Conservation Biology* 15: 370–377. <https://doi.org/10.1046/j.1523-1739.2001.015002370.x>
- Homburg K, Drees C, Boutaud E, Nolte D, Schuett W, Zumstein P, von Ruschkowski E, Assmann T (2019) Where have all the beetles gone? Long-term study reveals carabid species decline in a nature reserve in Northern Germany. *Insect Conservation and Diversity* 12: 268–277. <https://doi.org/10.1111/icad.12348>
- Hu Z, Yu M (2008) Study on successions sequence of evergreen broad-leaved forest in Gutian Mountain of Zhejiang, Eastern China: species diversity. *Frontiers of Biology in China* 3: 45–49. <https://doi.org/10.1007/s11515-008-0011-4>
- Hülsmann M, Boutaud E, Buse J, Schuldt A, Assmann T (2019) Land-use legacy and tree age in continuous woodlands: weak effects on overall ground beetle assemblages, but strong effects on two threatened species. *Journal of Insect Conservation* 23: 623–633. <https://doi.org/10.1007/s10841-019-00156-8>
- Hunting W, Yang MM (2019) A taxonomic review of the pericaline ground-beetles in Taiwan, with descriptions of new species (Coleoptera: Carabidae, Lebiini). *ZooKeys* 816: 1–164. <https://doi.org/10.3897/zookeys.816.29738>

- Jelaska LS, Dumbovic V, Kucinic M (2011) Carabid beetle diversity and mean individual biomass in beech forests of various ages. *ZooKeys* 100: 393–405. <https://doi.org/10.3897/zookeys.100.1536>
- Jouveau S, Toïgo M, Giffard B, Castagneyrol B, van Halder I, Vétillard F, Jactel H (2020) Carabid activity-density increases with forest vegetation diversity at different spatial scales. *Insect Conservation and Diversity* 13: 36–46. <https://doi.org/10.1111/icad.12372>
- Koivula M (2011) Useful model organisms, indicators, or both? Ground beetles (Coleoptera: Carabidae) reflecting environmental conditions. *ZooKeys* 100: 287–317. <https://doi.org/10.3897/zookeys.100.1533>
- Koivula M, Kukkonen J, Niemelä J (2002) Boreal carabid-beetle (Coleoptera: Carabidae) assemblages along the clear-cut originated succession gradient. *Biodiversity and Conservation* 11: 1269–1288. <https://doi.org/10.1023/A:1016018702894>
- Korboulewsky N, Perez G, Chauvat M (2016) How tree diversity affects soil fauna diversity: a review. *Soil Biology and Biochemistry* 94: 94–106. <https://doi.org/10.1016/j.soilbio.2015.11.024>
- Krogerus R (1939) Zur Ökologie nordischer Moortiere. *Verhandlungen des 7. Internationalen Kongress für Entomologie Berlin 1938* 2: 1213–1231.
- Krogerus R (1960) Ökologische Studien über nordische Moorarthropoden. *Societas Scientiarum Fennica Commentationes Biologicae* 21: 1–238.
- Lang AC, Härdtle W, Bruelheide H, Kröber W, Schröter M, von Wehrden H, von Oheimb G (2012) Horizontal, but not vertical canopy structure is related to stand functional diversity in a subtropical slope forest. *Ecological Research* 27: 181–189. <https://doi.org/10.1007/s11284-011-0887-3>
- Lefcheck JS, Byrnes JEK, Isbell F, Gamfeldt L, Griffin JN, Eisenhauer N, Hensel MJS, Hector A, Cardinale BJ, Duffy JE (2015) Biodiversity enhances ecosystem multifunctionality across trophic levels and habitats. *Nature Communications* 6: e6936. <https://doi.org/10.1038/ncomms7936>
- Legendre P, Mi X, Ren H, Ma K, Yu M, Sun IF, He F (2009) Partitioning beta diversity in a subtropical broad-leaved forest of China. *Ecology* 90: 663–674. <https://doi.org/10.1890/07-1880.1>
- Liu JL, Li FR, Sun TS, Ma LF, Liu LL, Yang K (2016) Interactive effects of vegetation and soil determine the composition and diversity of carabid and tenebrionid functional groups in an arid ecosystem. *Journal of Arid Environments* 128: 80–90. <https://doi.org/10.1016/j.jaridenv.2016.01.009>
- Liu WY, Fox JED, Xu ZF (2003) Litterfall and nutrient dynamics in a montane moist evergreen broad-leaved forest in Ailao Mountains, SW China. *Plant Ecology* 164: 157–170. <https://doi.org/10.1023/A:1021201012950>
- Löbl I, Löbl D [Eds] (2017) *Catalogue of Palaearctic Coleoptera. Archostemata-Myxophaga-Adephaga, Revised and Updated Edition. Volume 1*. Brill, Leiden, 1443 pp.
- Loreau M, Naeem S, Inchausti P, Bengtsson J, Grime JP, Hector A, Hooper DU, Huston MA, Raffaelli D, Schmid B, Tilman D, Wardle DA (2001) Biodiversity and Ecosystem Functioning: current knowledge and future challenges. *Science* 294: 804–808. <https://doi.org/10.1126/science.1064088>

- Magura T, Tóthmérész B, Molnár T (2001) Forest edge and diversity: carabids along forest-grassland transects. *Biodiversity and Conservation* 10: 287–300. <https://doi.org/10.1023/A:1008967230493>
- Marrec R, Le Roux V, Martin L, Lenoir J, Brunet J, Cousins SAO, De Smedt P, Deconchat M, Diekmann M, Ehrmann S, Gallet-Moron E, Giffard B, Liira J, Lindgren J, Valdes A, Verheyen K, Wulf M, Decocq G (2021) Multiscale drivers of carabid beetle (Coleoptera: Carabidae) assemblages in small European woodlands. *Global Ecology and Biogeography* 30: 165–182. <https://doi.org/10.1111/geb.13208>
- Matern A, Drees C, Kleinwächter M, Assmann T (2007) Habitat modelling for the conservation of the rare ground beetle species *Carabus variolosus* (Coleoptera: Carabidae) in the riparian zones of headwaters. *Biological Conservation* 136: 618–627. <https://doi.org/10.1016/j.biocon.2007.01.006>
- Maveety S, Browne R, Erwin TL (2011) Carabidae diversity along an altitudinal gradient in a Peruvian cloud forest (Coleoptera). *ZooKeys* 147: 651–666. <https://doi.org/10.3897/zookeys.147.2047>
- Meyer P (2008) Distribution géographique des sous-genres actualisés du genre *Carabus*. Magelanes, Conflans-Sainte-Honorine, 106 pp.
- Mossakowski D (1970) Ökologische Untersuchungen an epigäischen Coleopteren atlantischer Moor- und Heidestandorte. *Zeitschrift für wissenschaftliche Zoologie* 181: 233–316.
- Mottl O, Fibich P, Klimes P, Volf M, Tropek R, Anderson-Teixeira K, Auga J, Blair T, Butterill P, Carscallen G, Gonzalez-Akre E, Goodman A, Kaman O, Lamarre GPA, Libra M, Losada ME, Manumbor M, Miller SE, Molem K, Nichols G, Plowman NS, Redmond C, Seifert CL, Vrana J, Weiblen GD, Novotny V (2020) Spatial covariance of herbivorous and predatory guilds of forest canopy arthropods along a latitudinal gradient. *Ecology Letters* 23: 1499–1510. <https://doi.org/10.1111/ele.13579>
- Nolte D, Schuldt A, Gossner MM, Ulrich W, Assmann T (2017) Functional traits drive ground beetle community structures in Central European forests: implications for conservation. *Biological Conservation* 213: 5–12. <https://doi.org/10.1016/j.biocon.2017.06.038>
- O'Brien MJ, Brezzi M, Schuldt A, Zhang JY, Ma KP, Schmid B, Niklaus PA (2016) Tree diversity drives diversity of arthropod herbivores, but successional stage mediates detritivores. *Ecology and Evolution* 7: 8753–8760. <https://doi.org/10.1002/ece3.3411>
- Oliver I, Beattie AJ (1996) Designing a cost-effective invertebrate survey: a test of methods for rapid assessment of biodiversity. *Ecological Applications* 6: 594–607. <https://doi.org/10.2307/2269394>
- Paje F, Mossakowski D (1984) pH-preferences and habitat selection in carabid beetles. *Oecologia* 64: 41–46. <https://doi.org/10.1007/BF00377541>
- Qodri A, Raffiudin R, Noerdjito W (2016) Diversity and abundance of Carabidae and Staphylinidae (Insecta: Coleoptera) in four montane habitat types on Mt. Bawakaraeng, South Sulawesi. *HAYATI Journal of Biosciences* 23: 22–28. <https://doi.org/10.1016/j.hjb.2016.04.002>
- Rainio J, Niemelä J (2003) Ground beetles (Coleoptera: Carabidae) as bioindicators. *Biodiversity and Conservation* 12: 487–506. <https://doi.org/10.1023/A:1022412617568>
- Riihimäki J, Kaitaniemi P, Koricheva J, Vehviläinen H (2005) Testing the enemies hypothesis in forest stands: the important role of tree species composition. *Oecologia* 142: 90–97. <https://doi.org/10.1007/s00442-004-1696-y>

- Root RB (1973) Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (*Brassica oleracea*). Ecological Monographs 43: 95–124. <https://doi.org/10.2307/1942161>
- Schuldt A, Baruffol M, Böhnke M, Bruelheide H, Härdtle W, Lang AC, Nadrowski K, von Oheimb G, Voigt W, Zhou H, Assmann T (2010) Tree diversity promotes insect herbivory in subtropical forests of south-east China. Journal of Ecology 98: 917–926. <https://doi.org/10.1111/j.1365-2745.2010.01659.x>
- Schuldt A, Bruelheide H, Buscot F, Assmann T, Erfmeier A, Klein AM, Ma K, Scholten T, Staab M, Wirth C, Zhang J, Wubet T (2017) Belowground top-down and aboveground bottom-up effects structure multitrophic community relationships in a biodiverse forest. Scientific Reports 7: e4222. <https://doi.org/10.1038/s41598-017-04619-3>
- Schuldt A, Wubet T, Buscot F, Staab M, Assmann T, Böhnke-Kammerlander M, Both S, Erfmeier A, Klein AM, Ma KP, Pietsch KA, Schultze S, Wirth C, Zhang JY, Zumstein P, Bruelheide H (2015a) Multitrophic diversity in a biodiverse forest is highly nonlinear across spatial scales. Nature Communications 6: e10169. <https://doi.org/10.1038/ncomms10169>
- Schuldt A, Ebeling A, Kunz M, Staab M, Guimarães-Steinicke C, Bachmann D, Buchmann N, Durka W, Fichtner A, Fornoff F, Härdtle W, Hertzog LR, Klein AM, Roscher C, Schaller J, von Oheimb G, Weigelt A, Weisser W, Wirth C, Zhang J, Bruelheide H, Eisenhauer N (2019) Multiple plant diversity components drive consumer communities across ecosystems. Nature Communications 10: e1460. <https://doi.org/10.1038/s41467-019-09448-8>
- Schuldt A, Both S, Bruelheide H, Härdtle W, Schmid B, Zhou H, Assmann T (2011) Predator diversity and abundance provide little support for the enemies hypothesis in forests of high tree diversity. PLoS ONE 6: e22905. <https://doi.org/10.1371/journal.pone.0022905>
- Schuldt A, Bruelheide H, Härdtle W, Assmann T, Li Y, Ma KP, von Oheimb G, Zhang JY (2015b) Early positive effects of tree species richness on herbivory in a large-scale forest biodiversity experiment influence tree growth. Journal of Ecology 103: 563–571. <https://doi.org/10.1111/1365-2745.12396>
- Schwerk A, Szyszko J (2007) Mittlere Individuen Biomasse (MIB) von epigäischen Laufkäfern (Coleoptera: Carabidae) als Indikator für Sukzessionsstadien. Angewandte Carabidologie 8: 69–72.
- Shook G, Wu XJ (2007) Tiger beetles of Yunnan. Yunnan Publishing Group, Yunnan Science and Technology Press, Kunming, 119 pp.
- Staab M, Liu X, Assmann T, Bruelheide H, Buscot F, Durka W, Erfmeier A, Klein AM, Ma K, Michalski S, Wubet T, Schmid B, Schuldt A (2021) Tree phylogenetic diversity structures multitrophic communities. Functional Ecology 35: 521–534. <https://doi.org/10.1111/1365-2435.13722>
- Staab M, Schuldt A (2020) The influence of tree diversity on natural enemies – a review of the “Enemies” hypothesis in forests. Current Forestry Reports 6: 243–259. <https://doi.org/10.1007/s40725-020-00123-6>
- Staab M, Schuldt A, Assmann T, Klein AM (2014) Tree diversity promotes predator but not omnivore ants in a subtropical Chinese forest. Ecological Entomology 39: 637–647. <https://doi.org/10.1111/een.12143>
- Stork NE (1980) A scanning electron microscope study of tarsal adhesive setae in the Coleoptera. Zoological Journal of the Linnean Society 68: 173–306. <https://doi.org/10.1111/j.1096-3642.1980.tb01121.x>

- Stork NE (2018) How many species of insects and other terrestrial arthropods are there on earth? *Annual Review of Entomology* 63: 31–45. <https://doi.org/10.1146/annurev-ento-020117-043348>
- Stork NE, Grimbacher PS (2006) Beetle assemblages from an Australian tropical rainforest show that the canopy and the ground strata contribute equally to biodiversity. *Proceedings of the Royal Society B-Biological Sciences* 273: 1969–1975. <https://doi.org/10.1098/rspb.2006.3521>
- Strong DR, Lawton JH, Southwood TRE (1984) *Insects on plants: community patterns and mechanisms*. Blackwell, Oxford, 313 pp.
- Taboada A, Kotze DJ, Tarrega R, Salgado JM (2008) Carabids of differently aged reforested pinewoods and a natural pine forest in a historically modified landscape. *Basic and Applied Ecology* 9: 161–171. <https://doi.org/10.1016/j.baae.2007.01.004>
- Taboada A, Tarrega R, Calvo L, Marcos E, Marcos JA, Salgado JM (2010) Plant and carabid beetle species diversity in relation to forest type and structural heterogeneity. *European Journal of Forest Research* 129: 31–45. <https://doi.org/10.1007/s10342-008-0245-3>
- Tang CQ (2006) Forest vegetation as related to climate and soil conditions at varying altitudes on a humid subtropical mountain, Mount Emei, Sichuan, China. *Ecological Research* 21: 174–180. <https://doi.org/10.1007/s11284-005-0106-1>
- Terrell-Nield C (1990) Is it possible to age woodlands on the basis of their carabid beetle diversity? *Entomologist* 109: 136–145.
- Thiele HU (1977) *Carabid beetles in their environments: a study on habitat selection by adaptations in Physiology and Behaviour*. Springer-Verlag Berlin Heidelberg, 372 pp. <https://doi.org/10.1007/978-3-642-81154-8>
- Thorn S, Bußler H, Fritze MA, Goeder P, Müller J, Weiß I, Seibold S (2016) Canopy closure determines arthropod assemblages in microhabitats created by windstorms and salvage logging. *Forest Ecology and Management* 381: 188–195. <https://doi.org/10.1016/j.foreco.2016.09.029>
- Vehviläinen H, Koricheva J, Ruohomäki K (2008) Effects of stand tree species composition and diversity on abundance of predatory arthropods. *Oikos* 117: 935–943. <https://doi.org/10.1111/j.0030-1299.2008.15972.x>
- Vennila S (2000) Pitfall trap sampling of tropical carabids (Carabidae: Coleoptera) – evaluation of traps, preservatives and sampling frequency. *Journal of Bombay Natural History Society* 97: 241–246.
- Vennila S (1999) Optimum sampling effort for study of tropical ground beetles (Carabidae: Coleoptera) using pitfall traps. *Current Science* 77: 281–283.
- Wang MQ, Li Y, Chesters D, Anttonen P, Bruehlheide H, Chen JT, Durka W, Guo PF, Härdtle W, Ma KP, Michalski SG, Schmid B, von Oheimb G, Wu CS, Zhang NL, Zhou QS, Schuldt A, Zhu CD (2019) Multiple components of plant diversity loss determine herbivore phylogenetic diversity in a subtropical forest experiment. *Journal of Ecology* 107: 2697–2712. <https://doi.org/10.1111/1365-2745.13273>
- Whitehead DR, Ball GE (1997) The middle American genus *Onypterygia* Dejean (Insecta: Coleoptera, Carabidae, Platynini): a taxonomic revision of the species, with notes about their way of life and geographical distribution. *Annals of the Carnegie Museum* 66: 289–409. <https://www.biodiversitylibrary.org/part/215140>.

- Wolters V, Bengtsson J, Zaitsev AS (2006) Relationship among the species richness of different taxa. *Ecology* 87: 1886–1895. [https://doi.org/10.1890/0012-9658\(2006\)87\[1886:RATSRO\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1886:RATSRO]2.0.CO;2)
- Work T, Shorthouse D, Spence J, Volney W, Langor D (2011) Stand composition and structure of the boreal mixedwood and epigaeic arthropods of the Ecosystem Management Emulating Natural Disturbance (EMEND) landbase in northwestern Alberta. *Canadian Journal of Forest Research* 34: 417–430. <https://doi.org/10.1139/x03-238>
- Worthen WB, Merriman DCG (2013) Relationships between carabid beetle communities and forest stand parameters: Taxon congruence or habitat association? *Southeastern Naturalist* 12: 379–386. <https://doi.org/10.1656/058.012.0211>
- Wu YT, Wubet T, Trogisch S, Both S, Scholten T, Bruelheide H, Buscot F (2013) Forest age and plant species composition determine the soil fungal community composition in a Chinese subtropical forest. *PLoS ONE* 8: e66829. <https://doi.org/10.1371/journal.pone.0066829>
- Yeeles P, Lach L, Hobbs RJ, van Wees M, Didham RK (2017) Woody plant richness does not influence invertebrate community reassembly trajectories in a tree diversity experiment. *Ecology* 98: 500–511. <https://doi.org/10.1002/ecy.1662>
- Yu M, Hu Z, Yu J, Ding B, Fang T (2001) Forest vegetation types in Gutianshan Natural Reserve in Zhejiang. *Journal of Zhejiang University (Agriculture and Life Sciences)* 27: 375–380. <http://europemc.org/abstract/CBA/351032>
- Yu XD, Luo TH, Zhou HZ (2009) Distribution of carabid beetles (Coleoptera: Carabidae) across ecotones between regenerating and mature forests in southwestern China. *Environmental Entomology* 38: 1053–1060. <https://doi.org/10.1603/022.038.0412>
- Yu XD, Liu CL, Lu L, Bearer SL, Luo TH, Zhou HZ (2017) Does selective logging change ground-dwelling beetle assemblages in a subtropical broad-leaved forest of China? *Insect Science* 24: 303–313. <https://doi.org/10.1111/1744-7917.12283>
- Zou Y, Sang W, Bai F, Brennan E, Dickman M, Lius Y, Li L, Marples A, Shi H, Sui Z, Sun X, Wang C, Wang X, Warren-Thomas E, Yang X, Yu Z, Axmacher JC (2019) Large-scale alpha-diversity patterns in plants and ground beetles (Coleoptera: Carabidae) indicate a high biodiversity conservation value of China's restored temperate forest landscapes. *Diversity and Distributions* 25: 1613–1624. <https://doi.org/10.1111/ddi.12964>
- Zou Y, Sang WG, Bai F, Axmacher JC (2013) Relationships between plant diversity and the abundance and alpha-diversity of predatory ground beetles (Coleoptera: Carabidae) in a mature Asian temperate forest ecosystem. *PLoS ONE* 8: e82792. <https://doi.org/10.1371/journal.pone.0082792>
- Zou Y, Sang W, Wang S, Warren-Thomas E, Liu Y, Yu Z, Wang C, Axmacher JC (2015) Diversity patterns of ground beetles and understory vegetation in mature, secondary, and plantation forest regions of temperate northern China. *Ecology and Evolution* 5: 531–542. <https://doi.org/10.1002/ece3.1367>
- Zuur A, Ieno EN, Walker N, Saveliev A, Smith GM (2009) Mixed effects models and extensions in ecology with R. Springer-Verlag, New York, 574 pp. https://doi.org/10.1007/978-0-387-87458-6_1