



Editorial

Jacobson, Nicholas C.; Funk, Burkhardt; Abdullah, Saeed

Published in:
Frontiers in Psychology

DOI:
[10.3389/fpsyg.2022.1126971](https://doi.org/10.3389/fpsyg.2022.1126971)

Publication date:
2023

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

Citation for published version (APA):
Jacobson, N. C., Funk, B., & Abdullah, S. (2023). Editorial: Quantitative modeling of psychopathology using passively collected data. *Frontiers in Psychology*, 13, Article 1126971.
<https://doi.org/10.3389/fpsyg.2022.1126971>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



OPEN ACCESS

EDITED AND REVIEWED BY
Stefano Barlati,
ASST Spedali Civili of Brescia, Italy

*CORRESPONDENCE
Nicholas C. Jacobson
✉ Nicholas.C.Jacobson@dartmouth.edu

SPECIALTY SECTION
This article was submitted to
Psychopathology,
a section of the journal
Frontiers in Psychology

RECEIVED 18 December 2022
ACCEPTED 19 December 2022
PUBLISHED 24 January 2023

CITATION
Jacobson NC, Funk B and Abdullah S
(2023) Editorial: Quantitative modeling
of psychopathology using passively
collected data.
Front. Psychol. 13:1126971.
doi: 10.3389/fpsyg.2022.1126971

COPYRIGHT
© 2023 Jacobson, Funk and Abdullah.
This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Editorial: Quantitative modeling of psychopathology using passively collected data

Nicholas C. Jacobson^{1*}, Burkhardt Funk² and Saeed Abdullah³

¹Center for Technology and Behavioral Health, Departments of Biomedical Data Science, Psychiatry, and Computer Science, Geisel School of Medicine, Dartmouth College, Hanover, NH, United States, ²Leuphana University Lüneburg, Lüneburg, Germany, ³College of Information Sciences and Technology, Penn State University, State College, PA, United States

KEYWORDS

digital phenotyping, digital biomarker, mobile sensing, machine learning, passive sensing

Editorial on the Research Topic

Quantitative modeling of psychopathology using passively collected data

Traditional mental health assessment collects data at a single point in time, which does not provide a comprehensive picture of an individual's mental health over an extended period. To address this issue, methods have been developed that involve repeatedly asking people to fill out surveys. However, these methods can be burdensome and as such, might not support granular assessment. Advances in technology have made it possible to collect a wide range of behavioral, social, and physiological data about an individual without requiring their active participation. This data can be collected from various sources, including smartphone and wearable sensors, electronic health records, social media data, and internet search data, with minimal burden to the individual. As a result, this data can provide a more comprehensive and nuanced understanding of an individual's mental health over time.

For instance, smartphone sensors can collect data on a person's social interactions, physical activity, and environment to gain insight into how these factors may affect their mental health. This data can be used to study the relationship between mental health measures and outcomes, develop predictive models, and test theoretical models of mental health. For instance, data from smartphone sensors could be used to explore the link between social isolation and changes in depression symptoms.

The goal of this Research Topic is to explore how passively collected data from sources like smartphones and wearable sensors can be used in combination with quantitative methods to improve our understanding and prediction of mental health. The manuscripts in this collection show how these data can be easily collected with minimal burden and can be used to gain new insights into mental health and to predict changes in mental health conditions.

Asgari et al. have developed a method for automatically detecting prosodic abnormalities in speech that are associated with autism. Using a machine learning model trained on a dataset of speech samples from individuals with autism and typically developing controls, they were able to accurately distinguish between the two groups. Their approach could be used as a new measure in autism treatment research and for early detection of the condition in preverbal infants or toddlers who are at risk.

Ponzo et al. have developed a new interoceptive task, the CARDiac Elevation Detection (CARED) task, which aims to improve upon existing methods. The task involves using a wearable device to record participants' heart rate and sending notifications to their mobile devices over a period of 4 weeks. Participants were asked to report on their heart rate and rate their confidence in their responses. The study found that the task was easy to administer and that participants had good insight into their interoceptive abilities.

Ding et al. developed a machine learning model that can predict changes in state anxiety levels with high temporal resolution. The model was trained on data collected from participants who were induced to experience state anxiety through exposure to aversive stimuli. The data included dimensional emotion ratings, electrocardiogram readings, and galvanic skin response measurements. The model was able to accurately predict self-reported state anxiety levels, providing a sensitive and fine-grained measure of state anxiety that could be useful for future studies of affective brain-computer interaction and anxiety modulation.

Liu and Shi present a novel hybrid feature selection and ensemble approach for detecting depression on social media. The method uses a combination of recursive elimination and extremely randomized trees to select the optimal subset of features for a stacking ensemble model. Their proposed method achieved an accuracy of 90.27% in identifying individuals with depression using online interaction data, resulting in a better performance than recent machine learning algorithms. This approach has potential applications for developing new methods to identify depression on social media.

Lin et al. evaluated the feasibility and effectiveness of using a machine learning-based smartphone app, the Ellipsis Health App, for detecting depression and anxiety in a senior population. The app uses semantic information from recorded speech to screen for these conditions. The study data shows a high completion rate of the app and a good performance in detecting depression and anxiety among seniors and various age ranges. Following the study findings, the authors suggest that the Ellipsis Health App is a promising tool for mental health screening in a senior population.

Maatoug et al. review the potential use of digital phenotyping in the diagnosis of mood disorders. Digital phenotyping involves the use of real-time data collected from digital sensors, wearable devices, and smartphones to determine the digital signature

of a particular pathology. The study found that individuals with mood disorders often exhibit decreases in functional and biological parameters, such as decreased activities and walking, fewer calls and text messages, and lower body temperature and heart rate variability. These findings suggest that digital phenotyping could be a valuable addition to traditional clinical interviews in the diagnosis of mood disorders, providing objective data to supplement subjective symptoms.

The goal of this Research Topic is to examine how passively available data, such as data from smartphones and wearable sensors, can be used in combination with quantitative methods to improve our understanding and prediction of mental health. The manuscripts in this collection demonstrate how these data can be collected with minimal burden, used to enhance our knowledge of mental health, and/or used to predict mental health. They present a range of approaches, including using machine learning to detect prosodic abnormalities in speech, developing a wearable device for interoceptive research, and using multi-modal data to predict changes in anxiety levels. These approaches have the potential to improve the detection and monitoring of mental health conditions.

Author contributions

All authors listed have made a substantial, direct, and intellectual contribution to the work and approved it for publication.

Funding

This work was funded by the National Institute of Mental Health and the National Institute of General Medical Sciences under grant 1 R01 MH123482-01.

Conflict of interest

BF is a shareholder in HelloBetter.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.